Maths Explore-Notes- Terminal-12

Course notes

Amounts and changes

For a finite random variable X, taking the values x_i with probabilities p_i,

$$E(X) = \sum_i x_i p_i, \qquad V(X) = \sum_i (x_i - E(X))^2 p_i = E(X^2) - E(X)^2.$$

The standard deviation is $\sigma(X) = \sqrt{V(X)}$. Developing the square in $V(aX+b)$, the lag b seliminate and the factor a is raised squared. For finite random variables X, Y and Real a, b,

$$E(X+Y) = E(X) + E(Y), \quad E(aX+b) = aE(X) + b, \quad V(aX+b) = a^2 V(X).$$

The linearity of expectation does not require independence. X, Y are independent, $V(X+Y) = V(X) + V(Y)$ (admitted). Without this hypothesis, this formula cannot be applied.
$aX + b$ value $|a|\sigma(X)$.

Example of addiction. If $Y = X$, then $V(X+Y) = V(2X) = 4V(X)$, not 2 $V(X)$ as soon as $V(X) \neq 0$.

Demonstration for the binomial. Write $X = B_1 + \cdots + B_n$ with independent indicators of success. B_i checks $E(B_i) = p$, $E(B_i^2) = p$ and $V(B_i) = p - p^2$. By linearity and additivity under independence, $E(X) = np$, $V(X) = np(1-p)$.

Sample and mean

A size sample n of a law is a family of n variables of the same law.
have expectation μ and variance σ^2, we note $S_n = \sum_{i=1}^{n} X_i$ and $M_n = \frac{S_n}{n}$.

Sum and average

$$E(S_n) = n\mu, \quad V(S_n) = n\sigma^2, \qquad E(M_n) = \mu, \quad V(M_n) = \frac{\sigma^2}{n}, \quad \sigma(M_n) = \frac{\sigma}{\sqrt{n}}.$$

To divide the standard deviation of the average by 2, multiply the size by 4. The variance of the sum increase; the average is decreasing.

Concentration

Inequality of Bienaymé-Chebychev

For a variable of expectation μ, variance σ^2, and $\delta > 0$,

$$P(|X - \mu| \geq \delta) \leq \frac{\sigma^2}{\delta^2}.$$

A limit of more than 1 remains true, but does not provide additional information.
may be replaced by 1.

Proof in the finished frame. In $V(X) = \sum_x (x-\mu)^2 P(X = x)$, only keep the values checking $|x - \mu| \geq \delta$. Each square is then at least δ^2. So $V(X) \geq \delta^2 P(|X - \mu| \geq \delta)$.

Law of large numbers

Applied to the average of a sample,

$$P(|M_n - \mu| \geq \delta) \leq \frac{\sigma^2}{n\delta^2} \xrightarrow[n\to\infty]{} 0.$$

This is the law of the large numbers in the given framework: for a specified accuracy, the probability of at least equal to this precision tends to zero. This result does not say that each achievement is close to μ in a monotonous manner.

Select a sufficient size

To guarantee $P(|M_n - \mu| < \delta) \geq 1 - \alpha$, just choose

$$n \geq \frac{\sigma^2}{\alpha\delta^2}.$$

Take the ceiling. It is the smallest satisfying integer this bound , not necessarily of minimum size for the exact probability.

For a frequency of success, $\mu = p$ and $\sigma^2 = p(1-p) \leq \frac{1}{4}$. If p is unknown, $n \geq \frac{1}{4\alpha\delta^2}$ is a guarantee uniform.

From individual law to average

14.1 Describe a non-binomial variable

First use: S62.

Example resolved · A repair cost

A cost X in euros takes values 0, 20 and 50 with probabilities 0 ,5, 0,3 and 0,2. $E(X) = 16$, $E(X^2) = 620$, therefore $V(X) = 620 - 16^2 = 364$ in square euros. $\sqrt{364} \simeq 19{,}08$ Euros. Average 16 is none of the possible values of X; it describes an average theoretical on many achievements.

A1 · Immediate application X takes values 1, 2, 4 with probabilities 1 /2, 1/4, 1/4. Check the Law and calculate expectation, variance and standard deviation.

14.2 Remission also changes dispersion

First use: S62.

Property · Affine processing

For $Y = aX + b$, $E(Y) = aE(X) + b$ and $V(Y) = a^2V(X)$. The standard deviation is $|a|\sigma(X)$. Adding a constant translates all values without changing their dispersion.

Example resolved · Cost after remission

For the previous cost, a discount of 20% and then 5 euros of costs gives $Y = 0{,}8X + 5$. Thus $E(Y) = 17.8$ and $V(Y) = 0{,}64 \times 364 = 232.96$. The variance is neither $0.8V(X) + 5$ or $0.8\,V(X)$.

A2 · Partial guidance For the law of the previous application, calculate E and V of $Y = 3X - 2$ and then $Z = -X$. Explain why multiply by -1 does not change the variance.

14.3 Independence intervenes in the variance

First use: S64.

Property · Sum and average

For variables $X_1, \dots, X_n$ of the same expectation μ and even variance σ^2, average M_n a for expectation μ. If independent, $V(M_n) = \sigma^2/n$ and $\sigma(M_n) = \sigma/\sqrt{n}$. Linearity expectation does not require independence.

Demonstration. With $S_n = \sum X_i$, linearity gives $E(S_n) = n\mu$. Under independence, additivity Allowed variances give $V(S_n) = n\sigma^2$. Multiply by 1 /n for the average multiply expectation by 1 /n and variance by 1 /n^2. For Bernoulli, E=p and $V = p(1-p)$; the binomial, sum of n Bernoulli independent, therefore has E=np and $V = np(1-p)$. □

Example resolved · Average of 100 costs

Per cent independent costs of the same law, of expectation 16 and of variance 364, the total has for expectation 1600 and variance 36400. The average cost has for expectation 16 and variance 3.64, hence a Standard deviation of around EUR 1.91.

A3 · Counterexample X is 0 or 1 with probability 1 /2 each. Compare variances of $(X_1 + \cdots + X_n)/n$ for independent copies, then for $X_1 = \cdots = X_n = X$.

14.4 Sufficient guarantee, not an exact probability

First use: S73.

Property · Bienaymé-Chebychev

For $\delta > 0$, $P(|X - \mu| \geq \delta) \leq V(X)/\delta^2$. For an independent average, $P(|M_n - \mu| \geq \delta) \leq \sigma^2/(n\delta^2)$. Any bound may be reduced to 1 if it exceeds it.

Example resolved · Size guaranteed by the bound

For a probability success frequency 1 /2, $V(M_n) = 1/(4n)$. We want a strict deviation less than 0 ,1 with a probability at least 0 ,95. It is sufficient to have $25/n \leq 0{,}05$, so $n \geq 500$. This gives the smallest integer satisfying this sufficient inequality; it does not prove that 499 observations actually fail.

A4 · Immediate application A variable has a variance of 9. independent guarantees, by this bound, an average deviation strictly less than 0 ,5 with one Like at least 0.99?

A5 · Compare two informations For $X \sim \mathcal{B}(10; 1/2)$ and $M = X/10$, calculate exactly $P(|M - 1/2| \geq 0{,}3)$ and compare to the Tchebychev bound. Work with the exact entire bounds.

14.5 Several samples, several averages

First use: S74 and S77.




Figure 14.1. Theoretical curve 1/ $(2\sqrt{n})$: multiply n by 4 divides standard deviation by 2, not by 4.

Example resolved · What is compared

With n=100, the theoretical standard deviation of a probability-success frequency 1/2 is 0.05.
series of 1000 experiments of 100 tests produces 1000 averages.
a series of averages compared to 0.05; the 100 000 individual results have another dispersion.

A6 · TP and interpretation With the program provided, simulate 1,000 samples independent of sizes 10,100 and 1000 for a Bernoulli parameter 1 /2. For each size, compare the standard deviation averages to 1 /(2 $\sqrt{n}$) and count deviations of at least 0 ,1. Why a single trajectory Does it not answer this question?

A7 · Refuter "The law of large numbers says the frequency is approaching expectation give a short sequence of results that temporarily removes the frequency of 1/2, then reformulate the result correctly.

Wait a minute at the courthouse. Make a sentence containing "accuracy", "risk" and "sufficient size", then distinguish this sentence from certainty for a particular sample.

Delayed review · S90. In a topic of synthesis, identify first the law or variance of the average, before choosing the threshold.

Research activity

Fluctuations of frequency · 40 min.

We independently launch a balanced piece. $n \geq 1$, note F_n the frequency of · pile » among the n first shots.

1. Write a program that simulates 1000 throws, then represent points $(n; F_n)$ for $1 \leq n \leq 1000$. The gap to $\frac{1}{2}$ Does it necessarily decrease at each pitch?
2. Express $E(F_n)$ and $V(F_n)$ according to the n.
3. With the help of Bienaymé-Chebychev, increase $P(|F_{1000} - \frac{1}{2}| \geq 0{,}05)$.
4. Determine an Integer n sufficient to ensure $P(|F_n - \frac{1}{2}| < 0{,}05) \geq 0{,}99$.
5. Explain what the simulated trajectory shows and what the inequality achieved guarantees. Is it proof of this guarantee?