
1.1 Defining and representing a sequence

Key point · Definition and results. A real sequence is a function whose variable is an integer: to each allowed index n corresponds a real number u_n. The notation (u_n) denotes the sequence, u_n a term. An explicit formula gives u_n directly in terms of n. A definition by recurrence gives a first term and a rule for passing from one term to the next. The representation is a set of points (n, u_n): these points are not necessarily joined up.

Status and justification. Definition. The uniqueness of the terms obtained from a first term and a deterministic rule is accepted; no formal proof by induction is required.

Example 1. For $u_n = n^2 + 1$, $u_0 = 1$, $u_3 = 10$. For $v_0 = 1$ and $v_{n+1} = 2v_n + 1$, $v_1 = 3$, $v_2 = 7$, $v_3 = 15$. We cannot replace v_3 by $2 \times 3 + 1$: the rule applies to the previous term.



Terms of $u_n = 2n + 2$: only integer indices are plotted.

Watch out. The index is not the value. A finite list of terms does not determine a unique rule.

Application. Calculate the first four terms of $w_0 = 2$, $w_{n+1} = w_n + 2n + 1$; suggest an explicit formula.

1.2 Arithmetic sequences

Key point · Definition and results. A sequence is arithmetic with common difference r if $u_{n+1} = u_n + r$ for every allowed index. Then $u_n = u_p + (n - p)r$ for $n \geq p$. If $r > 0$ it is strictly increasing; if $r < 0$, strictly decreasing; if $r = 0$, constant. The points lie on a straight line.

Status and justification. For fixed p, the sequence $v_n = u_p + (n - p)r$ satisfies $v_p = u_p$ and $v_{n+1} = v_n + r$. By uniqueness of the successive terms, $u_n = v_n$. Monotonicity follows from the sign of $u_{n+1} - u_n = r$.

Example 2. A tank holds 140 L and receives 12 L per minute. After n minutes, $u_n = 140 + 12n$. At 5 min: 200 L. If we start the indexing at 1 with 140 L, we write $u_n = 140 + 12(n - 1)$.

Watch out. Adding a constant percentage does not generally give an arithmetic sequence.

Application. We know $u_3 = 17$ and $u_8 = 37$ for an arithmetic sequence. Determine r and u_0.

1.3 Geometric sequences

Key point · Definition and results. A sequence is geometric with common ratio q if $u_{n+1} = qu_n$. For $n \geq p$, $u_n = u_p q^{n-p}$, with the convention that an empty product equals 1. A constant rate t gives $q = 1 + t$. Monotonicity is determined using $u_{n+1} - u_n = (q-1)u_n$. For $q < 0$, the signs may alternate.

Status and justification. The formula gives the first term, and multiplying it by q gives the next term. The uniqueness of the generation justifies the formula. For $q = 0$, all terms after the first are zero; treat this case directly.

Example 3. A quantity of 160 mg loses 25% at each step: $u_n = 160(\frac{3}{4})^n$. The terms are positive and $u_{n+1} - u_n = -\frac{u_n}{4} < 0$. We obtain $u_2 = 90$ mg, without confusing a 25% decrease with removing 25 mg.

Watch out. $q > 1$ is not enough to conclude that every geometric sequence is increasing: the sign of the first term matters.

Application. Study the monotonicity of $u_n = -2 \times 3^n$, then of $v_n = 2(-1)^n$.

1.4 Sums, indices and proofs

Key point · Definition and results. For $n \geq 1$, $1 + 2 + \cdots + n = \frac{n(n+1)}{2}$. For $q \neq 1$ and $n \geq 0$, $1 + q + \cdots + q^n = \frac{1 - q^{n+1}}{1 - q}$. For $q = 1$, the sum equals $n + 1$. Hence, from u_p to u_n, an arithmetic sum equals $\frac{(n - p + 1)(u_p + u_n)}{2}$; a geometric sum equals $\frac{u_p(1 - q^{n-p+1})}{1 - q}$ if $q \neq 1$.

Status and justification. Writing $S = 1 + \cdots + n$ in both directions, each pair in the sum $2S$ equals $n + 1$, and there are n pairs. For $T = 1 + q + \cdots + q^n$, subtract $qT = q + \cdots + q^{n+1}$: $(1 - q)T = 1 - q^{n+1}$. The case $q = 1$ must come before the division.

Example 4. For $u_n = 3 + 2n$, $u_2 + \cdots + u_6 = \frac{5(7 + 15)}{2} = 55$. For $v_n = 2 \times 3^n$, $v_0 + \cdots + v_3 = \frac{2(1 - 3^4)}{1 - 3} = 80$.

Watch out. Count the terms before using the formula: from 0 to n there are n+1.

Application. Calculate $\sum_{k=2}^{5} 2^k$ without the summation symbol in the answer.

1.5 Conjecturing a limit and programming a threshold

Key point · Definition and results. A limit is approached here intuitively. The terms $\frac{1}{n+1}$ get closer to 0; n^2 grows without bound; $(-1)^n$ does not settle. A numerical experiment suggests a conclusion but does not prove a behaviour for all indices. A threshold algorithm must specify the direction of the inequality and provide a stopping condition.

Status and justification. Illustrative observations, without a formal theory of limits. The correctness of an integer threshold is checked by examining the index found and the previous index, once monotonicity is established.

Example 5. For $u_n = 80(\frac{1}{2})^n$, the first index with $u_n < 6$ is 4: $u_3 = 10 \geq 6$ and $u_4 = 5 < 6$. The program starts with $n = 0, u = 80$ and repeats the division by 2 while $u \geq 6$.

Watch out. A loop with no bound may never stop if the threshold cannot be reached; $<$ and $\leq$ are not interchangeable.

Application. For $u_n = 48(\frac{1}{2})^n$, determine the first indices where $u_n \leq 6$ and where $u_n < 6$.