Unit no. ° 4

REAL NUMBERS AND INTERVALS


2-13 November 2026

I Sets of numbers

Definitions

- $\mathbb{N}$: natural numbers; $\mathbb{Z}$: integers.
- $\mathbb{D}$: decimal numbers, of the form $a/10^n$ with $a \in \mathbb{Z}$, $n \in \mathbb{N}$.
- $\mathbb{Q}$: rational numbers, of the form a/b with $a, b \in \mathbb{Z}$, $b \neq 0$.
- $\mathbb{R}$: real numbers, represented by the x-coordinates of the points of a number line. A real number not rational is **irrational** .

$$\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{D} \subsetneq \mathbb{Q} \subsetneq \mathbb{R}.$$



Example 4.1

Give the smallest of the five sets containing each of the numbers: -95 ; $7{,}29$; $5/3$; $\sqrt{25}$; $\sqrt{2}$; π ; $0{,}0002$.

Solution:

$-95 \in \mathbb{Z}$; $7{,}29 \in \mathbb{D}$; $5/3 \in \mathbb{Q}$; $\sqrt{25} = 5 \in \mathbb{N}$; $\sqrt{2}, \pi \in \mathbb{R} \setminus \mathbb{Q}$; $0{,}0002 = 2/10^4 \in \mathbb{D}$. The irrationality of π is assumed here.

II Two proofs by contradiction

Proof by contradiction

To prove a statement, assume its negation, deduce a contradiction, then conclude that the statement is true.

Property

$\frac{1}{3}$ is not a decimal number.

Write a proof.

Solution:

Suppose $1/3 = a/10^n$ with $a \in \mathbb{Z}$, $n \in \mathbb{N}$. Then $10^n = 3a$ would be a multiple of 3. Yet $10 \equiv 1$ modulo 3, so 10^n leaves a remainder of 1 on division by 3 (we can also use the test for divisibility by 3). Contradiction. So $1/3 \notin \mathbb{D}$.

Property

$\sqrt{2}$ is irrational.

Write a proof.

Solution:

Suppose $\sqrt{2} = p/q$ with $p, q \in \mathbb{N}$, $q \neq 0$, and p/q irreducible. Then $p^2 = 2q^2$, therefore p^2 is even. If p were odd, its square would be odd too: thus p is even. Let us write $p = 2k$. We get $4k^2 = 2q^2$, then $q^2 = 2k^2$; q is therefore even in turn. So 2 divides p and q, contradicting irreducibility. Hence $\sqrt{2} \notin \mathbb{Q}$.

Guidance for the teacher

Have students identify separately the starting hypothesis, the implication used through the contrapositive on parity and the final contradiction. Do not present a calculator decimal value as a proof of irrationality.

III Intervals of the real line

Notation

For two real numbers $a < b$, intervals describe the real numbers between two bounds, with or without the bounds. A closed bracket includes the bound, an open bracket excludes it. The symbols $-\infty, +\infty$ are not real numbers and are always accompanied by an open bracket.

Condition	Interval	Description
$a \leq x \leq b$	$[a; b]$	both bounds included
$a < x \leq b$	$]a; b]$	a excluded, b included
$a \leq x < b$	$[a; b[$	a included, b excluded
$a < x < b$	$]a; b[$	both bounds excluded
$x \geq a$	$[a; +\infty[$	greater than or equal to a
$x > a$	$]a; +\infty[$	strictly greater than a
$x \leq b$	$]-\infty; b]$	less than or equal to b
$x < b$	$]-\infty; b[$	strictly less than b

$$\mathbb{R} =]-\infty; +\infty[, \quad \mathbb{R}^* =]-\infty; 0[\cup]0; +\infty[.$$

Watch out. The strictly negative real numbers form $]-\infty; 0[$: 0 is excluded. $[a; a] = \{a\}$; $]a; a[= \varnothing$.

Example 4.2

Write as an interval and then represent: $2 \leq x \leq 5$; $x > 4$; $-6 < x \leq 2$; $x < 0$.

Solution:

$[2; 5]$; $]4; +\infty[$; $]-6; 2]$; $]-\infty; 0[$. On the number line, mark the endpoints, then show each as included or excluded.

IV Intersection and union of intervals

Definition

$I \cap J$ corresponds to the condition "belonging to I **and** to J" ; $I \cup J$ corresponds to "belonging to I **or** to J ". A union of intervals is not always an interval.

Example 4.3

With $I = [0; 12]$ and $J = [3; 20]$, determine $I \cap J$ and $I \cup J$.

Solution:

$I \cap J = [3; 12]$ and $I \cup J = [0; 20]$.

Example 4.4

Complete, justifying with diagrams on a number line.

I	J	$I \cap J$	$I \cup J$
$]-4;3]$	$[1;5]$	$[1;3]$	$]-4;5]$
$]-\infty;2]$	$[-4;+\infty[$	$[-4;2]$	$\mathbb{R}$
$]-\infty;7]$	$[7;+\infty[$	$\{7\}$	$\mathbb{R}$
$]-\infty;5]$	$[-3;+\infty[$	$[-3;5]$	$\mathbb{R}$
$]-2;0[$	$]0;3]$	$\varnothing$	$]-2;0[\cup]0;3]$

Solution:

For each intersection, keep the points common to both representations; for each union, keep all the points of at least one of them. In the last line, 0 belongs to neither interval: the union cannot be replaced by $]-2;3]$.

Fluency · 5 to 10 minutes

Simplify $\overrightarrow{AB} + \overrightarrow{BC}$; solve $3x - 6 = 0$; calculate 10^{-2}.

Answers: $\overrightarrow{AC}$; $x = 2$; $0{,}01$.

V Absolute value and distance

Definition and properties

$|x|$ is the distance from the real number x to 0 ; $|x| \geq 0$ and

$$|x| = \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

The distance between a and b is $|b-a|$. For $a \in \mathbb{R}$ and $r \geq 0$:

$$|x-a| \leq r \iff x \in [a-r; a+r], \qquad |x-a| < r \iff x \in]a-r; a+r[.$$

Example 4.5

Calculate $|3{,}2|$, $|-1{,}6|$, $|\sqrt{3}-2|$, then the distances between 2 and 5, 2 and -5, -2 and -5.

Solution:

$3{,}2$; $1{,}6$; $2-\sqrt{3}$ since $\sqrt{3} < 2$. The distances are 3, 7 and 3.

Example 4.6

Solve: $|x-7/3| = 14$; $|x-2| \leq 3$; $|x+3| < 5$.

Solution:

For the first equation, $x - 7/3 = 14$ or $x - 7/3 = -14$, so $S = \{-35/3; 49/3\}$. Then $|x-2| \leq 3 \iff x \in [-1; 5]$. Finally $|x+3| = |x-(-3)| < 5 \iff x \in]-8; 2[$.

Example 4.7

Write $x \in [4; 10]$ as an inequality with an absolute value.

Solution:

The centre is $(4 + 10)/2 = 7$ and the radius $(10 - 4)/2 = 3$: $|x-7| \leq 3$.

VI Bounds and approximations

Vocabulary

To bound x, means giving $a \leq x \leq b$; the width is $b - a$. A lower approximation is less than or equal to x ; an upper approximation is greater than or equal to it. Rounding chooses the closest value to the required precision. Significant figures are counted from the first non-zero digit; the precision displayed must suit the context.

Example 4.8

Bound $\sqrt{2}$ between two consecutive hundredths. Give its value rounded to the nearest hundredth. Round 0 ,004567 to three significant figures, then 12345 to three significant figures in scientific notation.

Solution:

$1{,}41^2 = 1{,}9881 < 2 < 2{,}0164 = 1{,}42^2$, therefore $1{,}41 < \sqrt{2} < 1{,}42$ (amplitude 0 ,01). Rounding to the hundredth is 1 ,41. The other rounded values are 0 ,00457 and $1{,}23 \times 10^4$.

```
def encadre_racine2(p):
    # p natural number ; exact integer arithmetic
    q = 10**p
    k = 0
    while (k+1)**2 <= 2*q*q:
        k = k+1
    return k/q, (k+1)/q
```

Example 4.9

Explain why this function bounds $\sqrt{2}$. What does it return for $p = 2$? Why does it terminate?

Solution:

The loop looks for the largest integer k such that $k^2 \leq 2q^2$. On exit, $k^2 \leq 2q^2 < (k+1)^2$, therefore $k/q \leq \sqrt{2} < (k+1)/q$. For $p = 2$, it returns (1 ,41; 1,42). Since (2 $q)^2 > 2q^2$, the counter finally reaches a value that makes the test false. The final divisions are floating-point approximations ; the tests on squares are exact.