CHAPTER 8

Integral calculus

S59, S61, S63, S66

Before you begin · Antiderivatives, sign and sums

Without a calculator, in six minutes. Answer briefly. Use areas and the derivative of a product. Then choose the targeted review indicated below the questions.

1. Give an antiderivative of $2x$.

2. Determine the sign of x on $[-1;2]$.

3. Calculate the area of a rectangle with sides 3 and 5.

4. Differentiate xe^x.

5. Write $\sum_{k=1}^{n} k$.

6. Calculate the difference between the areas 2 and 5; can it be negative?

Answers

$x^2 + C$; negative then positive; 15 units of area; $(x+1)e^x$; $\frac{n(n+1)}{2}$; $2 - 5 = -3$, yes.

Targeted review after the diagnostic check

Redo only the questions you got wrong. For each question: **1.** Diff. the candidate answer; R64. **2.** Split the interval at 0, then read the sign. **3.** Multiply the lengths and state a squared unit. **4.** Write both terms of the derivative of the product; R35. **5.** Count the terms and pair the first and last. **6.** Perform the ordered subtraction and distinguish area from difference. The R questions serve as review material with the teacher, even if their whole-class routine is planned later.

Objectives

Calculate and bound an integral; distinguish area and integral; use integration by parts; study a sequence of integrals.

Course notes

Area and antiderivative

For f continuous and positive on $[a,b]$, $a \leq b$, the integral $\int_a^b f(x)\,\mathrm{d}x$ is the area under the curve, above the axis, between $x = a$ and $x = b$, expressed in units of area.

Fundamental theorem

If f is continuous on an interval I and $a \in I$, then $F_a(x) = \int_a^x f(t)\,\mathrm{d}t$ is the antiderivative of f that is zero at a. For every antiderivative F,

$$\int_a^b f(x)\,\mathrm{d}x = F(b) - F(a).$$

Every function continuous on an interval has antiderivatives. An integral is a number; an antiderivative is a function.

Proof in the positive increasing case. For $h > 0$, the rectangles give $hf(x) \leq F_a(x+h) - F_a(x) \leq hf(x+h)$. After division by h and taking the limit, the rate of change tends to $f(x)$. For $h < 0$, monotonicity gives $f(x+h) \leq f(t) \leq f(x)$ on $[x+h, x]$. By integrating,

$$(-h)f(x+h) \leq F_a(x) - F_a(x+h) \leq (-h)f(x).$$

Dividing by $-h > 0$, we get

$$f(x+h) \leq \frac{F_a(x+h) - F_a(x)}{h} \leq f(x).$$

Both bounds tend to $f(x)$. By continuity, $F_a' = f$. Any other antiderivative is $F_a + C$, hence the formula for evaluating at the bounds.

Properties

For continuous functions and bounds in the same interval:

$$\int_a^a f = 0, \quad \int_b^a f = -\int_a^b f, \quad \int_a^c f = \int_a^b f + \int_b^c f, \quad \int_a^b (\lambda f + \mu g) = \lambda \int_a^b f + \mu \int_a^b g.$$

If $a \leq b$ and $f \leq g$, then $\int_a^b f \leq \int_a^b g$. In particular, if $m \leq f \leq M$, then $m(b-a) \leq \int_a^b f \leq M(b-a)$. For a function of any sign, the integral is defined using antiderivatives. The geometric area between the curve and the axis is $\int_a^b |f|$, obtained by splitting at the sign changes. Between two curves, it equals $\int_a^b |f - g|$.

Example. $\int_{-1}^1 x\,\mathrm{d}x = 0$, but the area between the line and the axis on this interval is 1.

Mean value and integration by parts

For $a < b$, the mean value is $\mu = \frac{1}{b-a}\int_a^b f(x)\,\mathrm{d}x$. The rectangle of height μ and width $b-a$ has the same algebraic area.

Integration by parts

If u, v are of class $\mathcal{C}^1$ on $[a, b]$,

$$\int_a^b u(x)v'(x)\,\mathrm{d}x = [u(x)v(x)]_a^b - \int_a^b u'(x)v(x)\,\mathrm{d}x.$$


Proof. Integrate $(uv)' = u'v + uv'$ then isolate the required integral. The bracket is indispensable.

Example. With $u = x$ and $v' = e^x$, $\int_0^1 xe^x\,\mathrm{d}x = [xe^x]_0^1 - \int_0^1 e^x\,\mathrm{d}x = e - (e-1) = 1$.

Rectangles and sequences of integrals

For f continuous and increasing on $[a,b]$, with $a<b$ and $n\geq 1$ and $h=\frac{b-a}{n}$, the sums $h\sum_{k=0}^{n-1} f(a+kh)$ & $h\sum_{k=1}^{n} f(a+kh)$ bound the integral. Their difference is $h(f(b)-f(a))$. For $I_n=\int_0^1 x^n f(x)\,\mathrm{d}x$, start with the sign and comparisons of the integrand. If $0\leq f\leq M$, then $0\leq I_n\leq \frac{M}{n+1}$; the squeeze theorem gives $I_n\to 0$. Integration by parts can also give a relation between I_n and I_{n+1}.

Convention for index zero. In $I_n=\int_0^1 x^n f(x)\,\mathrm{d}x$, the expression x^0 denotes the constant function 1, including at the bound 0. Thus $I_0=\int_0^1 f(x)\,\mathrm{d}x$; no indeterminate form needs to be resolved.



The area between $y=x$ and $y=x^2$ on $[0,1]$ is $\frac{1}{6}$.

Linking area, antiderivative and approximation

8.1 Two areas, one integral

First use: S59 and S61.

Worked example · A change of sign

On $[0;3]$, $f(x)=x-1$ is negative before 1, positive after. The integral equals $[x^2/2-x]_0^3=3/2$. The geometric area is $-\int_0^1 f+\int_1^3 f=1/2+2=5/2$. The integral involves cancellation between two signed contributions.




Figure 8.1. For a geometric area, the two regions add up; for the integral, their signs come into play.

A1 · Immediate application For $g(x)=2x-2$ on $[0;3]$, calculate the integral and the area between the curve and the axis. How does this relate to the values in the example?

8.2 Why the area gives an antiderivative

First use: S61.

Property · Fundamental theorem

For f continuous on an interval I and $a \in I$, $A(x) = \int_a^x f(t)\,\mathrm{d}t$ is an antiderivative of f. So, for any antiderivative F, $\int_a^b f = F(b) - F(a)$. The general case is accepted without proof; the positive increasing case is proved below.

Proof. Suppose f is continuous, positive and increasing. For h>0, the comparison with two rectangles gives $f(x) \leq [A(x+h) - A(x)]/h \leq f(x+h)$. For h<0, it gives $f(x+h) \leq [A(x+h) - A(x)]/h \leq f(x)$. By continuity and squeezing, the difference quotient tends to f(x); thus $A' = f$. Any other antiderivative F differs from A by a constant; since A(a)=0, $A(b) = F(b) - F(a)$.□

Worked example · Recovering a derivative

For $A(x) = \int_0^x 2t\,\mathrm{d}t$, the area is x^2 when x is positive. The antiderivative formula also gives $A(x) = x^2$ for negative x, with the orientation of the integral. In both cases $A'(x) = 2x$.

A2 · Partial guidance On $[0;+\infty[$, set $A(x) = \int_0^x e^{-t}\,\mathrm{d}t$. Give A(x), A'(x), the direction of variation and the limit. Interpret the limit as an area.

8.3 An equivalent height, with its units

First use: S63.

Definition · Mean value

For $a < b$, the mean value of f on $[a;b]$ is $m = \frac{1}{b-a}\int_a^b f$. If f is positive, the rectangle of width b-a and height m has the same area as the region under the curve.

Worked example · Flow rate and volume

A flow rate $d(t) = 2 + t$ L/min for three minutes delivers $\int_0^3 (2+t)\,\mathrm{d}t = 10{,}5$ L. Its mean flow rate is 10.5/3 = 3.5 L/min. The volume and the mean flow rate do not have the same unit.

A3 · Immediate application Calculate the mean of x^2 on $[0;1]$. Draw the equivalent rectangle. Explain why this mean value is not the mean of the two endpoint values.

8.4 Draw and check the bounds

First use: S63 and S66.




Figure 8.2. Left-hand rectangles. Complete the drawing with the right-hand rectangles and the line at height 1/3.

Worked example · Two intervals

For x^2 on [0;1], two left-hand rectangles give $L_2 = \frac{1}{2}(0^2 + (1/2)^2) = 1/8$; the right-hand ones give $U_2 = \frac{1}{2}((1/2)^2 + 1^2) = 5/8$. The integral 1/3 does lie between these bounds.

A4 · Investigate For $f(x) = x^2$ on $[0;1]$, divide into four equal intervals. Calculate the left and right sums; justify their positions relative to the integral. Then show that, for n intervals, their difference is 1/n.

A first step: In the difference of the two sums, the interior values cancel.

8.5 Choosing what to differentiate

First use: S66.

Property · Integration by parts

For u, v of class $\mathcal{C}^1$ on $[a;b]$, $\int_a^b u'v = [uv]_a^b - \int_a^b uv'$. The formula comes from integrating $(uv)' = u'v + uv'$.

Worked example · Two successive integrations by parts

For $I = \int_0^1 x^2 e^x \,\mathrm{d}x$, differentiating a polynomial lowers its degree. First $I = [x^2 e^x]_0^1 - 2\int_0^1 xe^x \,\mathrm{d}x$. Then $\int_0^1 xe^x \,\mathrm{d}x = [xe^x]_0^1 - \int_0^1 e^x \,\mathrm{d}x = e - (e-1) = 1$. Hence I=e-2. Writing the bounds at each step avoids forgetting a constant.

A5 · Choice of strategy Calculate $\int_1^e \ln x \,\mathrm{d}x$. Justify your choice in the integration by parts and check the result by differentiating an antiderivative.

8.6 Bound before looking for a formula

First use: S66.

Worked example · Bounds without an explicit antiderivative

Let $J_n = \int_0^1 x^n/(1+x)\,dx$. Since $1/2 \le 1/(1+x) \le 1$, $1/[2(n+1)] \le J_n \le 1/(n+1)$. Both bounds tend to 0, so $J_n \to 0$, without calculating Jn for each n.

A6 · Synthesis For $n \in \mathbb{N}$, $I_n = \int_0^1 x^n e^{-x}\,dx$. Show that (I_n) is positive and decreasing. Establish $e^{-1}/(n+1) \le I_n \le 1/(n+1)$, then its limit. Does integration by parts give a relation between I_n and I_{n+1} ?

A7 · Refute Give a non-zero continuous function on $[-1;1]$ whose integral is zero. Which additional hypothesis, together with continuity, would allow us to conclude that a function with zero integral is identically zero?

One minute, spoken aloud. Explain on the figure why "zero integral" and "zero area" do not mean the same thing.

Delayed review · S79. Calculate an area between trigonometric curves, still looking for the intersections and the sign.

Investigation

Bounding an integral with rectangles - 35 min.

We want to bound $I = \int_0^1 x^2\,dx$. For an integer $n \ge 1$, we split $[0;1]$ in n intervals of the same length. Let L_n be the sum of the areas of the rectangles whose heights are the values of x^2 at left endpoints, and U_n the one obtained at the right-hand endpoints.

1. For $n = 4$, draw both families of rectangles, then calculate L_4 and U_4.
2. Justify that $L_4 \le I \le U_4$.
3. Write a Python function with argument n which returns L_n and U_n. Determine algebraically $U_n - L_n$.
4. Determine the smallest integer n for which these bounds are at most 0.01 apart.
5. Calculate the exact value of I using an antiderivative and check that it lies within the bounds obtained for $n = 4$.

Progressive exercises

E1 · Direct application. Calculate $\int_0^2 (3x^2 - 2x + 1)\,dx$ by finding an antiderivative.

E2 · Application and justification. In an orthonormal coordinate system, calculate the area of the region bounded by the curves $y = x$ and $y = x^2$, between x = 0 and x = 1. Give the result in units of area.

E3 · Method to organise. Calculate the mean value of the exponential function over the interval $[0;2]$.

E4 · Choice of strategy. Using integration by parts, calculate $\int_1^e \ln x \, dx$.

E5 · Synthesis or proof. For every $n \in \mathbb{N}$, set $I_n = \int_0^1 \frac{x^n}{1+x} \, dx$, with $x^0 = 1$ in the integrand. Prove that $0 \leq I_n \leq \frac{1}{n+1}$, then deduce the limit of (I_n).

Work from the textbook or a past paper

Barbazo, p. 260, Exercises 28 and 29. In S66: exercise 28, stating the two functions chosen for integration by parts. At home: exercise 29, to adapt this choice to a product containing a logarithm.

Homework

DM8 - A sequence of integrals - 60 min.

For every $n \in \mathbb{N}$, set $I_n = \int_0^1 x^n e^x \, dx$. For $n = 0$, the integrand is the function $x \mapsto e^x$.

1. Calculate I_0, then I_1 using integration by parts.
2. Prove that, for every $n \in \mathbb{N}$, $I_{n+1} = e - (n+1)I_n$.
3. By comparing the integrands, prove that (I_n) is positive and decreasing.
4. Show that $\frac{1}{n+1} \leq I_n \leq \frac{e}{n+1}$ for all $n \in \mathbb{N}$. Deduce the limit of (I_n).
5. In the calculation of I_{n+1} using the relation in question 2, by what factor does an error in I_n get multiplied if the other rounding errors are neglected? Explain the risk of repeating the calculation in this way.

MCQ A · integral, area and mean value

After S63; we always assume $a < b$.

For each question, only one option is correct. Justify your choice orally before checking the solutions.

1. If $F' = f$ on $[a;b]$, then $\int_a^b f(x) \, dx$ is:

A. $F(a) + F(b)$ B. $F(b) - F(a)$

C. $f(b) - f(a)$ D. 0

2. The integral of a negative function on $[a;b]$ is:

A. positive B. zero

C. negative or zero D. always a positive area

3. The average value of f on $[a;b]$ is:

A. $\frac{1}{b-a}\int_a^b f$ B. $\int_a^b f$

C. $f(a) + f(b)$ D. $b - a$

4. The geometric area between the curve and the x-axis requires:

A. preserving the signs B. splitting where the sign changes

C. a second derivative D. a random draw

5. An antiderivative of e^x is:

A. xe^x B. e^x

C. $\ln x$ D. $1/e^x$

MCQ B · integration by parts and bounds

After S66.

For each question, only one option is correct. Justify your choice orally before checking the solutions.

6. For integration by parts, you must know:

A. two functions of class $\mathcal{C}^1$ on $[a;b]$ B. a geometric sequence

C. a fixed point D. a binomial distribution

7. $\int_a^b u'v\,\mathrm{d}x$ is equal to:

A. $[uv]_a^b - \int_a^b uv'\,\mathrm{d}x$ B. $\int_a^b uv\,\mathrm{d}x$

C. $[u+v]_a^b$ D. 0

8. If f is increasing and positive on $[0;1]$, left-endpoint rectangles give:

A. an upper bound B. a lower bound

C. always the exact integral D. a negative value

9. A sequence of integrals $I_n = \int_0^1 x^n f(x)\,\mathrm{d}x$, with f continuous and positive on $[0;1]$, may often be bounded using:

A. $0 \leq x^n \leq 1$ B. $x^n = n$

C. $\ln x \geq 0$ D. the derivative of n

10. A unit of the integral of a flow rate (L/min) over a time in min is:

A. $\mathrm{L/min^2}$ B. L

C. $\mathrm{min/L}$ D. without a unit

True or false · justification expected. For each statement, indicate whether it is true or false, then write a proof or a counterexample.

1. An integral is always positive.

2. Two antiderivatives of the same function on an interval differ by a constant.

3. A rectangle approximation removes the need to check the direction of the bounds.

Review of prior work

I can interpret sign, area and mean value; choose an antiderivative or integration by parts, and find bounds.

To watch out for: confusing geometric area with algebraic integral; forgetting the bounds in $[uv]_a^b$.

MCQ solutions

1. **B.** The fundamental theorem gives $F(b) - F(a)$. It applies to an antiderivative of f, not to f itself.

2. **C.** For $a < b$, the integral keeps the sign of a function that is negative or zero. The geometric area, for its part, is positive or zero.
3. **A.** The integral is divided by the length of the interval $b - a > 0$. For a constant function, we recover this constant.
4. **B.** We calculate the areas on the intervals where the sign is constant and add their positive values. Without splitting, areas can cancel each other out.
5. **B.** The derivative of e^x is e^x. The other proposed functions have different derivatives.
6. **A.** The usual integration by parts formula applies to two functions of class $\mathcal{C}^1$ on the interval. This framework guarantees continuous derivatives and well-defined integrals.
7. **A.** The formula comes from $(uv)' = u'v + uv'$. Integrating and then isolating the first term gives $\int_a^b u'v = [uv]_a^b - \int_a^b uv'$.
8. **B.** An increasing function lies above the height taken at the left end of each interval. The left rectangles therefore give a lower bound.
9. **A.** For $x \in [0;1]$, $0 \leq x^n \leq 1$; we can multiply by $f(x) \geq 0$, then integrate. The bound may be refined using an upper bound for f.
10. **B.** The product of a flow rate in litres per minute and a duration in minutes is expressed in litres. The integral represents a cumulative volume.

True/false. False: $\int_0^1 (-1)\,\mathrm{d}x = -1$. True: the difference of two antiderivatives has zero derivative on an interval, so is constant. False: whether the function is increasing or decreasing determines which side gives the lower bound.

Solutions to the applications and guidance

Application 1. $g = 2f$: integral 3, area 5. The factor 2 is positive, so the signs are preserved and both areas are doubled.

Application 2. $A(x) = 1 - e^{-x}$, $A' = e^{-x} > 0$, hence increasing with limit 1. The area under e^{-t} between 0 and x increases and approaches 1.

Application 3. $m = 1/3$. The rectangle has width 1 and height 1/3. The mean of the endpoint values would be 1/2: it measures the area of the upper trapezium, not the area under the curve.

Application 4. $L_4 = \frac{1}{4}(0 + 1/16 + 1/4 + 9/16) = 7/32$; $U_4 = 15/32$. Monotonicity gives $L_4 \leq 1/3 \leq U_4$. $U_n - L_n = \frac{1}{n}[1^2 - 0^2] = 1/n$.

Application 5. Take u=x and $v = \ln x$, therefore $u' = 1$, $v' = 1/x$. The integral equals $[x \ln x]_1^e - \int_1^e 1 \, \mathrm{d}x = e - (e - 1) = 1$. An antiderivative is $x \ln x - x$, whose derivative is ln x.

Application 6. On [0;1], 0 $\leq x^{n+1} \leq x^n$; multiplying by $e^{-x} > 0$, then integrating proves positivity and decrease. From $e^{-1} \leq e^{-x} \leq 1$, we get the bounds, then $I_n \to 0$. Integration by parts applied to I_{n+1} gives $I_{n+1} = -e^{-1} + (n+1)I_n$. This relation is exact but is not enough on its own to calculate the limit.

Application 7. $f(x) = x$ works. If f is continuous and positive or zero, a strictly positive value would, by continuity, give a strictly positive area on a small interval, contradiction. Thus a continuous nonnegative function with zero integral is identically zero.

Session plan

Keep setting up the signed area separate from integration by parts. Work on rectangles is done first on the drawing, then with sums. In the sequence of integrals, let students choose between monotone convergence and bracketing; the second gives the value of the limit directly here.

Teaching notes and solutions

Guidance for the teacher


Start with a positive area, then show an example whose positive and negative areas cancel out. For a mean value, require a unit and an interpretation. The sequences of integrals come in S66; set aside a genuine research phase for choosing the comparison. The proof of the fundamental theorem is limited to the increasing positive case stated in the curriculum. Trigonometric calculations will be revisited after May.

Solutions to the activity


1. The four intervals have width $\frac{1}{4}$. The left sum is

$$L_4 = \frac{1}{4}(0^2 + (\frac{1}{4})^2 + (\frac{1}{2})^2 + (\frac{3}{4})^2) = \frac{7}{32}.$$

On the right, $U_4 = \frac{1}{4}((\frac{1}{4})^2 + (\frac{1}{2})^2 + (\frac{3}{4})^2 + 1) = \frac{15}{32}$. The heights of the rectangles are the values at the corresponding endpoints.



Left-hand rectangles: $L_4 = \frac{7}{32}$.



Right-hand rectangles: $U_4 = \frac{15}{32}$.

2. For $x \in [\frac{k}{4}, \frac{k+1}{4}]$, $(\frac{k}{4})^2 \leq x^2 \leq (\frac{k+1}{4})^2$ because the square function is increasing on $[0,1]$. Integration over the four intervals, then summing, gives $\frac{7}{32} \leq I \leq \frac{15}{32}$.

3.

```
def rectangles_carre(n):
    h = 1/n
    bas = h* sum((k*h)**2 for k in range (n))
    haut = h* sum((k*h)**2 for k in range (1, n+1))
    return bas, haut
```

The intermediate terms cancel out: $U_n - L_n = (\frac{1}{n})(1^2 - 0^2) = \frac{1}{n}$.

4. $\frac{1}{n} \leq 0{,}01 \iff n \geq 100$. One hundred subdivisions suffice; it is the first integer giving this width with these two sums.

5. An antiderivative of x^2 is $\frac{x^3}{3}$. Hence $I = [\frac{x^3}{3}]_0^1 = \frac{1}{3}$, indeed between $\frac{7}{32}$ & $\frac{15}{32}$.

E1. An antiderivative is $F(x) = x^3 - x^2 + x$. The integral equals $F(2) - F(0) = 8 - 4 + 2 = 6$.

E2. On $[0,1]$, $x - x^2 = x(1-x) \geq 0$, so the line lies above the parabola. The area is $\int_0^1 (x - x^2)\,\mathrm{d}x = [\frac{x^2}{2} - \frac{x^3}{3}]_0^1 = \frac{1}{6}$ unit of area.

E3. The length of the interval is 2. The mean value is $(\frac{1}{2})\int_0^2 e^x\,\mathrm{d}x = (\frac{1}{2})[e^x]_0^2 = \frac{e^2-1}{2}$.

E4. Let us choose $u = \ln x$, $u' = \frac{1}{x}$, $v' = 1$ and $v = x$ on $[1, e]$. By parts, $\int_1^e \ln x \, dx = [x \ln x]_1^e - \int_1^e 1 \, dx = e - (e-1) = 1$.

E5. On $[0,1]$, $1 + x \geq 1$ and $x^n \geq 0$, hence $0 \leq \frac{x^n}{1+x} \leq x^n$. Integration gives $0 \leq I_n \leq [\frac{x^{n+1}}{n+1}]_0^1 = \frac{1}{n+1}$. The squeeze theorem gives $I_n \to 0$.

Solutions to homework DM8

1. $I_0 = \int_0^1 e^x \, dx = e - 1$. For I_1, by parts with $u = x$, $v' = e^x$ gives $I_1 = [xe^x]_0^1 - \int_0^1 e^x \, dx = e - (e-1) = 1$.

2. For $n \geq 0$, let us choose $u = x^{n+1}$, $u' = (n+1)x^n$, $v' = e^x$ & $v = e^x$. The functions are differentiable with continuous derivatives on $[0,1]$. Then

$$I_{n+1} = [x^{n+1}e^x]_0^1 - (n+1)\int_0^1 x^n e^x \, dx = e - (n+1)I_n.$$

3. On $[0,1]$, $0 \leq x^{n+1} \leq x^n$. By multiplying by $e^x > 0$ then integrating, $0 \leq I_{n+1} \leq I_n$: the sequence is positive and non-increasing. It is even strictly positive and strictly decreasing, since the integrands of I_n and of $I_n - I_{n+1}$ are strictly positive on $]0,1[$.

4. Since $1 \leq e^x \leq e$ and $x^n \geq 0$ on $[0,1]$, we can integrate $x^n \leq x^n e^x \leq e x^n$. Hence

$$\frac{1}{n+1} \leq I_n \leq \frac{e}{n+1}.$$

Both bounds tend to 0; the squeeze theorem gives $I_n \to 0$.

5. If $\widetilde{I}_n = I_n + \varepsilon_n$ and the relation is then applied without further error, then

$$\widetilde{I}_{n+1} - I_{n+1} = -(n+1)(\widetilde{I}_n - I_n) = -(n+1)\varepsilon_n.$$

The absolute value of the error is multiplied by $n+1$. In machine arithmetic, new errors add up at each step. The decrease of the exact integrals therefore does not guarantee stability of this method of calculation.

Solutions to the textbook or past-paper work

Exercise 28. On $[0;1]$, let us choose $u(x) = x$ and $v(x) = e^x$. These functions are of class $\mathcal{C}^1$, $u'(x) = 1$ & $v'(x) = e^x$. Integration by parts gives

$$\int_0^1 xe^x \, dx = [xe^x]_0^1 - \int_0^1 e^x \, dx = e - (e-1) = 1.$$

Exercise 29. On $[1;e]$, let us choose $u(x) = \ln x$ and $v(x) = \frac{x^2}{2}$. Then $u'(x) = \frac{1}{x}$ & $v'(x) = x$; these functions are of class $\mathcal{C}^1$. Hence

$$\begin{aligned}\int_1^e x \ln x \, dx &= \left[\frac{x^2}{2}\ln x\right]_1^e - \frac{1}{2}\int_1^e x \, dx \\ &= \frac{e^2}{2} - \frac{e^2-1}{4} = \frac{e^2+1}{4}.\end{aligned}$$