Sequence 11

CROSS-TABULATING TWO QUALITATIVE VARIABLES

1–12 March 2027

I Two-way tables and marginal frequencies

Definitions

A qualitative variable describes a category. It is nominal if its categories have no natural order (means of transport), ordinal if they are ordered (level of satisfaction). A two-way table counts individuals according to two variables. The sums of the rows and columns are the margins. A marginal frequency divides a margin by the total count.

Example 11.1

In a population of 100 individuals, two characteristics A, B are observed:

	A	$\overline{A}$	Total
B	20	15	35
$\overline{B}$	12	53	65
Total	32	68	100

Calculate the marginal frequencies of A, B, $\overline{B}$, then the frequency of individuals satisfying both A and B.

Solution:

Example 11.2

A Year 12 class of 32 students is split as follows. Complete the margins and calculate the marginal frequencies "CS" and "girls".

	Boys	Girls	Total
CS	12	9	.....
No CS	6	5	.....
Total	.....	.....	32

Solution:

II Conditional frequencies and reading sentences

Definition

The frequency of A **among the individuals of** B, if this subpopulation is nonempty, is

$$f_B(A) = \frac{\text{count satisfying } A \text{ and } B}{\text{count of } B}.$$

The denominator corresponds to the population introduced by "among".

Example 11.3

In the CS table, calculate the frequency of girls among the students without CS, then the frequency of students without CS among the girls. Are the two results equal?

Solution:

Example 11.4

In the table of 100 individuals, calculate $f_B(A)$, $f_A(B)$ and $f_{\overline{A}}(B)$. Translate each result into a sentence.

Solution:

Example 11.5

Does "60 % of cyclists are students" allow us to conclude that 60 % of students are cyclists?

Solution:

III Completing a table from frequencies

Example 11.6: Fictional situation

A group has 200 people, of whom 120 use a bus (B). Among these 120 people, 75 % have a pass (A). Among the other 80, 25 % have a pass.

1. Construct the two-way table $B/\overline{B}$ and $A/\overline{A}$ with its margins.
2. Calculate the marginal frequency of A.
3. Calculate the frequency of B among A.

Solution:

Example 11.7

A headline announces “Three pass holders in four take the bus”, based on $f_B(A) = 75\,\%$. Explain the error and propose a correct sentence.

Solution:

IV Processing a file and collecting real data

```
# Exemple fictif : (transport, satisfaction)
donnees = [("bus", "oui"), ("marche", "oui"),
           ("bus", "non"), ("velo", "oui"),
           ("bus", "oui"), ("marche", "non")]

bus = [r for r in donnees if r[0] == "bus"]
bus_satisfaits = [r for r in donnees
                  if r[0] == "bus" and r[1] == "oui"]
autres = [r for r in donnees if not r[0] == "bus"]
```

Example 11.8

Give the sizes of the three lists. Calculate the frequency of satisfaction among bus users. Replace `and` by `or` in the second filter: how many rows are then selected?

Solution:

Activity with the class data

Collect anonymously two variables: main means of transport and journey time in ordered categories (< 15 min; from 15 to under 30 min; ≥ 30 min). One row per response, without names. Construct the two-way table, then two conditional frequencies in opposite directions. State the count, the collection date and any non-responses.

Example 11.9

Carry out the collection and write two justified conclusions. Explain why the results do not automatically describe all students in Madagascar.

Solution: