Sequence 13

SYNTHESIS PROBLEMS, REASONING AND FLUENCY

30 March–9 April 2027

I Implication, converse and counterexample

Tools to use in context

"If P, then Q" is an implication $P \Rightarrow Q$. Its converse is $Q \Rightarrow P$; it is not automatically true. An equivalence $P \iff Q$ requires both implications. A counterexample is enough to refute a universal statement; a few examples are not enough to prove it. The contrapositive not $Q \Rightarrow$ not P is equivalent to $P \Rightarrow Q$.

Example 13.1

For an integer n: "If n is a multiple of 4, then n is even." Prove the implication, write its converse and its contrapositive, then say which are true.

Solution:

Example 13.2

For $x \in \mathbb{R}$, study "$x = 3 \Rightarrow x^2 = 9$" and its converse. Give a correct equivalence.

Solution:

II Negation, case analysis and proof

Example 13.3

Negate correctly: "$x > 2$"; "$x \in [1;4]$"; "All students play sport"; "A and B both occur".

Solution:

Example 13.4

Show by case analysis that $n(n+1)$ is even for every integer n. Then show that $n^2 + n + 1$ is odd.

Solution:

Example 13.5

A student writes "n^2 even implies n even, because I checked it for $n = 2, 4, 6$". Replace this justification with a proof.

Solution:


III Combining algebra, functions and percentages

Example 13.6: Original situation

Two fictional rental rates are offered for x hours, with $x \in [0; 8]$: $A(x) = 3000+2000x$ Ar and $B(x) = 3500x$ Ar.

1. Calculate the costs for 1 h and 4 h.
2. Find where the rates are equal, then the durations for which A is cheaper.
3. A 10 % discount is granted on the whole of rate A. Give the new rate A_r and the new comparison threshold.

Solution:



```
def tarif_moins_cher(x):
    a = 3000 + 2000*x
    b = 3500*x
    if a < b:
        return "A"
    elif b < a:
        return "B"
    else:
        return "egalite"
```

Example 13.7

Predict the results for $x = 1, 2, 4$. Modify the line that computes a to take the discount into account.

Solution:

