Sequence 14

PROBABILITY: CONDITIONING AND WEIGHTED TREES

26 April–24 May 2027

I Sample space, outcomes and events

Vocabulary

A random experiment has several possible results, without it being known in advance which will occur. An outcome is a possible result; the sample space Ω is the set of them. An event is a subset of Ω; an elementary event contains a single outcome. The impossible event is $\varnothing$; the certain event is Ω.

Operations on events

$\overline{A}$ is the complementary event of A. $A \cap B$ means "A and B"; $A \cup B$, "A or B", possibly both. Two events are mutually exclusive if $A \cap B = \varnothing$; otherwise, they are compatible.

Example 14.1

A six-sided die is rolled. Write Ω, A: "get an even number", B: "get 4", then $\overline{A}$, $A \cap B$ and $A \cup B$. Are A, B compatible?

Solution:

Example 14.2

With $A = \{a; b; c; d; e\}$ and $B = \{b; e; f; g; h\}$, give the intersection and the union. Can $\overline{A}$ be determined without specifying the sample space?

Solution:

II Probability model and equally likely outcomes

Probability on a finite sample space

To each outcome, a model assigns a probability that is positive or zero; the sum is 1. The probability of an event is the sum of the probabilities of its outcomes. We have

$$P(\Omega) = 1, \quad P(\varnothing) = 0, \quad P(\overline{A}) = 1 - P(A).$$

With equally likely outcomes, all outcomes have the same probability and


$$P(A) = \frac{\text{Card}(A)}{\text{Card}(\Omega)}.$$

Equal likelihood is an assumption of the model, not a consequence of the list of outcomes alone.

Example 14.3

Assume the die is fair. For $A =$ "even" and $C =$ "greater than or equal to 5", calculate $P(A)$, $P(C)$, $P(A \cap C)$ and $P(A \cup C)$.


Solution:



Example 14.4

A drawing pin can land point up or point down. Can we deduce that each outcome has probability 1/2? How can a model be proposed?

Solution:



III Conditional probability from a table

Definition

For $P(A) > 0$, the probability of B given A is

$$P_A(B) = \frac{P(A \cap B)}{P(A)}, \qquad P(A \cap B) = P(A)P_A(B).$$

For an equally likely draw within a finite population:

$$P_A(B) = \frac{\text{Card}(A \cap B)}{\text{Card}(A)}.$$

We also write $P(B \mid A)$. $P_A(B)$ and $P_B(A)$ are generally not equal.


Example 14.5

A student is drawn uniformly from the class of 32 students studied earlier. G: "boy", N: "takes CS".

	G	$\overline{G}$	Total
N	12	9	21
$\overline{N}$	6	5	11
Total	18	14	32

Calculate $P(G)$, $P(N)$, $P(G \cap N)$, $P_N(G)$ and $P_G(N)$.

Solution:




Watch out. Conditioning requires the known event to have nonzero probability; we do not divide by zero.

IV Weighted trees and the probability of a path

Rules

At each node, the sum of the probabilities of the outgoing branches is 1. A second-level branch carries a conditional probability. A path represents an intersection of events; its probability is the product of the probabilities written along that path.



Example 14.6

We know $P(A) = 0.6$, $P_A(B) = 0.7$ and $P_{\overline{A}}(\overline{B}) = 0.9$. Complete the other branches and calculate the probabilities of the four paths.

Solution:

Example 14.7

Justify the multiplication along the path A then B from the definition of $P_A(B)$.


Solution:

V From a table to a tree

Example 14.8

For the class of 32 students, construct a tree starting with CS / no CS, then boy / girl. Then construct the tree in the reverse order. Check, in both trees, the probability of choosing a girl who takes CS.


Solution:



Example 14.9

An urn contains 3 red balls and 2 blue balls. Two balls are drawn in succession **without replacement**. Construct the tree of colours and calculate the probability of drawing two red balls.

Solution:




VI Inverting conditioning and false positives

Example 14.10: Fictional component check

A batch contains 10000 components, of which 100 are defective. A test is positive for 99 of the defective components and for 495 of the non-defective components. A component is chosen uniformly. D: "defective"; T: "test positive".

1. Complete the two-way table of test / true state.
2. Calculate $P_D(T)$ and $P_T(D)$.
3. Give the number of false positives and false negatives.
4. Calculate the sensitivity $P_D(T)$ and the specificity $P_{\overline{D}}(\overline{T})$.

Solution:



Reading a test result

"Positive among the defective" and "defective among the positives" correspond to two different conditionings. Always name the reference group before the calculation.

VII Simulation and the law of large numbers

Popularised version

When an experiment is repeated a large number of times in the same model with independent repetitions, the observed frequency of an event is, barring exceptions, close to its probability. It fluctuates: it does not necessarily get closer with each new trial.

```
from random import Random

def frequences_six(n, graine=1):
    # n entier >= 1 ; modele de lancers independants
    rng = Random(graine)
    succes = 0
    frequences = []
    for k in range(1, n+1):
        if rng.randint(1, 6) == 6:
            succes = succes+1
        frequences.append(succes/k)
    return frequences
```

Example 14.11

Plot the frequencies for $n = 1000$, with several seeds. Compare with $1/6$. Why fix the seed? Can we require the final frequency to be exactly $1/6$?

Solution:

Example 14.12

Modify the program to track the frequency of the event "even result". Which reference probability should be used?

Solution: