Matrices

Lecture and worked examples

Objectives

Compute with matrices (sum, product by a real number, product); find the inverse of a matrix of order 2, and of order 3 with a calculator; compute a power, in particular by induction; solve a linear system in matrix form; represent a transformation of the plane by a matrix.

Introductory activity

We represent a point $M(x, y)$ by the column $X = \begin{pmatrix} x \\ y \end{pmatrix}$. For a matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, we define the product $AX = \begin{pmatrix} ax + by \\ cx + dy \end{pmatrix}$: it is the column of a new point.

Introductory activity (2)

1. Let $R = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Compute the images under R of the vertices of the triangle $O(0,0)$, $P(2,0)$, $Q(2,1)$. Draw both triangles. Which transformation do you recognise?
2. Let $E = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$. Same question.
3. Compute $R(EX)$ for any $X = \begin{pmatrix} x \\ y \end{pmatrix}$. Find a matrix C such that $R(EX) = CX$ for every X.
4. Compute likewise $E(RX)$ and the corresponding matrix. Are the two composite transformations the same? Check on the point Q.

Introductory activity (3)

5. Suggest a rule for computing C directly from R and E: it is the matrix product RE.

Vocabulary and operations

A *matrix* of size $n \times p$ is an array of real numbers with n rows and p columns. The entry in row i and column j of A is written a_{ij}. An $n \times 1$ matrix is a *column* matrix, a $1 \times p$ matrix a *row* matrix, an $n \times n$ matrix a *square* matrix of order n. The *identity* matrix I_n has 1s on the diagonal and 0s elsewhere; the *zero* matrix has only 0s. A square matrix is *diagonal* if its entries off the diagonal are zero.

Vocabulary and operations

For A and B of the same size and k real, $A + B$ and kA are computed entry by entry.

If A has size $n \times p$ and B has size $p \times q$, the product AB is the $n \times q$ matrix with entries

$$c_{ij} = \sum_{k=1}^{p} a_{ik} b_{kj} = a_{i1} b_{1j} + a_{i2} b_{2j} + \cdots + a_{ip} b_{pj}.$$

Vocabulary and operations

$$\begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 4 & -2 \end{pmatrix} = \begin{pmatrix} 1 \times 0 + 2 \times 4 & 1 \times 1 + 2 \times (-2) \\ -1 \times 0 + 3 \times 4 & -1 \times 1 + 3 \times (-2) \end{pmatrix} = \begin{pmatrix} 8 & -3 \\ 12 & -7 \end{pmatrix}.$$

Vocabulary and operations

When the products are defined: $(AB)C = A(BC)$, $A(B+C) = AB+AC$, $(A+B)C = AC+BC$, $(kA)B = k(AB)$, and $AI_n = I_nA = A$ for A square of order n.

Vocabulary and operations

1. In general $AB \neq BA$; consequently $(A+B)^2 = A^2 + AB + BA + B^2$, and not $A^2 + 2AB + B^2$.
2. A product can be zero without either factor being zero: $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$.

Powers

For A square of order n, let $A^0 = I_n$ and $A^{k+1} = A^k A$ for every natural number k.

Powers

For $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, we compute $A^2 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$, $A^3 = \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}$ and conjecture $A^k = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$.

This is true for $k = 0$; if it is true for k, then $A^{k+1} = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & k+1 \\ 0 & 1 \end{pmatrix}$.

Powers

If D is diagonal with diagonal entries $d_1, \ldots, d_n$, then D^k is diagonal with entries $d_1^k, \ldots, d_n^k$.

Inverse and systems

A square matrix A of order n is *invertible* if there is a matrix B such that $AB = BA = I_n$. This matrix is unique; it is written A^{-1}.

Inverse and systems

Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. The matrix A is invertible if and only if $ad - bc \neq 0$, and then

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

The number $ad - bc$ is the *determinant* of A, written $\det A$.

Inverse and systems

Let $M = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$. Computation gives $AM = MA = (ad - bc)I_2$. If $ad - bc \neq 0$, the matrix $\frac{1}{ad - bc} M$ is therefore the inverse of A.

If $ad - bc = 0$, then $AM = 0$. Suppose A is invertible: multiplying on the left by A^{-1} gives $M = 0$, so $a = b = c = d = 0$ and A is zero. But the zero matrix is not invertible ($0 \times B = 0 \neq I_2$): a contradiction.

Inverse and systems

A linear system of n equations in n unknowns can be written $AX = B$, where A is the matrix of coefficients, X the column of unknowns and B the column of right-hand sides. If A is invertible, the system has a unique solution: $X = A^{-1}B$.

Inverse and systems

If $AX = B$, then $A^{-1}AX = A^{-1}B$, that is $X = A^{-1}B$. Conversely, $A\left(A^{-1}B\right) = B$.

Inverse and systems

For a matrix of order 3, obtain A^{-1} with a calculator or software, then check one or two entries of AA^{-1} by hand.

The rotation with centre O and angle θ maps the point with column X to the point with column $R_\theta X$, where

$$R_\theta = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}.$$

Transformations of the plane and sequences

The point $M(x, y)$ has affix $z = x + iy$, and its image has affix $e^{i\theta}z = (\cos\theta + i\sin\theta)(x + iy) = (x\cos\theta - y\sin\theta) + i(x\sin\theta + y\cos\theta)$ (chapter 3): these are the entries of $R_\theta X$.

Transformations of the plane and sequences

If $X_{k+1} = AX_k$ for every k, then $X_k = A^k X_0$ (immediate induction). A sequence defined by $u_{k+2} = au_{k+1} + bu_k$ reduces to this with $X_k = \begin{pmatrix} u_{k+1} \\ u_k \end{pmatrix}$ and $A = \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix}$.