CHAPTER 10

Graphs

Vocabulary, degrees, adjacency matrix, counting walks

Objectives. Model a situation with a graph; use the vocabulary of graphs (order, degree, walk, cycle, connectedness, complete graph, directed graph); write the adjacency matrix of a graph; count the walks of given length with the powers of this matrix.

Prerequisites. Chapter 9 (product and powers of matrices); proof by induction; counting.

Fluency

Fluency · No calculator, five minutes.

1. Compute $\binom{4}{2}$ and $\binom{8}{2}$.
2. Compute $1+2+\cdots+7$.
3. Compute $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^2$.
4. In a group of 5 people, each shakes hands with all the others. How many handshakes are there?
5. A sum of numbers is odd. What can be said about the number of odd terms?
6. Write the entry in row i and column j of M^2 in terms of the entries of M, for M of order n.

Introductory activity

Activity 1 · A network of bush taxis 40 min . Five towns A, B, C, D, E are linked by direct bush-taxi routes, in both directions: A-B, A-C, A-D, B-C, C-D and D-E.

1. Represent the network by a diagram: one point per town, one line per route.
2. For each town, count the number of routes leaving it. Compute the sum of these numbers and compare it with the number of routes. Explain.
3. List the journeys from A to C that use exactly two routes (a town may be visited again). How many are there?
4. Same question for the journeys from A to E using exactly three routes.
5. We store the information in a table M with 5 rows and 5 columns: the entry in row X and column Y is 1 if there is a direct route between X and Y, 0 otherwise. Write M, then compute M^2 and M^3 with a calculator. Compare some entries with questions 3 and 4.
6. State a conjecture.


Observing is not proving: the properties conjectured here are established in the course.

Course notes

Vocabulary

Definition · Graph. A (undirected) *graph* consists of a finite set of *vertices* and a set of *edges*, each edge joining two distinct vertices; two joined vertices are *adjacent*. We consider graphs without multiple edges. The *order* of the graph is its number of vertices. The *degree* of a vertex is the number of edges of which it is an end. A graph is *complete* if any two distinct vertices are adjacent.

Definition · Walk, cycle, connectedness. A *walk* is a sequence of vertices in which any two consecutive vertices are adjacent; its *length* is its number of edges (a vertex or an edge may be used again). A walk is *closed* if its ends are equal. A *cycle* is a closed walk of length at least 3 whose edges are distinct. A graph is *connected* if any two vertices are joined by a walk.



The network of the activity: a graph of order 5 with 6 edges.

Property · Sum of the degrees. In a graph, the sum of the degrees of the vertices is twice the number of edges. Consequently, the number of vertices of odd degree is even.

Proof. Each edge has two ends: it is counted twice in the sum of the degrees. The sum of the degrees is even; now a sum of integers is even if and only if it contains an even number of odd terms. □

Property · Complete graph. The complete graph of order n has $\binom{n}{2} = \frac{n(n-1)}{2}$ edges.

Definition · Directed graph, weighted graph. In a *directed* graph, the links, called *arcs*, have a direction: the arc (X, Y) goes from X to Y. A *path* is a sequence of vertices in which each is joined to the next by an arc. A graph is *weighted* when each edge or arc carries a number, called its weight.

Adjacency matrix

Definition · Adjacency matrix. The vertices of a graph of order n being numbered from 1 to n, its *adjacency matrix* is the square matrix M of order n whose entry m_{ij} is 1 if the vertices i and j are joined (by an edge, or by an arc from i to j for a directed graph), and 0 otherwise. For an undirected graph, M is symmetric: $m_{ij} = m_{ji}$.

Property · Number of walks of length n. Let M be the adjacency matrix of a graph and $n \geq 1$ an integer. The entry in row i and column j of M^n equals the number of walks (of paths, for a directed graph) of length n from vertex i to vertex j.

Proof. Let $m_{ij}^{(n)}$ be the entries of M^n and $c_{ij}^{(n)}$ the number of walks of length n from i to j. For $n = 1$, a walk of length 1 from i to j is an edge: $c_{ij}^{(1)} = m_{ij}$.

Assume $m_{ij}^{(n)} = c_{ij}^{(n)}$ for all i, j. A walk of length $n+1$ from i to j consists of a walk of length n from i to a vertex k, followed by an edge from k to j. Sorting these walks by the second-to-last vertex k, we obtain

$$c_{ij}^{(n+1)} = \sum_{k=1}^{p} c_{ik}^{(n)} \, m_{kj} = \sum_{k=1}^{p} m_{ik}^{(n)} \, m_{kj} = m_{ij}^{(n+1)},$$

where p is the order of the graph, since $M^{n+1} = M^n M$. The property holds for $n+1$. □

Key point · Consequences. The diagonal entry $\left(M^2\right)_{ii}$ is the degree of vertex i (undirected graph).

Beyond the syllabus · Connectedness and powers of the adjacency matrix. A graph of order $p \geq 2$ is connected if and only if every off-diagonal entry of $M + M^2 + \cdots + M^{p-1}$ is strictly positive. Indeed, by the theorem, the entry with indices $i \neq j$ counts the walks from i to j of length 1 to $p-1$. If it is strictly positive, i and j are linked. Conversely, if i and j are linked, choose a walk from i to j of minimal length. It never passes twice through the same vertex: otherwise, removing the loop between the two visits would give a shorter walk. So it visits at most p distinct vertices and its length is at most $p-1$. This criterion is not part of the syllabus and is not assessed.

Exercises

Exercise 10.1 · Vocabulary. A graph G has vertices 1, 2, 3, 4, 5, 6 and edges 1-2, 1-3, 2-3, 3-4, 4-5, 4-6, 5-6.

1. Give the order of G and the degree of each vertex, and check the property of the sum of the degrees.
2. Is G complete? connected?
3. Give a walk of length 3 from 1 to 6 and a cycle through 5.

Exercise 10.2 · Handshakes.

1. Can one build a graph of order 7 all of whose vertices have degree 3?
2. Can six people each shake hands with exactly three others? If so, how many handshakes are there?

Exercise 10.3 · A directed graph. A directed graph has vertices A, B, C and arcs $A \to B$, $A \to C$, $B \to C$ and $C \to A$.

1. Write its adjacency matrix M (order A, B, C). Is it symmetric?
2. Compute M^2 and M^3.
3. How many paths of length 3 are there from A to A? Write them out.

Exercise 10.4 · A tournament. Eight teams play a tournament: each meets each of the others once. Model with a graph and determine the number of matches.

Exercise 10.5 · Degrees and the square of the matrix. A graph has adjacency matrix

$$M = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}.$$

1. Draw it.
2. Compute M^2 and compare its diagonal with the degrees of the vertices.
3. Prove that for every undirected graph, $\left(M^2\right)_{ii}$ is the degree of vertex i.

Exercise 10.6 · Testing connectedness. A graph has vertices 1 to 5 and edges 1-2, 2-3, 3-1, 4-5. Let M be its adjacency matrix.

1. Compute, with a calculator, $S = M + M^2 + M^3 + M^4$.
2. Is the graph connected? Justify directly, then check that this result can be read from the entries of S.

Exercise 10.7 · **Counting triangles.** A triangle of a graph is a set of three pairwise adjacent vertices.

1. Prove that the number of triangles is $\frac{1}{6}$ of the sum of the diagonal entries of M^3.
2. Apply this to the complete graph of order 4 and to the graph of exercise 10.5.

Problem

Problem · Drawing without lifting the pencil. An *Eulerian trail* is a walk that uses every edge of the graph exactly once.

1. Assume that a graph has an Eulerian trail whose ends are distinct. Prove that its two ends have odd degree and that all the other vertices have even degree. What can be said if the trail is closed?
2. We take the converse for granted (Euler's theorem): a connected graph all of whose vertices have even degree, except zero or two, has an Eulerian trail; if there are two vertices of odd degree, it starts at one and ends at the other.

   Consider the "house": a square $abdc$ (a and b at the bottom, c and d at the top) with its two diagonals, topped by a roof c-e-d. List the edges and the degrees.
3. Can the house be drawn without lifting the pencil or going over a line twice? If so, where must one start? Give such a trail.
4. Can it be done returning to the starting point?

Beyond the syllabus · Euler's theorem. The converse stated in question 2 (Euler's theorem) is not part of the syllabus: it is admitted and is not assessed.