
Chapter 4

Reciprocal (inverse) function

Skills: finding a domain, differentiating, studying a sign, drawing up a table of variations, graphing

4.1 Reference notes

The reciprocal function is defined by $f(x) = \dfrac{1}{x}$ on $\mathbb{R} \setminus \{0\}$.



Key points

$f'(x) = -\dfrac{1}{x^2} < 0$ for every $x \neq 0$. The reciprocal function is strictly decreasing on each of the intervals $]-\infty, 0[$ and $]0, +\infty[$. The lines $x = 0$ and $y = 0$ are asymptotes.

Common mistake

One cannot write that the reciprocal function is decreasing "on $\mathbb{R} \setminus \{0\}$" without specifying the two intervals. For example $-1 < 1$ but $f(-1) = -1 < 1 = f(1)$, which would contradict the global definition of decreasing.

For a function $g(x) = P(x) + \dfrac{k}{x}$, proceed in this order: domain; derivative; common denominator and factorisation; sign of the derivative; variations; limits or behaviour at the boundaries; useful points; graph.

Worked example

On $]0, +\infty[$, $g(x) = x + \dfrac{4}{x}$ satisfies $g'(x) = 1 - \dfrac{4}{x^2} = \dfrac{(x-2)(x+2)}{x^2}$. The denominator is positive; g' is negative on $]0, 2[$ then positive on $]2, +\infty[$. The function has a minimum $g(2) = 4$.

Session 23 — Definition and graph of the reciprocal function

Reciprocal (inverse) function

15 Dec 2026 • B: 2 h – C: 1 h

Session objectives

- Find the domain.
- Draw the curve.
- Use symmetries.

Fluency – 6 to 8 minutes, no calculator


1. Simplify $3x + 5 - x + 2$. $2x + 7$
2. Give the excluded value of $\dfrac{x+1}{x}$. $x = 0$
3. Calculate $\dfrac{1}{5}$ and $\dfrac{1}{-5}$. $\dfrac{1}{5}$ **and** $-\dfrac{1}{5}$
4. Differentiate $3x^2 - 5x + 1$. $6x - 5$
5. Solve $(x-2)(x+1) = 0$. $x = -1$ **or** $x = 2$

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: find the domain and draw the curve.

Investigation

Calculate $\dfrac{1}{x}$ for $x = -4, -2, -1, -\dfrac{1}{2}, \dfrac{1}{2}, 1, 2, 4$. Plot the points and describe the symmetries. Why is $x = 0$ missing?



Detailed solution – teacher edition

The values are respectively $-\frac{1}{4}, -\frac{1}{2}, -1, -2, 2, 1, \frac{1}{2}, \frac{1}{4}$. The domain excludes 0; the curve is symmetric about the origin. Have students say aloud which piece of data triggers the choice of method.

Written record

The reciprocal function is $f(x) = \dfrac{1}{x}$ on $\mathbb{R} \setminus \{0\}$. Its curve is a hyperbola symmetric about the origin.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Detailed solution – teacher edition

The values are respectively $-\frac{1}{4}, -\frac{1}{2}, -1, -2, 2, 1, \frac{1}{2}, \frac{1}{4}$. The domain excludes 0; the curve is symmetric about the origin.

Practice – essential

1. Give the domain and the asymptotes of $f(x) = 5 + \dfrac{7}{x}$.
2. Calculate $f'(x)$ and study its sign.
3. Study the function $g(x) = x + \dfrac{25}{x}$ on $]0, +\infty[$.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$; asymptotes $x = 0$ and $y = 5$. $f'(x) = -\dfrac{7}{x^2} < 0$. For g, $g'(x) = 1 - \dfrac{25}{x^2}$ and the minimum is reached at $x = 5$, with value 10.

Textbook references

Further reading: Calao, course pp. 88–93; exercises p. 94, no. 1–5. The exercise reproduced in this session is identified below by its page and number; the reference also lets you find the figures and digital resources of the textbook.

Calao textbook – p. 101, exercise 42

The aim of this exercise is to draw the curves representing the functions f and g defined on $\mathbb{R}^*$ by: $f(x) = \dfrac{2}{x}$ and $g(x) = 4 - \dfrac{3}{x}$.

1. Complete the table of values for $x \in \{-5; -4; -2; -0.5; -0.2; -0.1; 0.1; 0.2; 0.5; 1; 2; 4; 5\}$.
2. Using a table of values, construct the curves representing the functions f and g.

Detailed solution – teacher edition

Calculate each image, respecting the excluded value 0. Both curves have the vertical asymptote $x = 0$. The horizontal asymptote of f is $y = 0$; that of g is $y = 4$. To check the drawing, verify for example $f(1) = 2$, $f(-1) = -2$, $g(1) = 1$ and $g(-1) = 7$.

Homework

Redo practice exercise 2 with the following data: replace the initial value by 235 and the rate or common ratio by 7. Write it up in four lines: data, formula, calculation, conclusion.

Detailed solution – teacher edition

The answer depends on the exercise revisited. Check above all the consistency of the model, the number of iterations, the rounding and the concluding sentence.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1, 2 and 3.

Method

1 h session: 7 min of fluency; 10 min of pooling and written record; 28 min of essential exercises; 5 min of correction; 3 min of exit ticket.
2 h session: add 20 min of investigation, 20 min of formalisation, a second independent exercise and 10 min of comparative correction.
Differentiation: give struggling students a four-box sheet (data, model, calculation, conclusion). Ask quick students to produce a counterexample, a justification or a second method.

Session 24 — Behaviour at the boundaries and asymptotes

Reciprocal (inverse) function

17 Dec 2026 • B: 1 h – C: 2 h

Session objectives

- Describe the behaviour at the boundaries.
- Identify the asymptotes.
- Read a graph.

Fluency – 6 to 8 minutes, no calculator


1. Calculate 5 % of 120. 6
2. Calculate $\frac{1}{6}$ and $\frac{1}{-6}$. $\frac{1}{6}$ **and** $-\frac{1}{6}$
3. Differentiate $3x^2 - 5x + 1$. $6x - 5$
4. Solve $(x-2)(x+1) = 0$. $x = -1$ **or** $x = 2$
5. Give the excluded value of $\frac{x+1}{x}$. $x = 0$

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: describe the behaviour at the boundaries and identify the asymptotes.

Investigation

Using a table of values near 0 and for large values of $|x|$, describe $\frac{1}{x}$. Translate each behaviour into an asymptote and a sentence.



Detailed solution – teacher edition

Near 0, the values become very large in absolute value: asymptote $x = 0$. As $|x|$ grows, $\frac{1}{x}$ gets closer to 0: asymptote $y = 0$. Have students say aloud which piece of data triggers the choice of method.

Written record

For $f(x) = \dfrac{1}{x}$, $x = 0$ is a vertical asymptote and $y = 0$ a horizontal asymptote. The branches approach them without crossing them.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Detailed solution – teacher edition

Near 0, the values become very large in absolute value: asymptote $x = 0$. As $|x|$ grows, $\dfrac{1}{x}$ gets closer to 0: asymptote $y = 0$.

Practice – essential

1. Give the domain and the asymptotes of $f(x) = 6 + \dfrac{8}{x}$.
2. Calculate $f'(x)$ and study its sign.
3. Study the function $g(x) = x + \dfrac{36}{x}$ on $]0, +\infty[$.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$; asymptotes $x = 0$ and $y = 6$. $f'(x) = -\dfrac{8}{x^2} < 0$. For g, $g'(x) = 1 - \dfrac{36}{x^2}$ and the minimum is reached at $x = 6$, with value 12.

Textbook references

Further reading: Calao, course pp. 88–93; exercises p. 94, no. 4–9. The exercise reproduced in this session is identified below by its page and number; the reference also lets you find the figures and digital resources of the textbook.

Calao textbook – p. 95, exercise 16

We want to draw the curve representing the function g defined on $\mathbb{R}^*$ by: $g(x) = 3 - \dfrac{2}{x}$.

1. Complete a table of values for

   $$x \in \{-1\,000; -500; -200; -100; -50; 50; 100; 200; 500; 1\,000\}.$$

   What can be deduced about the behaviour of g at $-\infty$ and at $+\infty$?
2. Complete a table of values for $x = -0.5; -0.1; -0.01; 0.01; 0.1; 0.5$. What can be deduced about the behaviour of g at 0?
3. Copy and complete the table of values from the textbook, then draw the curve representing the function g.

Detailed solution – teacher edition

As $x \to \pm\infty$, $-2/x \to 0$, so $g(x) \to 3$. As $x \to 0^-$, $-2/x \to +\infty$; as $x \to 0^+$, $-2/x \to -\infty$. The asymptotes are $x = 0$ and $y = 3$.

Homework

Redo practice exercise 3 with the following data: replace the initial value by 240 and the rate or common ratio by 2. Write it up in four lines: data, formula, calculation, conclusion.

Detailed solution – teacher edition

The answer depends on the exercise revisited. Check above all the consistency of the model, the number of iterations, the rounding and the concluding sentence.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1 and 2.

Method

1 h session: 7 min of fluency; 10 min of pooling and written record; 28 min of essential exercises; 5 min of correction; 3 min of exit ticket.
2 h session: add 20 min of investigation, 20 min of formalisation, a second independent exercise and 10 min of comparative correction.
Differentiation: give struggling students a four-box sheet (data, model, calculation, conclusion). Ask quick students to produce a counterexample, a justification or a second method.

Session 25 — Derivative and direction of variation

Reciprocal (inverse) function

5 Jan 2027 • B: 2 h – C: 1 h

Session objectives

- Differentiate the reciprocal function.
- Justify the variations.
- Draw up a table.

Fluency – 6 to 8 minutes, no calculator

1. Write the multiplier for an increase of 10 %. $1 + \frac{10}{100} = 1.10$
2. Differentiate $3x^2 - 5x + 1$. $6x - 5$
3. Solve $(x-2)(x+1) = 0$. $x = -1$ **or** $x = 2$
4. Give the excluded value of $\frac{x+1}{x}$. $x = 0$
5. Calculate $\frac{1}{7}$ and $\frac{1}{-7}$. $\frac{1}{7}$ **and** $-\frac{1}{7}$

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: differentiate the reciprocal function and justify the variations.

Investigation

Starting from the difference quotient of $f(x) = \frac{1}{x}$, recover $f'(x) = -\frac{1}{x^2}$. Study its sign and check consistency with the graph.

Detailed solution – teacher edition

The square x^2 is positive for $x \neq 0$, so the derivative is strictly negative on each interval of the domain. Have students say aloud which piece of data triggers the choice of method.

Written record

For $x \neq 0$, $f'(x) = -\frac{1}{x^2} < 0$. The reciprocal function decreases on each of the intervals $]-\infty, 0[$ and $]0, +\infty[$.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Detailed solution – teacher edition

The square x^2 is positive for $x \neq 0$, so the derivative is strictly negative on each interval of the domain.

Practice – essential

1. Give the domain and the asymptotes of $f(x) = 2 + \dfrac{4}{x}$.
2. Calculate $f'(x)$ and study its sign.
3. Study the function $g(x) = x + \dfrac{4}{x}$ on $]0, +\infty[$.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$; asymptotes $x = 0$ and $y = 2$. $f'(x) = -\dfrac{4}{x^2} < 0$. For g, $g'(x) = 1 - \dfrac{4}{x^2}$ and the minimum is reached at $x = 2$, with value 4.

Textbook references

Further reading: Calao, course pp. 89–93; exercises p. 94, no. 1–3, then p. 96, no. 10–13. The exercise reproduced in this session is identified below by its page and number; the reference also lets you find the figures and digital resources of the textbook.

Calao textbook – p. 94, exercise 3

Calculate the derivatives of the following functions defined on $\mathbb{R}^*$.

a. $f(x) = 2x - 3 + \dfrac{4}{x}$

b. $g(x) = 3x - 5 - \dfrac{1}{x}$

c. $f(x) = 7x^3 - 2x + \dfrac{1}{x}$

d. $g(x) = 3x^2 - 5x - 2 + \dfrac{4}{x}$

Detailed solution – teacher edition

The derivatives are: $f'(x) = 2 - 4/x^2$; $g'(x) = 3 + 1/x^2$; $f'(x) = 21x^2 - 2 - 1/x^2$; $g'(x) = 6x - 5 - 4/x^2$.

Homework

Redo practice exercise 2 with the following data: replace the initial value by 245 and the rate or common ratio by 3. Write it up in four lines: data, formula, calculation, conclusion.

Detailed solution – teacher edition

The answer depends on the exercise revisited. Check above all the consistency of the model, the number of iterations, the rounding and the concluding sentence.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1, 2 and 3.

Method

1 h session: 7 min of fluency; 10 min of pooling and written record; 28 min of essential exercises; 5 min of correction; 3 min of exit ticket.
2 h session: add 20 min of investigation, 20 min of formalisation, a second independent exercise and 10 min of comparative correction.
Differentiation: give struggling students a four-box sheet (data, model, calculation, conclusion). Ask quick students to produce a counterexample, a justification or a second method.

Session 26 — Combinations with polynomial functions

Reciprocal (inverse) function

7 Jan 2027 • B: 1 h – C: 2 h

Session objectives

- Differentiate a combination of functions.
- Simplify an expression.
- Organise a function study.

Fluency – 6 to 8 minutes, no calculator

1. Simplify $3x + 8 - x + 2$. $2x + 10$
2. Solve $(x - 2)(x + 1) = 0$. $x = -1$ **or** $x = 2$
3. Give the excluded value of $\dfrac{x+1}{x}$. $x = 0$
4. Calculate $\dfrac{1}{8}$ and $\dfrac{1}{-8}$. $\dfrac{1}{8}$ **and** $-\dfrac{1}{8}$
5. Differentiate $3x^2 - 5x + 1$. $6x - 5$

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: differentiate a combination of functions and simplify an expression.

Investigation

A company produces $x > 0$ hundred items and its average cost is $C(x) = 2x + 30 + \dfrac{200}{x}$. Calculate $C'(x)$, factorise its numerator and find the production level that minimises the average cost.

Detailed solution – teacher edition

$C'(x) = 2 - \dfrac{200}{x^2} = \dfrac{2(x-10)(x+10)}{x^2}$. The minimum is reached for $x = 10$, that is 1 000 items. Have students say aloud which piece of data triggers the choice of method.

Written record

To study $P(x) + \dfrac{k}{x}$, find the domain, calculate the derivative, put over a common denominator, study its sign, then draw up the variations.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Detailed solution – teacher edition

$C'(x) = 2 - \dfrac{200}{x^2} = \dfrac{2(x-10)(x+10)}{x^2}$. The minimum is reached for $x = 10$, that is 1 000 items.

Practice – essential

1. Give the domain and the asymptotes of $f(x) = 3 + \dfrac{5}{x}$.
2. Calculate $f'(x)$ and study its sign.
3. Study the function $g(x) = x + \dfrac{9}{x}$ on $]0, +\infty[$.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$; asymptotes $x = 0$ and $y = 3$. $f'(x) = -\dfrac{5}{x^2} < 0$. For g, $g'(x) = 1 - \dfrac{9}{x^2}$ and the minimum is reached at $x = 3$, with value 6.

Textbook references

Further reading: Calao, p. 96, no. 14–21. The exercise reproduced in this session is identified below by its page and number; the reference also lets you find the figures and digital resources of the textbook.

Calao textbook – p. 96, exercise 20

Let f be the function defined on $\mathbb{R}^*$ by $f(x) = 4x + 1 + \dfrac{9}{x}$.

1. Show that for every nonzero real number x: $f'(x) = \dfrac{(2x-3)(2x+3)}{x^2}$.
2. For every nonzero real number x, study the sign of $f'(x)$.
3. Deduce the variations of the function f.

Detailed solution – teacher edition

$f'(x) = 4 - 9/x^2 = (4x^2 - 9)/x^2 = ((2x-3)(2x+3))/x^2$. The denominator is positive: f' is positive on $]-\infty; -3/2[\cup]3/2; +\infty[$ and negative on $]-3/2; 0[\cup]0; 3/2[$. We deduce the variations on each of the intervals separated by $-3/2$, 0 and $3/2$.

Homework

Redo practice exercise 3 with the following data: replace the initial value by 250 and the rate or common ratio by 4. Write it up in four lines: data, formula, calculation, conclusion.

Detailed solution – teacher edition

The answer depends on the exercise revisited. Check above all the consistency of the model, the number of iterations, the rounding and the concluding sentence.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1 and 2.

Method

1 h session: 7 min of fluency; 10 min of pooling and written record; 28 min of essential exercises; 5 min of correction; 3 min of exit ticket.
2 h session: add 20 min of investigation, 20 min of formalisation, a second independent exercise and 10 min of comparative correction.
Differentiation: give struggling students a four-box sheet (data, model, calculation, conclusion). Ask quick students to produce a counterexample, a justification or a second method.

Session 27 — Sign and factorisation

Reciprocal (inverse) function

12 Jan 2027 • B: 2 h – C: 1 h

Session objectives

- Factorise.
- Build a sign table.
- Respect excluded values.

Fluency – 6 to 8 minutes, no calculator

1. Calculate 20 % of 180. 36
2. Give the excluded value of $\dfrac{x+1}{x}$. $x = 0$
3. Calculate $\dfrac{1}{9}$ and $\dfrac{1}{-9}$. $\dfrac{1}{9}$ **and** $-\dfrac{1}{9}$
4. Differentiate $3x^2 - 5x + 1$. $6x - 5$
5. Solve $(x-2)(x+1) = 0$. $x = -1$ **or** $x = 2$

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: factorise and build a sign table.

Investigation

Study the sign of $\dfrac{(x-2)(x+3)}{x}$. Build the table with the three critical values, distinguish a zero from an excluded value, then solve the inequality $\dfrac{(x-2)(x+3)}{x} \geq 0$.

Detailed solution – teacher edition

The critical values are $-3, 0, 2$. The solution is $[-3, 0[\cup [2, +\infty[$. Have students say aloud which piece of data triggers the choice of method.

Written record

In a quotient, the sign depends on the numerator and the denominator. Excluded values are left out of the table, even if a factor formally cancels.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Detailed solution – teacher edition

The critical values are $-3, 0, 2$. The solution is $[-3, 0[\cup [2, +\infty[$.

Practice – essential

1. Give the domain and the asymptotes of $f(x) = 4 + \dfrac{6}{x}$.
2. Calculate $f'(x)$ and study its sign.
3. Study the function $g(x) = x + \dfrac{16}{x}$ on $]0, +\infty[$.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$; asymptotes $x = 0$ and $y = 4$. $f'(x) = -\dfrac{6}{x^2} < 0$. For g, $g'(x) = 1 - \dfrac{16}{x^2}$ and the minimum is reached at $x = 4$, with value 8.

Textbook references

Further reading: Calao, pp. 96–101. The exercise reproduced in this session is identified below by its page and number; the reference also lets you find the figures and digital resources of the textbook.

Calao textbook – p. 96, exercise 21

Let g be the function defined on $\mathbb{R}^*$ by $g(x) = x + 10 - \dfrac{1}{x}$.

1. Show that for every nonzero real number x: $g'(x) = \dfrac{x^2 + 1}{x^2}$.
2. For every nonzero real number x, study the sign of $g'(x)$.
3. Deduce the variations of the function g.

Detailed solution – teacher edition

$g'(x) = 1 + 1/x^2 = (x^2 + 1)/x^2 > 0$ for every $x \neq 0$. The function is therefore strictly increasing on each of the intervals $]-\infty; 0[$ and $]0; +\infty[$.

Homework

Redo practice exercise 2 with the following data: replace the initial value by 255 and the rate or common ratio by 5. Write it up in four lines: data, formula, calculation, conclusion.

Detailed solution – teacher edition

The answer depends on the exercise revisited. Check above all the consistency of the model, the number of iterations, the rounding and the concluding sentence.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1, 2 and 3.

Method

1 h session: 7 min of fluency; 10 min of pooling and written record; 28 min of essential exercises; 5 min of correction; 3 min of exit ticket.
2 h session: add 20 min of investigation, 20 min of formalisation, a second independent exercise

and 10 min of comparative correction.
Differentiation: give struggling students a four-box sheet (data, model, calculation, conclusion). Ask quick students to produce a counterexample, a justification or a second method.

Session 28 — Derivatives of polynomials of degree at most 3

Reciprocal (inverse) function

14 Jan 2027 • B: 1 h – C: 2 h

Session objectives

- Differentiate a polynomial.
- Study a derivative.
- Deduce the variations.

Fluency – 6 to 8 minutes, no calculator

1. Write the multiplier for an increase of 25 %. $1 + \dfrac{25}{100} = 1.25$
2. Calculate $\dfrac{1}{3}$ and $\dfrac{1}{-3}$. $\dfrac{1}{3}$ **and** $-\dfrac{1}{3}$
3. Differentiate $3x^2 - 5x + 1$. $6x - 5$
4. Solve $(x-2)(x+1) = 0$. $x = -1$ **or** $x = 2$
5. Give the excluded value of $\dfrac{x+1}{x}$. $x = 0$

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: differentiate a polynomial and study a derivative.

Investigation

For $f(x) = 2x^3 - 3x^2 - 12x + 5$, calculate $f'(x)$, factorise, study its sign and deduce the variations of f.

Detailed solution – teacher edition

$f'(x) = 6x^2 - 6x - 12 = 6(x-2)(x+1)$. The function increases on $]-\infty, -1]$, decreases on $[-1, 2]$, then increases. Have students say aloud which piece of data triggers the choice of method.

Written record

$(ax^3 + bx^2 + cx + d)' = 3ax^2 + 2bx + c$. The sign of the derivative gives the direction of variation of the function.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Detailed solution – teacher edition

$f'(x) = 6x^2 - 6x - 12 = 6(x-2)(x+1)$. The function increases on $]-\infty, -1]$, decreases on $[-1, 2]$, then increases.

Practice – essential

1. Give the domain and the asymptotes of $f(x) = 5 + \dfrac{7}{x}$.
2. Calculate $f'(x)$ and study its sign.
3. Study the function $g(x) = x + \dfrac{25}{x}$ on $]0, +\infty[$.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$; asymptotes $x = 0$ and $y = 5$. $f'(x) = -\dfrac{7}{x^2} < 0$. For g, $g'(x) = 1 - \dfrac{25}{x^2}$ and the minimum is reached at $x = 5$, with value 10.

Textbook references

Further reading: Calao, pp. 94–96. The exercise reproduced in this session is identified below by its page and number; the reference also lets you find the figures and digital resources of the textbook.

Calao textbook – p. 94, exercise 13

Let f be the function defined on $\mathbb{R}^*$ by: $f(x) = \dfrac{1}{x}$.

1. Calculate $f'(x)$.
2. Show that for every nonzero real number x: $f'(x) = \dfrac{-1}{x^2}$.
3. After studying the sign of $f'(x)$, draw up the table of variations of f.

Detailed solution – teacher edition

$f'(x) = -1/x^2 < 0$ for every $x \neq 0$. The reciprocal function is therefore strictly decreasing on each of the intervals $]-\infty; 0[$ and $]0; +\infty[$. These two intervals must not be merged into a single interval of decrease.

Homework

Redo practice exercise 3 with the following data: replace the initial value by 260 and the rate or common ratio by 6. Write it up in four lines: data, formula, calculation, conclusion.

Detailed solution – teacher edition

The answer depends on the exercise revisited. Check above all the consistency of the model, the number of iterations, the rounding and the concluding sentence.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1 and 2.

Method

1 h session: 7 min of fluency; 10 min of pooling and written record; 28 min of essential exercises; 5 min of correction; 3 min of exit ticket.
2 h session: add 20 min of investigation, 20 min of formalisation, a second independent exercise

and 10 min of comparative correction.
Differentiation: give struggling students a four-box sheet (data, model, calculation, conclusion). Ask quick students to produce a counterexample, a justification or a second method.

Session 29 — Full study and graph

Reciprocal (inverse) function

19 Jan 2027 • B: 2 h – C: 1 h

Session objectives

- Carry out a full study.
- Draw a consistent curve.
- Interpret an extremum.

Fluency – 6 to 8 minutes, no calculator


1. Simplify $3x + 4 - x + 2$. $2x + 6$
2. Differentiate $3x^2 - 5x + 1$. $6x - 5$
3. Solve $(x - 2)(x + 1) = 0$. $x = -1$ **or** $x = 2$
4. Give the excluded value of $\dfrac{x+1}{x}$. $x = 0$
5. Calculate $\dfrac{1}{4}$ and $\dfrac{1}{-4}$. $\dfrac{1}{4}$ **and** $-\dfrac{1}{4}$

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: carry out a full study and draw a consistent curve.

Investigation

Study $g(x) = x + \dfrac{9}{x}$ completely on $]0, +\infty[$: domain, derivative, sign, variations, minimum, behaviour at the boundaries and sketch.



Detailed solution – teacher edition

$g'(x) = \dfrac{(x-3)(x+3)}{x^2}$; minimum $g(3) = 6$; $g(x)$ tends to $+\infty$ at 0^+ and at $+\infty$. Have students say aloud which piece of data triggers the choice of method.

Written record

A full study includes: domain, derivative, sign, variations, limits or behaviour at the boundaries, key points and a consistent graph.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Detailed solution – teacher edition

$g'(x) = \dfrac{(x-3)(x+3)}{x^2}$; minimum $g(3) = 6$; $g(x)$ tends to $+\infty$ at 0^+ and at $+\infty$.

Practice – essential

1. Give the domain and the asymptotes of $f(x) = 6 + \dfrac{8}{x}$.
2. Calculate $f'(x)$ and study its sign.
3. Study the function $g(x) = x + \dfrac{36}{x}$ on $]0, +\infty[$.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$; asymptotes $x = 0$ and $y = 6$. $f'(x) = -\dfrac{8}{x^2} < 0$. For g, $g'(x) = 1 - \dfrac{36}{x^2}$ and the minimum is reached at $x = 6$, with value 12.

Textbook references

Further reading: Calao, pp. 98–103. The exercise reproduced in this session is identified below by its page and number; the reference also lets you find the figures and digital resources of the textbook.

Calao textbook – p. 98, exercise 31

A company specialising in industrial ovens can manufacture between 20 and 80 of them depending on the year. Let x be the number of ovens manufactured per year.
The graph given in the textbook is that of a function C defined on the interval $[20; 80]$. For every integer value of x, $C(x)$ is the average unit production cost (in €).

1. What is the average unit production cost when 25 ovens are produced, then when 70 ovens are produced?
2. Which production levels correspond to an average unit cost of 20 000 €?
3. What is the minimum average unit production cost? Which production level does it correspond to?

Detailed solution – teacher edition

The numerical answers are read from the curve on p. 98. Require reading lines and units. The graphical minimum must be identified by the ordinate of the lowest point, then linked to its abscissa; a graphical reading remains approximate.

Homework

Redo practice exercise 2 with the following data: replace the initial value by 265 and the rate or common ratio by 7. Write it up in four lines: data, formula, calculation, conclusion.

Detailed solution – teacher edition

The answer depends on the exercise revisited. Check above all the consistency of the model, the number of iterations, the rounding and the concluding sentence.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1, 2 and 3.

Method

1 h session: 7 min of fluency; 10 min of pooling and written record; 28 min of essential exercises; 5 min of correction; 3 min of exit ticket.
2 h session: add 20 min of investigation, 20 min of formalisation, a second independent exercise and 10 min of comparative correction.
Differentiation: give struggling students a four-box sheet (data, model, calculation, conclusion). Ask quick students to produce a counterexample, a justification or a second method.

Session 30 — Review and assessment

Reciprocal (inverse) function

21 Jan 2027 • B: 1 h – C: 2 h

Session objectives

- Use domain, derivative and sign.
- Write up a study.
- Correct work methodically.

Fluency – 6 to 8 minutes, no calculator

1. Calculate 5 % of 100. 5
2. Solve $(x-2)(x+1)=0$. $x=-1$ **or** $x=2$
3. Give the excluded value of $\dfrac{x+1}{x}$. $x=0$
4. Calculate $\dfrac{1}{5}$ and $\dfrac{1}{-5}$. $\dfrac{1}{5}$ **and** $-\dfrac{1}{5}$
5. Differentiate $3x^2-5x+1$. $6x-5$

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: use domain, derivative and sign and write up a study.

Investigation

Correct a deliberately flawed study of $h(x)=x+\dfrac{4}{x}$: stated domain $\mathbb{R}$, derivative $1+\dfrac{4}{x^2}$, table with no break at 0. Rewrite a complete solution.

Detailed solution – teacher edition

The domain is $\mathbb{R}\setminus\{0\}$ and $h'(x)=1-\dfrac{4}{x^2}$. The table is split at 0 and takes into account -2 and 2. Have students say aloud which piece of data triggers the choice of method.

Written record

Check the four items separately: domain, expression of the derivative, sign of the derivative, variations. An error at one step must not hide the following ones.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$ and $h'(x) = 1 - \dfrac{4}{x^2}$. The table is split at 0 and takes into account -2 and 2.

Practice – essential

1. Give the domain and the asymptotes of $f(x) = 2 + \dfrac{4}{x}$.
2. Calculate $f'(x)$ and study its sign.
3. Study the function $g(x) = x + \dfrac{4}{x}$ on $]0, +\infty[$.

Detailed solution – teacher edition

The domain is $\mathbb{R} \setminus \{0\}$; asymptotes $x = 0$ and $y = 2$. $f'(x) = -\dfrac{4}{x^2} < 0$. For g, $g'(x) = 1 - \dfrac{4}{x^2}$ and the minimum is reached at $x = 2$, with value 4.

Textbook references

Further reading: Calao, pp. 100–103. The exercise reproduced in this session is identified below by its page and number; the reference also lets you find the figures and digital resources of the textbook.

Calao textbook – p. 98, exercise 32

Each week, a liquid detergent company estimates that the production cost (in euros) can be modelled by a function C given by: $C(x) = x^2 + 60x + 121$, where x is the volume of detergent (in m^3), $x \in [1;30]$.
Let f be the function representing the average production cost per m^3 of liquid detergent.

1. Show that for every $x \in [1;30]$, $f(x) = x + 60 + \dfrac{121}{x}$.
2. Calculate $f'(x)$. Show that for every $x \in [1;30]$, $f'(x) = \dfrac{(x-11)(x+11)}{x^2}$.
3. Draw up the table of variations of f.
4. What is the minimum average production cost? For what quantity of detergent is it reached?

Detailed solution – teacher edition

$f(x) = C(x)/x = x + 60 + 121/x$. Then $f'(x) = 1 - 121/x^2 = ((x-11)(x+11))/x^2$. On $[1;30]$, f' is negative up to 11, then positive. The minimum is reached for $x = 11$ and equals $f(11) = 82$ € per m^3.

Homework

Redo practice exercise 3 with the following data: replace the initial value by 270 and the rate or common ratio by 2. Write it up in four lines: data, formula, calculation, conclusion.

Detailed solution – teacher edition

The answer depends on the exercise revisited. Check above all the consistency of the model, the number of iterations, the rounding and the concluding sentence.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1 and 2.

Method

1 h session: 7 min of fluency; 10 min of pooling and written record; 28 min of essential exercises; 5 min of correction; 3 min of exit ticket.
2 h session: add 20 min of investigation, 20 min of formalisation, a second independent exercise and 10 min of comparative correction.
Differentiation: give struggling students a four-box sheet (data, model, calculation, conclusion). Ask quick students to produce a counterexample, a justification or a second method.