Course notes

Area and antiderivative

For f continuous and positive on $[a, b]$, $a \leq b$, the integral $\int_a^b f(x)\,\mathrm{d}x$ is the area under the curve, above the axis, between $x = a$ and $x = b$, expressed in area units.

Fundamental theorem

If f is continuous on an interval I and $a \in I$, then $F_a(x) = \int_a^x f(t)\,\mathrm{d}t$ is the antiderivative of f that vanishes at a. For every antiderivative F,

$$\int_a^b f(x)\,\mathrm{d}x = F(b) - F(a).$$

Every continuous function on an interval has antiderivatives. An integral is a number; an antiderivative is a function.

Proof in the positive increasing case. For $h > 0$, the rectangles give $hf(x) \leq F_a(x+h) - F_a(x) \leq hf(x+h)$. After dividing by h and taking the limit, the rate of change tends to $f(x)$. For $h < 0$, the increasing property gives $f(x+h) \leq f(t) \leq f(x)$ on $[x+h, x]$. Integrating,

$$(-h)f(x+h) \leq F_a(x) - F_a(x+h) \leq (-h)f(x).$$

Dividing by $-h > 0$, we obtain

$$f(x+h) \leq \frac{F_a(x+h) - F_a(x)}{h} \leq f(x).$$

Both bounds tend to $f(x)$. By continuity, $F_a' = f$. Every other antiderivative is $F_a + C$, hence the formula for evaluating between the bounds.

Properties

For continuous functions and bounds belonging to the same interval:

$$\int_a^a f = 0, \quad \int_b^a f = -\int_a^b f, \quad \int_a^c f = \int_a^b f + \int_b^c f, \quad \int_a^b (\lambda f + \mu g) = \lambda \int_a^b f + \mu \int_a^b g.$$

If $a \leq b$ and $f \leq g$, then $\int_a^b f \leq \int_a^b g$. In particular, if $m \leq f \leq M$, then $m(b-a) \leq \int_a^b f \leq M(b-a)$. For a function of arbitrary sign, the integral is defined through antiderivatives. The geometric area between the curve and the axis is $\int_a^b |f|$, obtained by splitting at the sign changes. Between two curves, it is $\int_a^b |f - g|$.

Example. $\int_{-1}^{1} x \, \mathrm{d}x = 0$, but the area between the line and the axis on this interval is 1.

Mean value and integration by parts

For $a < b$, the mean value is $\mu = \frac{1}{b-a} \int_a^b f(x) \, \mathrm{d}x$. The rectangle of height μ and width $b - a$ has the same algebraic area.

Integration by parts

If u, v are of class $\mathcal{C}^1$ on $[a, b]$,

$$\int_a^b u(x)v'(x) \, \mathrm{d}x = [u(x)v(x)]_a^b - \int_a^b u'(x)v(x) \, \mathrm{d}x.$$


Proof. Integrate $(uv)' = u'v + uv'$, then isolate the required integral. The square bracket is indispensable.

Example. With $u = x$ and $v' = e^x$, $\int_0^1 xe^x \, \mathrm{d}x = [xe^x]_0^1 - \int_0^1 e^x \, \mathrm{d}x = e - (e - 1) = 1$.

Rectangles and sequences of integrals

For f continuous and increasing on $[a, b]$, with $a < b$ and $n \geq 1$ and $h = \frac{b-a}{n}$, the sums $h \sum_{k=0}^{n-1} f(a + kh)$ and $h \sum_{k=1}^{n} f(a + kh)$ bound the integral. Their difference is $h(f(b) - f(a))$. For $I_n = \int_0^1 x^n f(x) \, \mathrm{d}x$, begin with the sign and comparisons of the integrand. If $0 \leq f \leq M$, then $0 \leq I_n \leq \frac{M}{n+1}$; the squeeze theorem gives $I_n \to 0$. An integration by parts can also provide a relation between I_n and I_{n+1}.

Convention at rank zero. In $I_n = \int_0^1 x^n f(x) \, \mathrm{d}x$, the expression x^0 denotes the constant function 1, including at the lower bound 0. Thus $I_0 = \int_0^1 f(x) \, \mathrm{d}x$; no indeterminate calculation is needed.



The area between $y = x$ and $y = x^2$ on $[0, 1]$ is $\frac{1}{6}$.