

The Facets of Geometry

Categorical

(Under heavy construction!!)

July 15, 2024

Contents

I	The Algebraic Viewpoint	1
1	Foundational Algebraic Geometry	3
1.1	A guiding example	5
1.2	Affine schemes and basic properties	6
1.3	Schemes and basic properties	18
1.4	First notions on schemes	25
1.5	Varieties	34
1.6	Fundamental constructions on schemes	55
1.7	Dimension of schemes	74
1.8	Projective schemes	79
1.9	\mathcal{O}_X -modules	86
1.10	Divisors	100
1.11	Smoothness & differential forms	109
1.12	Morphism of schemes	113
1.13	Coherent and quasicoherent sheaf cohomology	135
2	Varieties over an algebraically closed field	137
2.1	Notations	137
2.2	Intersection with hypersurfaces	139
2.3	Grassmannians	140
3	Elliptic Curves	141
4	Étale topology	143
5	Deformation Theory	145
6	Algebraic Geometry	147
6.1	Functor of points	147
6.2	Analytification and GAGA	147
6.3	Intersection theory	147
6.4	Overview of K-theory of schemes	149
6.5	The Riemann-Hilbert correspondence	150
6.6	Serre's intersection formula	151

II	The Arithmetic Viewpoint	153
7	Foundational Arithmetic	155
7.1	Fundamental properties of \mathbb{Z}	155
7.2	Algebraic number fields	155
III	The Topological Viewpoint	159
8	Foundational Geometry	161
8.1	Locally ringed spaces and manifolds	161
8.2	Linearization	169
8.3	Constructions on manifolds	169
8.4	Lie groups	169
8.5	Global algebra	170
8.6	Torsors and 1 st -Čech cohomology group	186
8.7	Bundles	187
8.8	Differential forms and de-Rham cohomology	188
9	Foundational Differential Geometry	193
9.1	Bundles in differential geometry and applications	193
9.2	Cohomological methods	193
9.3	Covariant derivative, connections, classes and curvatures	193
10	Foundational Homotopy Theory	195
10.1	Fundamental group and covering maps	199
10.2	Cofibrations and cofiber sequences	222
10.3	Fibrations and fiber sequences	233
10.4	Homology theories	246
10.5	Cohomology theories	261
10.6	Cohomology products and duality	262
10.7	CW-complexes & CW homotopy types	263
10.8	Homotopy and homology	264
10.9	Homotopy & algebraic structures	264
10.10	Model categories & abstract homotopy	266
10.11	Classifying spaces	266
10.12	Spectra	266
10.13	Lifting & extension problems	266
11	Stable Homotopy Theory	267
12	Classical Ordinary Differential Equations	269
12.1	Initial value problems	270
12.2	Linear systems	279
12.3	Stability of linear systems in \mathbb{R}^2	282
12.4	Autonomous systems	283

12.5 Linearization and flow analysis	285
12.6 Second order ODE	291
13 K-Theory of Vector Bundles	297
14 Jet Bundles	299
IV The Analytic Viewpoint	301
15 Analysis on Complex Plane	303
15.1 Holomorphic functions	304
15.2 La théorie des cartes holomorphes	311
15.3 Singularities	325
15.4 Cauchy's theorem - II	327
15.5 Residues and meromorphic maps	329
15.6 Riemann mapping theorem	331
16 Riemann Surfaces	333
16.1 Introduction	333
16.2 Ramified coverings & Riemann-Hurwitz formula	342
16.3 Monodromy & analytic continuation	343
16.4 Holomorphic & meromorphic forms	344
16.5 Riemann-Roch theorem	349
17 Foundational Analytic Geometry	359
V The Categorical Viewpoint	361
18 Classical Topoi	365
18.1 Towards the axioms of a Topos	366
18.2 Grothendieck Topologies & Sheaves	371
18.3 Basic Properties and Results in Topoi	381
18.4 Sheaves in an arbitrary Topos	401
18.5 Geometric Morphisms	415
18.6 Categorical Semantics	421
18.7 Topoi and Logic	431
19 Language of ∞-Categories	443
19.1 Simplicial sets	443
19.2 Classical homotopical algebra	458
20 Algèbre Commutative Dérivée	461

VI Special Topics	463
23 Commutative Algebra	471
23.1 General algebra	473
23.2 Graded rings & modules	487
23.3 Noetherian modules and rings	490
23.4 $\text{Supp}(M)$, $\text{Ass}(M)$ and primary decomposition	493
23.5 Tensor, symmetric & exterior algebras	496
23.6 Field theory	508
23.7 Integral dependence and normal domains	543
23.8 Dimension theory	551
23.9 Completions	554
23.10 Valuation rings	555
23.11 Dedekind domains	558
23.12 Tor and Ext functors	560
23.13 Projective and injective modules	561
23.14 Multiplicities	566
23.15 Kähler differentials	567
23.16 Depth, Cohen-Macaulay & regularity	570
23.17 Filtrations	572
23.18 Flatness	572
23.19 Lifting properties : Étale maps	573
23.20 Lifting properties : Unramified maps	574
23.21 Lifting properties : Smooth maps	575
23.22 Simple, semisimple and separable algebras	576
23.23 Miscellaneous	578
24 K-Theory of Rings	591
24.1 K_0	591
24.2 K_1	603
24.3 K_2	610
24.4 Higher K -theory of rings-I	619
24.5 K -theory & étale cohomology	636
25 Abstract Analysis	639
25.1 Integration theory	640
25.2 Banach spaces	709
25.3 Hilbert spaces	725
25.4 Extension problems-I : Hahn-Banach theorem	726
25.5 Major theorems : UBP, OMT, BIT, CGT	726
25.6 Strong & weak convergence	735
25.7 Spectral theory	737
25.8 Compact operators	737

26 Homological Methods	739
26.1 The setup : abelian categories	739
26.2 Homology, resolutions and derived functors	741
26.3 Results for $\mathbf{Mod}(R)$	750
27 Foundational Sheaf Theory	751
27.1 Recollections	751
27.2 The sheafification functor	752
27.3 Morphisms of sheaves	754
27.4 Sheaves are étale spaces	760
27.5 Direct and inverse image	764
27.6 Category of sheaves	767
27.7 Classical Čech cohomology	773
27.8 Derived functor cohomology	778

Part V

The Categorical Viewpoint

Chapter 19

Language of ∞ -Categories

Contents

19.1 Simplicial sets	443
19.1.1 Extension of functor by colimits-I	444
19.1.2 Categories Δ and \mathbf{sSet}	446
19.1.3 Operations on simplicial sets	449
19.1.4 Basic properties	450
19.1.5 Eilenberg-Zilber categories	452
19.1.6 Kan complexes and homotopy groups	453
19.1.7 The fundamental group of a Kan complex	457
19.1.8 ∞ -categories	458
19.1.9 Theorem of Boardman-Vogt	458
19.2 Classical homotopical algebra	458
19.2.1 Model categories	458
19.2.2 Basic constructions	459
19.2.3 Construction of model structures on $\mathbf{PSh}(\mathbf{C})$	459
19.2.4 From one model structure to many	459

In this chapter we give an overview of the language of ∞ -categories. The ultimate goal of this chapter is to define the ∞ -category of ∞ -groupoids and state the Yoneda lemma in ∞ -categorical setting. Therefore, if you believe that ∞ -groupoids are spaces (homotopy hypothesis), then we would construct the ∞ -category of spaces by the end of this chapter. In the process, we will learn about the techniques involved in manipulating ∞ -categories. In this chapter, we will only work with the category of compactly generated spaces and by **Top** we mean category of compactly generated spaces.

19.1 Simplicial sets

The goal of simplicial sets is to obtain a combinatorial approximation of topological spaces. We describe simplicial sets as a presheaf over simplex category. One can motivate themselves why this

definition is the correct definition for combinatorially handling topological spaces by reading the review paper cite[Bergner’s simplicial set paper]. We give some basic properties of such objects. We also give a general important result about presheaves, ”every presheaf is a colimit of representable presheaves”. This is of fundamental importance in the development of ∞ -categories.

Remark 19.1.0.1. (*Notations*)

1. As we will be frequently constructing and dealing with presheaf category over a category \mathbf{C} , therefore instead of only denoting it by $\mathbf{PSh}(\mathbf{C})$, we will also be denoting it by $\widehat{\mathbf{C}}$, depending on the convenience of the given situation.
2. We will denote objects of an ordinary 1-category \mathbf{C} by lowercase alphabets like a, b, c, \dots, x, y, z , morphisms in \mathbf{C} by lowercase letters like f, g, h, \dots and functors $\mathbf{C} \rightarrow \mathbf{D}$ by uppercase alphabets like A, B, C, \dots, X, Y, Z and also by lowercase alphabets like f, g, h, \dots .
3. Let \mathbf{C} be a category and $c \in \mathbf{C}$ be an object. We denote by $h_c : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ the contravariant hom-functor given by $a \mapsto \text{Hom}_{\mathbf{C}}(a, c)$.
4. We will denote ∞ -categories by $\mathcal{A}, \mathcal{B}, \mathcal{C}, \dots, \mathcal{X}, \mathcal{Y}, \mathcal{Z}$.

Let us also as a reminder put the usual Yoneda lemma.

Lemma 19.1.0.2. (*Yoneda lemma*) *Let \mathbf{A} be a category, $a \in \mathbf{A}$ be an object and F be a presheaf over \mathbf{A} . Then, there is a natural isomorphism*

$$\text{Hom}_{\widehat{\mathbf{A}}}(h_a, F) \xrightarrow{\cong} F(a).$$

Proof. Consider the following maps

$$\begin{aligned} \varphi : \text{Hom}_{\widehat{\mathbf{A}}}(h_a, F) &\longrightarrow F(a) \\ \alpha &\longmapsto \alpha_a(\text{id}_a). \end{aligned}$$

and

$$\begin{aligned} \xi : F(a) &\longrightarrow \text{Hom}_{\widehat{\mathbf{A}}}(h_a, F) \\ x &\longmapsto \beta : h_a \rightarrow F \end{aligned}$$

where β is defined as follows. Consider any $a' \in \mathbf{A}$ and any $f \in h_a(a')$. Define $\beta_{a'} : h_a(a') \rightarrow F(a')$ by mapping as $f \mapsto F(f)(x)$.

One then sees that $\xi \circ \varphi = \text{id}$ by naturality of α and that $\varphi \circ \xi = \text{id}$ by functoriality of F . \square

Corollary 19.1.0.3. *Let \mathbf{A} be a category and define $\text{Yon} : \mathbf{A} \rightarrow \widehat{\mathbf{A}}$ to be the Yoneda functor given by $a \mapsto h_a$ and $a \rightarrow b \mapsto h_a \rightarrow h_b$. Then, Yon is fully-faithful.*

Proof. We have by Yoneda lemma (Lemma 19.1.0.2) that $\text{Hom}_{\widehat{\mathbf{A}}}(h_a, h_b) \cong h_b(a) = \text{Hom}_{\mathbf{A}}(a, b)$. \square

19.1.1 Extension of functor by colimits-I

The following is a fundamental result in category theory.

Theorem 19.1.1.1. *Let \mathbf{A} be a small category and \mathbf{C} be a locally small category with all small colimits. Then, for any functor*

$$f : \mathbf{A} \rightarrow \mathbf{C}$$

we get two functors

$$f_! : \widehat{\mathbf{A}} \rightarrow \mathbf{C}$$

and

$$\begin{aligned} f^* : \mathbf{C} &\rightarrow \widehat{\mathbf{A}} \\ c &\mapsto \text{Hom}_{\mathbf{C}}(f(-), c) \end{aligned}$$

such that

1. $f_!$ is left adjoint of f^*

$$\widehat{\mathbf{A}} \begin{array}{c} \xrightarrow{f_!} \\ \perp \\ \xleftarrow{f^*} \end{array} \mathbf{C} .$$

The functor $f_!$ is called the extension of f by colimits.

2. *Denoting $\text{Yon} : \mathbf{A} \hookrightarrow \widehat{\mathbf{A}}$ to be the Yoneda embedding, we get that the following commutes upto a unique natural isomorphism*

$$\begin{array}{ccc} \widehat{\mathbf{A}} & \xrightarrow{f_!} & \mathbf{C} \\ \text{Yon} \uparrow & \searrow f & \\ \mathbf{A} & & \end{array} .$$

That is, $f_!(h_a) \cong f(a)$ where $h_a = \text{Hom}_{\mathbf{A}}(-, a)$.

Proof. The main heart of the proof is the fact that every presheaf is a colimit of representables (Proposition 1.1.8 of cite[Cis]).

1. We define $f_! : \widehat{\mathbf{A}} \rightarrow \mathbf{C}$ as follows. Pick any $X \in \widehat{\mathbf{A}}$. Consider the functor $\varphi_X : \mathbf{A}/X \rightarrow \widehat{\mathbf{A}}$ from the category of elements of X given by $(a, s) \mapsto h_a$ and on maps by $f : (a, s) \rightarrow (b, t)$ by $h_f : h_a \rightarrow h_b$. By Proposition 1.1.8 of cite[Cis], we have $\varinjlim_{(a,s)} h_a = X$. Define $f_!(X) = \varinjlim_{(a,s)} f(a)$ which exists in \mathbf{C} as \mathbf{C} has all small colimits. Consequently, we obtain the following natural isomorphisms by Yoneda lemma (Lemma 19.1.0.2) and limit preserving properties of contravariant homs

$$\begin{aligned} \text{Hom}_{\mathbf{C}}(f_!(X), c) &\cong \text{Hom}_{\mathbf{C}}\left(\varinjlim_{(a,s)} f(a), c\right) \cong \varprojlim_{(a,s)} \text{Hom}_{\mathbf{C}}(f(a), c) \\ &\cong \varprojlim_{(a,s)} \text{Hom}_{\widehat{\mathbf{A}}}(h_a, f^*(c)) \cong \text{Hom}_{\widehat{\mathbf{A}}}\left(\varinjlim_{(a,s)} h_a, f^*(c)\right) \\ &\cong \text{Hom}_{\widehat{\mathbf{A}}}(X, f^*(c)). \end{aligned}$$

2. Since $\text{Hom}_{\mathbf{C}}(f_!(h_a), c) \cong \text{Hom}_{\widehat{\mathbf{A}}}(h_a, f^*(c)) \cong f^*(c)(a) = \text{Hom}_{\mathbf{C}}(f(a), c)$ for all objects $c \in \mathbf{C}$, therefore the result follows by Corollary 19.1.0.3. \square

The above theorem will take a central place in constructions of this chapter. Indeed, let us point the following applications of this theorem before stating its proof.

Corollary 19.1.1.2. *Let \mathbf{A} be a small category and \mathbf{C} be a category with small colimits. If $F : \widehat{\mathbf{A}} \rightarrow \mathbf{C}$ is a colimit preserving functor. Then*

1. *there exists $f : \mathbf{A} \rightarrow \mathbf{C}$ such that $f_! \cong F$ naturally,*
2. *F has a right adjoint.*

Proof. We first define the required f . For any object $a \in \mathbf{A}$, define $f(a) := F(h_a)$. Then by Theorem 19.1.1.1, 1, it follows that $f_!(h_a) \cong f(a) = F(h_a)$. Since we know that every presheaf is presented as a colimit of representable functors indexed by its category of elements, thus we get that for any $X \in \widehat{\mathbf{A}}$, $F(X) \cong f_!(X)$ naturally. The second conclusion also follows from Theorem 19.1.1.1, 2. \square

The following result will be important in order to define simplicial mapping spaces.

Proposition 19.1.1.3. *For any small category \mathbf{A} , the presheaf category $\widehat{\mathbf{A}}$ is Cartesian closed where the internal hom object is defined by*

$$\underline{\mathrm{Hom}}(X, Y)(a) := \mathrm{Hom}_{\widehat{\mathbf{A}}}(X \times h_a, Y)$$

and on morphisms as

$$\underline{\mathrm{Hom}}(X, Y)(f) := \mathrm{Hom}_{\widehat{\mathbf{A}}}(X \times h_f, Y).$$

Proof. We wish to show that $\underline{\mathrm{Hom}}(-, -)$ acts as internal hom object in $\widehat{\mathbf{A}}$. This can be seen by establishing following natural bijections

$$\mathrm{Hom}_{\widehat{\mathbf{A}}}(T \times X, Y) \cong \mathrm{Hom}_{\widehat{\mathbf{A}}}(T, \underline{\mathrm{Hom}}(X, Y))$$

This follows from contemplating the functor $f_X : \mathbf{A} \rightarrow \widehat{\mathbf{A}}$ for all $X \in \widehat{\mathbf{A}}$ given by $a \mapsto X \times \mathrm{Yon}(-)$, together with Theorem 19.1.1.1. Indeed, first observe that $\widehat{\mathbf{A}}$ is locally small with all small colimits. Second, observe from the proof of Theorem 19.1.1.1 that for any $Z \in \widehat{\mathbf{A}}$, we have that $f_{X!} : \widehat{\mathbf{A}} \rightarrow \widehat{\mathbf{A}}$ takes any $Z \in \widehat{\mathbf{A}}$ and maps it to $f_{X!}(Z) = \varinjlim_{(a,s)} f_X(a)$ where (a, s) varies over \mathbf{A}/Z , the category of elements of Z . Since $f_X(a) = X \times h_a$ and filtered colimits commute with finite limits, therefore we get a natural isomorphism $f_{X!}(Z) \cong X \times Z$. Consequently, the adjunction of Theorem 19.1.1.1 completes the proof. \square

19.1.2 Categories Δ and \mathbf{sSet}

Consider the category of all finite sets $[n]$ with $n + 1$ elements with linear order $0 < 1 < \dots < n$ and mappings being the non-decreasing maps. Denote this category by Δ and call it the *simplex category*. A *simplicial set* is then a presheaf over Δ . Let the category of all simplicial sets be denoted \mathbf{sSet} .

There are two important class of maps in Δ .

Definition 19.1.2.1 (Face and degeneracy maps). For each $n \in \mathbb{N}$, we have $n + 1$ face maps

$$d^i : [n - 1] \rightarrow [n]$$

$$j \mapsto \begin{cases} j & \text{if } j \leq i \\ j + 1 & \text{if } j > i \end{cases}$$

where $0 \leq i \leq n$ and n degeneracy maps

$$s^i : [n] \rightarrow [n - 1]$$

$$j \mapsto \begin{cases} j & \text{if } j \leq i \\ j - 1 & \text{if } j > i \end{cases}$$

where $0 \leq i \leq n - 1$.

Remark 19.1.2.2. By Yoneda embedding, we will allow ourselves to abuse the notation by writing $d^i : [n - 1] \rightarrow [n]$ as the unique map $d^i : \Delta^{n-1} \rightarrow \Delta^n$ and $s^i : [n] \rightarrow [n - 1]$ as the unique map $s^i : \Delta^n \rightarrow \Delta^{n-1}$ in **sSet**. For a simplicial set X we may thus interpret $x \in X_n$ as $x : \Delta^n \rightarrow X$. The above give maps which we denote as

$$d_i : X_n \longrightarrow X_{n-1}$$

$$x \longmapsto x \circ d_i$$

for each $0 \leq i \leq n$ also called the face maps and

$$s_i : X_{n-1} \longrightarrow X_n$$

$$x \longmapsto x \circ s_i$$

for each $0 \leq i \leq n - 1$ also called degeneracy maps.

It is quite easy to observe, but very important for applications, the following relations satisfied by face and degeneracy maps. All these are immediate from definition given above.

Proposition 19.1.2.3. *The following relations hold in Δ (and thus in **sSet**):*

1. *If $i < j$, then $d^j d^i = d^i d^{j-1}$.*
2. *If $i \leq j$, then $s^j s^i = s^i s^{j+1}$.*
3. *We have*

$$s^j d^i = \begin{cases} d^i s^{j-1} & \text{if } i < j \\ \text{id} & \text{if } i = j, j + 1 \\ d^{i-1} s^j & \text{if } j + 1 < i. \end{cases}$$

□

Consequently, for a simplicial set, dual relations hold.

Proposition 19.1.2.4. *Let X be a simplicial set. Then the face and degeneracy maps of X satisfies the following relations:*

1. *If $i < j$, then $d_i d_j = d_{j-1} d_i$.*

2. If $i \leq j$, then $s_i s_j = s_{j+1} s_i$.
3. We have

$$d_i s_j = \begin{cases} s_{j-1} d_i & \text{if } i < j \\ \text{id} & \text{if } i = j, j+1 \\ s_j d_{i-1} & \text{if } j+1 < i. \end{cases}$$

□

Remark 19.1.2.5. One may keep the following picture in mind while working with a simplicial set X :

$$\begin{array}{ccccc} [0] & \begin{array}{c} \rhd \xrightarrow{d^0} d^1 \xrightarrow{\quad} \\ \ll \xleftarrow{\quad} s^0 \end{array} & [1] & \begin{array}{c} \rhd \xrightarrow{d^0} d^1 \xrightarrow{d^2} \\ \ll \xleftarrow{s^1} s^0 \end{array} & [2] & \begin{array}{c} \rhd \xrightarrow{d^i} \\ \vdots \\ \ll \xleftarrow{s^i} \end{array} & \dots \\ & & & & & & \\ X_0 & \begin{array}{c} \xleftarrow{\quad} d_1 \xleftarrow{d_0} \\ \xrightarrow{-s_0} \end{array} & X_1 & \begin{array}{c} \xleftarrow{\quad} d_2 \xleftarrow{d_1} \\ \xrightarrow{-s_0} \xrightarrow{s_1} \end{array} & X_2 & \begin{array}{c} \xleftarrow{d_i} \\ \vdots \\ \xrightarrow{s_i} \end{array} & \dots \end{array}$$

Remark 19.1.2.6. There is a functor

$$\begin{aligned} |-| : \mathbf{\Delta} &\longrightarrow \mathbf{Top} \\ [n] &\longmapsto |\Delta^n| \end{aligned}$$

where $|\Delta^n|$ is the standard topological n -simplex in \mathbb{R}^{n+1} and for $f : [n] \rightarrow [m]$ in $\mathbf{\Delta}$, we have

$$\begin{aligned} |f| : |\Delta^n| &\rightarrow |\Delta^m| \\ (t_0, \dots, t_n) &\longmapsto \left(\sum_{i \in f^{-1}(0)} i, \dots, \sum_{i \in f^{-1}(m)} i \right). \end{aligned}$$

Example 19.1.2.7 (*Singular chains*). An important example of a simplicial set is that of $\text{Sing}(X)$ defined as

$$\begin{aligned} \text{Sing}(X) : \mathbf{\Delta}^{op} &\rightarrow \mathbf{Set} \\ [n] &\mapsto \text{Hom}_{\mathbf{Top}}(|\Delta^n|, X). \end{aligned}$$

The main point is that $\text{Sing}(X)$ as a simplicial set knows all about the homotopy type of space X . Consequently, we will denote $X([n]) := \text{Sing}(X)_n = \text{Hom}_{\mathbf{Top}}(|\Delta^n|, X)$.

Example 19.1.2.8 (*Nerve of a category*). Let \mathbf{C} be a category. Define the nerve of \mathbf{C} as

$$\begin{aligned} N\mathbf{C} : \mathbf{\Delta}^{op} &\rightarrow \mathbf{Set} \\ [n] &\mapsto \text{Hom}_{\mathbf{Cat}}([n], \mathbf{C}) \end{aligned}$$

where $[n]$ is a category as it is a poset. Consequently, $N\mathbf{C}_n$ is the set of all n -composable arrows of \mathbf{C} .

Theorem 19.1.2.9. *Nerve construction is a fully-faithfull embedding of categories into simplicial sets.*

19.1.3 Operations on simplicial sets

Define product, coproducts, subspaces, unions, quotients, limits, colimits and mapping objects. **TODO!!**

Example 19.1.3.1 (Standard Δ^n , boundaries $\partial_i \Delta^n$, $\partial \Delta^n$ and horn Λ_i^n). Denote $\Delta^n = N[n]$ to be the nerve of the category $[n]$. That is,

$$\Delta^n([m]) = \Delta_m^n = \text{Hom}_{\mathbf{\Delta}}([m], [n]) = h_{[n]}([m])$$

which are exactly all representable presheaves over $\mathbf{\Delta}$. These are the combinatorial analogues of the topological n -simplex $|\Delta^n|$ and we tend to think about Δ^n using the intuition gained from the topological one. There are some important simplicial subsets of Δ^n .

Let $E \subseteq [n]$ be a totally ordered subset. Define $\Delta^E = NE$ to be a simplicial subset of Δ^n . From this we derive the following simplicial subsets of Δ^n . The first is the i^{th} -boundary of Δ^n for $0 \leq i \leq n$ given by

$$\partial_i \Delta^n = \bigcup_{i \notin E \subsetneq [n]} \Delta^E \cong \Delta^{n-1}.$$

The second is the *boundary* of Δ^n given by

$$\partial \Delta^n = \bigcup_{E \subsetneq [n]} \Delta^E = \bigcup_{i=0}^n \partial_i \Delta^n$$

The third is the i^{th} -horn of Δ^n denoted Λ_i^n given by

$$\Lambda_i^n = \bigcup_{i \in E \subsetneq [n]} \Delta^E.$$

Remark 19.1.3.2. Let $n \geq 1$, $0 \leq i \leq n$ and $m \geq 0$. Note that we have

$$\begin{aligned} (\partial_i \Delta^n)_m &= \left\{ \begin{array}{l} \text{Order preserving maps } f : [m] \rightarrow [n] \\ \text{which are not surjective and } i \notin \text{Im}(f). \end{array} \right\} \\ (\partial \Delta^n)_m &= \left\{ \begin{array}{l} \text{Order preserving maps } f : [m] \rightarrow [n] \\ \text{which are not surjective.} \end{array} \right\} \\ (\Lambda_i^n)_m &= \left\{ \begin{array}{l} \text{Order preserving maps } f : [m] \rightarrow [n] \\ \text{which are not surjective and } i \in \text{Im}(f). \end{array} \right\}. \end{aligned}$$

For two simplicial sets, we can define the internal hom using the Proposition [19.1.1.3](#).

Definition 19.1.3.3 (Homotopy & mapping complex). Let S, T be a simplicial set. Then $\underline{\text{Hom}}(S, T)$ denotes the following simplicial set

$$[n] \mapsto \text{Hom}_{\mathbf{sSet}}(S \times \Delta^n, T).$$

An n -simplex of $\underline{\text{Hom}}(S, T)$ is defined to be an n -homotopy from S to T . A 1-homotopy H is also referred to as a homotopy from $H|_{S \times \{0\}} =: f$ to $H|_{S \times \{1\}} =: g$.

19.1.4 Basic properties

Do results and exercises from various sources to showcase the combinatorial arguments used while working with simplicial sets.

We discuss some properties of simplicial sets which would be useful later on. Some of these might be taken as exercises on combinatorial manipulations with simplicial sets.

We first begin with a simple observation.

Definition 19.1.4.1 (*n -degenerate*). A simplicial set X is said to be n -degenerate if for all $m > n$, all m -simplices in X_m are degenerate.

Example 19.1.4.2. Each standard simplicial sets Δ^n are n -degenerate. Indeed, its m -simplices for $m > n$ are

$$\Delta_m^n = \{\text{Order preserving maps } f : [m] \rightarrow [n]\}.$$

But as $m > n$, therefore every such f is necessarily non-injective. It follows that each simplex in Δ_m^n is in the image of $s_i : X_{m-1} \rightarrow X_m$ for some $0 \leq i \leq m-1$.

For similar reasons, the i^{th} -boundary $\partial_i \Delta^n$, boundary $\partial \Delta^n$ and horns Λ_i^n for $0 \leq i \leq n$ are all $n-1$ -degenerate.

Lemma 19.1.4.3. *Let X be an n -degenerate simplicial set and Y be a simplicial set. Then any collection of functions $\{\varphi_m : X_m \rightarrow Y_m\}_{0 \leq m \leq n}$ such that for any $f : [k] \rightarrow [l]$ in Δ with $0 \leq k, l \leq n$, the following square commutes*

$$\begin{array}{ccc} X_l & \xrightarrow{\varphi_l} & Y_l \\ f^* \downarrow & & \downarrow f^* \\ X_k & \xrightarrow{\varphi_k} & Y_k \end{array},$$

the collection $\{\varphi_m\}_{0 \leq m \leq n}$ lifts to a unique map of simplicial sets $\varphi : X \rightarrow Y$.

Remark 19.1.4.4. As a consequence of Lemma 19.1.4.3, in order to give a map of simplicial sets from an n -degenerate simplicial set S , it suffices to construct the required map only on m -simplices for $0 \leq m \leq n$.

Proof of Lemma 19.1.4.3. Let $\{\varphi_m\}_{0 \leq m \leq n}$ be as given. We wish to define $\varphi_{m'}$ for $m' > n$. We proceed by induction on m' . Suppose $\varphi_{m'-1}$ is given to us. Since we have the following diagram

$$\begin{array}{ccc} X_{m'-1} & \xrightarrow{\varphi_{m'-1}} & Y_{m'-1} \\ s_i \downarrow & & \downarrow s_i \\ X_{m'} & \xrightarrow{\varphi_{m'}} & Y_{m'} \end{array}$$

and that every element of $X_{m'}$ is in image of s_i (guaranteed by Proposition 19.1.2.4, 3), it follows that there is a unique choice of $\varphi_{m'}$ to fit in the above diagram, as required. \square

Lemma 19.1.4.5. *Let $n \geq 1$ and $0 \leq i \leq n$. Then,*

1. $\partial_i \Delta^n$ has exactly 1 non-degenerate $n-1$ -simplex,

2. $\partial\Delta^n$ has exactly $n + 1$ non-degenerate $n - 1$ -simplices,
3. Λ_i^n has exactly n non-degenerate $n - 1$ -simplices.

Proof. These three items are immediate from Remark 19.1.3.2. □

The following is an important adjunction which will be consistently used in later sections. Moreover, it is generally good to keep in mind all the time while working with simplicial sets so that one transfer intuition from topological spaces to that of simplicial sets, as we would need to do time and time again (for example when dealing with homotopy of simplicial sets).

Theorem 19.1.4.6 (*Geometric realization*). *The singular functor $\text{Sing} : \mathbf{Top} \rightarrow \mathbf{sSet}$ has a left adjoint $|-| : \mathbf{sSet} \rightarrow \mathbf{Top}$*

$$\mathbf{sSet} \begin{array}{c} \xrightarrow{|-|} \\ \xleftarrow[\text{Sing}]{\perp} \end{array} \mathbf{Top} .$$

The functor $|-|$ is called the *geometric realization* and for a simplicial set X , we have

$$|X| = \coprod_{n \geq 0} X_n \times |\Delta^n| / \sim$$

where \sim is generated by $(f^*x, t) \sim (x, |f|t)$ for all $f : [n] \rightarrow [m]$ in Δ , $x \in X_m$ and $t \in |\Delta^n|$.

Proof. The main idea is that any map $X \rightarrow \text{Sing}(Y)$ is a natural transform of presheaves. One observes that the naturality conditions on this morphism is equivalently represented in terms of a map $|X| \rightarrow Y$. **TODO.** □

Example 19.1.4.7. As an example, we calculate $\Delta^1 \times \Delta^1$ and show that

$$\Delta^1 \times \Delta^1 \cong \Delta^2 \cup_{\Delta^1} \Delta^2.$$

TODO.

Observe that even though Δ^1 has all simplices of dimension ≥ 2 as degenerate, yet $\Delta^1 \times \Delta^1$ has two non-degenerate 2-dimensional simplices.

A consequence of the above isomorphism is that the geometric realization of $\Delta^1 \times \Delta^1$ is exactly I^2 , the unit square, which is the product of the geometric realization of Δ^1 with itself. Indeed, this is an instantiation of the general result in Theorem 19.1.4.9.

Remark 19.1.4.8. It is immediate to observe from Theorem 19.1.4.6 that geometric realization of Δ^n is exactly the standard topological n -simplex. Similarly, $\partial\Delta^n$ and $|\Lambda_i^n|$ are homeomorphic to exactly the pictures that we used in our mind to understand them.

Finally, the main result is as follows.

Theorem 19.1.4.9. *Let X, Y be simplicial sets. Then the natural map*

$$|X \times Y| \rightarrow |X| \times |Y|$$

is a homeomorphism.

Proof. **TODO.** □

We show that nerve of a category satisfies lifting property which would prove to be useful later on while discussing ∞ -categories.

Proposition 19.1.4.10. *Let $X = N\mathbf{C}$ be the nerve of a small category \mathbf{C} . Then for any $0 < i < n$, the following lifting problem is uniquely filled:*

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & \nearrow \text{!} & \\ \Delta^n & & \end{array} .$$

Proof. **TODO.** □

The following is an important result which will find its use later on.

Proposition 19.1.4.11. *Let S be a simplicial set and X a Kan complex. Then $\underline{\text{Hom}}(S, X)$ is a Kan complex, denoted $\text{Map}(S, X)$.*

Proof. **TODO.** □

19.1.5 Eilenberg-Zilber categories

This is an abstraction of the type of combinatorial proofs that we would like to make in the simplex category Δ . Indeed, as we would have to work with surjections and injections in Δ primarily, which interacts with the size of $[n]$ (which we would define to be n in a minute) as an injection would only increase the size and a surjection would only decrease it, therefore we need a systematic toolset to work with these things. In particular, if we denote Δ_+ to be the subcategory of all injective maps, Δ_- to be the subcategory of all surjective maps in Δ and $d : \text{Ob}(\Delta) \rightarrow \mathbb{N}$ the size map, then we have the following properties about the tuple $(\Delta, \Delta_+, \Delta_-, d)$:

1. All bijections are in both Δ_+ and Δ_- .
2. The dimension map d takes bijective cardinals to the same natural.
3. Let $\sigma : [n] \rightarrow [m]$ in Δ which is not bijective. If σ is injective (i.e. in Δ_+), then $d(a) < d(b)$ and if σ is surjective (i.e. in Δ_-), then $d(a) > d(b)$.
4. Any map $\sigma : [n] \rightarrow [m]$ in Δ factors as a surjection followed by an injection.
5. For any surjective map $\sigma : [n] \rightarrow [m]$ in Δ , there exists a section $\pi : [m] \rightarrow [n]$, i.e. such that $\pi\sigma = \text{id}_{[m]}$.
6. If $\sigma : [n] \rightarrow [m]$ is surjective, then the set of sections of σ uniquely determines the map σ .

Remark 19.1.5.1. We need this abstraction of properties of Δ so that with the same techniques, we can work with bisimplicial sets, which is important if one wishes to consider simplicial objects in \mathbf{sSet} .

These considerations about Δ motivates the following definition.

Definition 19.1.5.2. (Eilenberg-Zilber categories) A category \mathbf{A} is said to be an Eilenberg-Zilber category (or much simply, *EZ category*), if there exists subcategories \mathbf{A}_+ , \mathbf{A}_- and a function $d : \text{Ob}(\mathbf{A}) \rightarrow \mathbb{N}$ which satisfies the following axioms:

1. All isomorphisms of \mathbf{A} are in both \mathbf{A}_+ and \mathbf{A}_- .
2. If a, a' are isomorphic objects in \mathbf{A} , then $d(a) = d(a')$.

3. Let $\sigma : a \rightarrow a'$ not be an isomorphism. If σ is in \mathbf{A}_+ , then $d(a) < d(a')$. If σ is in \mathbf{A}_- , then $d(a) > d(a')$.
4. For any map $\sigma : a \rightarrow a'$ in \mathbf{A} , there exists unique factorization of σ into maps $p : a \rightarrow c$ in \mathbf{A}_- and $i : c \rightarrow a'$ in \mathbf{A}_+

$$\begin{array}{ccc} & c & \\ p \nearrow & & \searrow i \\ a & \xrightarrow{\sigma} & a' \end{array} .$$

5. If $\sigma : a \rightarrow a'$ is a map in \mathbf{A}_- , then there exists a section $\pi : a' \rightarrow a$, i.e. $\pi\sigma = \text{id}_a$.
6. If σ, σ' are two maps in \mathbf{A}_- such both of them has the same collection of sections, then $\sigma = \sigma'$.

The main thrust of this section is to discuss presheaves over an EZ category \mathbf{A} , keeping in mind the prototypical case of $\mathbf{A} = \mathbf{\Delta}$. Indeed, the main result and its corollaries will serve first as a practice for the type of arguments we shall need later and also as a tool to be consistently used later in constructions with simplicial sets (which are, presheaves over $\mathbf{\Delta}$).

Example 19.1.5.3. By previous discussion, it is clear that the simplex category $\mathbf{\Delta}$ is an EZ category.

Let \mathbf{A} be an EZ category and X a presheaf over \mathbf{A} . Then the category of elements of X , \mathbf{A}/X , is an EZ category again. Indeed, define $(\mathbf{A}/X)_+$ exactly as those pairs (a, s) where $a \in \mathbf{A}_+$ and $(\mathbf{A}/X)_-$ exactly as those pairs (a, s) where $a \in \mathbf{A}_-$. Further for an object (a, s) , define $d(a, s) = d(a)$, where the latter d is coming from the EZ structure on \mathbf{A} .

19.1.6 Kan complexes and homotopy groups

In the beginning section, we showed how simplicial sets can be viewed as combinatorial version of usual spaces. However, in order to do homotopy theory in this combinatorial setting, we need to isolate a class of simplicial sets which are right for this kind. Indeed, these will be Kan complexes.

We wish to define $\pi_n(X, x)$ where $x \in X_0$ is a 0-simplex of a simplicial set. To this end, we immediately run into problems as $\pi_0(X, x)$ should be the equivalence class of all those 0-simplices which are boundaries of a 1-simplex. But it is immediately clear that this is not an equivalence relation! Indeed, consider Δ^1 . Then the above relation is not symmetric as we have $0 \rightarrow 1$ as a 1-simplex but there is no $1 \rightarrow 0$ in Δ^1 . Similarly, if we try to prove transitivity of the above relation, we land up in the following situation. Let $x \rightarrow y, y \rightarrow z$ be two 1-simplices in X . Then, we wish to find a 1-simplex $x \rightarrow z$ in X . Note that given $x \rightarrow y$ and $y \rightarrow z$, we have a map $\Lambda_1^2 \rightarrow X$ and we wish to fill the following diagram

$$\begin{array}{ccc} \Lambda_1^2 & \longrightarrow & X \\ \downarrow & \nearrow \text{dashed} & \\ \Delta^2 & & \end{array} .$$

So we need those simplicial sets, where the above dotted arrow always exists.

Definition 19.1.6.1 (Kan complex). A simplicial set X is a Kan complex if for any $n \geq 0$ and

any $0 \leq i \leq n$, the following lifting diagram is filled

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & \nearrow & \\ \Delta^n & & \end{array}.$$

Example 19.1.6.2. Clearly, Δ^n are not Kan complexes for $n \geq 0$. Moreover, $N\mathbf{C}$ is a Kan complex if and only if \mathbf{C} is a groupoid. This is immediate.

The prototypical (and in some sense, the only example) of Kan complexes are those obtained by Theorem 19.1.4.6.

Proposition 19.1.6.3. *Let X be a topological space. Then $\text{Sing}(X)$ is a Kan complex.*

Proof. Let $n \geq 0$, $0 \leq i \leq n$. Then a map $\Lambda_i^n \rightarrow \text{Sing}(X)$ is equivalent to a map $|\Lambda_i^n| \rightarrow X$ by adjunction of Theorem 19.1.4.6. Consequently, we wish to fill the following diagram

$$\begin{array}{ccc} |\Lambda_i^n| & \longrightarrow & X \\ \downarrow & \nearrow & \\ |\Delta^n| & & \end{array}.$$

But this is immediate as $|\Lambda_i^n| \hookrightarrow |\Delta^n|$ is a retraction. \square

Another example of a Kan complex is the mapping complex.

Definition 19.1.6.4 (Mapping complex). Let X, Y be two spaces. The mapping complex, denoted $\text{Map}(X, Y)$, is the one whose n -simplices are

$$[n] \mapsto \text{Hom}_{\mathbf{Top}}(X \times |\Delta^n|, Y).$$

Note that $\text{Map}(X, Y)$ has 0-simplices as continuous maps, 1-simplices as homotopies and so on. The name is justified by the following result.

Corollary 19.1.6.5. *Let X, Y be spaces. Then the mapping complex $\text{Map}(X, Y)$ is a Kan complex.*

Proof. As all spaces are compactly generated, therefore we have $\text{Hom}_{\mathbf{Top}}(X \times |\Delta^n|, Y) \cong \text{Hom}_{\mathbf{Top}}(|\Delta^n|, Y^X)$. Consequently, $\text{Map}(X, Y) \cong \text{Sing}(Y^X)$, which is a Kan complex by Proposition 19.1.6.3. \square

One can now check that the relation mentioned in the beginning on X_0 is indeed an equivalence relation and would thus yield the definition of $\pi_0(X, x)$, however, we would like to define all homotopy groups in one go. We first define the notion of two simplices being homotopic.

Definition 19.1.6.6 (Homotopic rel ∂). Let X be a Kan complex and $\sigma, \tau \in X_n$ be two n -simplices. Then σ and τ are homotopic rel ∂ if there exists an $n+1$ -simplex $\psi \in X_{n+1}$ such that $\sigma|_{\partial\Delta^n} = \tau|_{\partial\Delta^n}$ and $\partial_n\psi = \sigma$, $\partial_{n+1}\psi = \tau$ and for all $0 \leq i \leq n-1$, $\partial_i\psi = \sigma d^i s^{n-1} = \tau d^i s^{n-1}$. We can diagrammatically represent these conditions as the commutativity of the diagrams

$$\begin{array}{ccc} X & \xleftarrow{\tau} & \Delta^n \\ \sigma \uparrow & & \uparrow \\ \Delta^n & \xleftarrow{\quad} & \partial\Delta^n \end{array},$$

$$\begin{array}{ccccc}
\Delta^{n+1} & \xrightarrow{\psi} & X & \xleftarrow{\psi} & \Delta^{n+1} \\
d^n \uparrow & \nearrow \sigma & & \nwarrow \tau & \uparrow d^{n+1} \\
\Delta^n & & & & \Delta^n
\end{array}$$

and for $0 \leq i \leq n-1$

$$\begin{array}{ccccc}
\Delta^{n+1} & \xrightarrow{\psi} & X & \xleftarrow{\tau} & \Delta^n \\
d^i \uparrow & & & & \uparrow d^i \\
\Delta^n & \xrightarrow{s^{n-1}} & \Delta^{n-1} & &
\end{array}$$

in **sSet**.

Indeed, we see that this generalizes our previous notion of homotopy relative to boundary as follows.

Lemma 19.1.6.7. *Let $X = \text{Sing}(Y)$ be the Kan complex associated to a space Y . Then two n -simplices $\sigma, \tau \in X_n$ are homotopic rel ∂ if and only if $\sigma, \tau : |\Delta^n| \rightarrow Y$ are homotopic relative to $|\partial\Delta^n| \hookrightarrow |\Delta^n|$ in the classical sense.*

Proof. Let $\sigma, \tau : \Delta^n \rightarrow X$ be two n -simplices. These are homotopic rel ∂ if the diagrams in Definition 19.1.6.6 commutes in **sSet**. By the adjunction of Theorem 19.1.4.6, this is equivalent to commutativity of the following diagrams in **Top**

$$\begin{array}{ccc}
Y & \xleftarrow{\tau} & |\Delta^n| \\
\sigma \uparrow & & \uparrow \\
|\Delta^n| & \xleftarrow{\quad} & |\partial\Delta^n|
\end{array},$$

$$\begin{array}{ccccc}
|\Delta^{n+1}| & \xrightarrow{\psi} & X & \xleftarrow{\psi} & |\Delta^{n+1}| \\
d^n \uparrow & \nearrow \sigma & & \nwarrow \tau & \uparrow d^{n+1} \\
|\Delta^n| & & & & |\Delta^n|
\end{array}$$

and for $0 \leq i \leq n-1$

$$\begin{array}{ccccc}
|\Delta^{n+1}| & \xrightarrow{\psi} & Y & \xleftarrow{\tau} & |\Delta^n| \\
d^i \uparrow & & & & \uparrow d^i \\
|\Delta^n| & \xrightarrow{s^{n-1}} & |\Delta^{n-1}| & &
\end{array}.$$

But this is equivalent to the data of a homotopy $H : |\Delta^n| \times I \rightarrow Y$ rel boundary. Indeed, since we want $H_0 = \sigma$, $H_1 = \tau$ and $H_t|_{|\partial\Delta^n|} = \sigma|_{|\partial\Delta^n|} = \tau|_{|\partial\Delta^n|}$ for all $t \in I$, therefore we naturally get that H factors through $|\Delta^{n+1}|$. This can be visualized for a 3-simplex immediately. \square

Finally, when X is a Kan complex, then this is an equivalence relation. This is where combinatorial relations between face and degeneracy maps of a simplicial set as provided in Proposition 19.1.2.3 becomes handy.

Proposition 19.1.6.8. *Let X be a Kan complex. The homotopy rel ∂ is an equivalence relation on each X_n for $n \geq 1$.*

Proof. We first show reflexivity. Consider $\sigma : \Delta^n \rightarrow X$ an n -simplex. We claim that $\psi = \sigma s^n : \Delta^{n+1} \rightarrow X$ works as the homotopy from σ to σ . Indeed, we see that $\psi d^n = \sigma s^n d^n = \sigma$ and $\psi d^{n+1} = \sigma s^n d^{n+1} = \sigma$ by Proposition 19.1.2.3. Similarly, for $0 \leq i \leq n-1$, we have $\psi d^i = \sigma s^n d^i = \sigma d^i s^{n-1}$ by the same result. This establishes reflexivity.

We now show symmetry. Let $\sigma \sim \tau$ for some $\sigma, \tau : \Delta^n \rightarrow X$ with $\sigma|_{\partial \Delta^n} = \tau|_{\partial \Delta^n}$. Then, there exists $\psi : \Delta^{n+1} \rightarrow X$ with $\psi d^n = \sigma$, $\psi d^{n+1} = \tau$ and $\psi d^i = \sigma d^i s^{n-1} = \tau d^i s^{n-1}$ for $0 \leq i \leq n-1$. We wish to show that $\tau \sim \sigma$. We will use the fact that X is a Kan complex (so that all horns can be filled). In particular, we will construct a horn $\kappa : \Lambda_n^{n+2} \rightarrow X$, filling which will give us the required homotopy from τ to σ . Indeed, let κ be obtained from ψ by adding degeneracies such that $\kappa d^{n+2} = \psi$ and $\kappa d^{n+1} = \sigma s^n$, that is, the degenerate $n+1$ -simplex obtained by repeating the n^{th} -vertex of σ . For $0 \leq i \leq n-1$, we keep $\kappa d^i = \psi d^i s^{n-1}$. Filling the horn κ yields an $n+2$ -simplex $\tilde{\kappa} : \Delta^{n+2} \rightarrow X$ whose n^{th} -face is exactly a homotopy from τ to σ . Indeed, denote $\phi = \tilde{\kappa} d^n$. Then, $\phi d^n = \tilde{\kappa} d^n d^n = \tilde{\kappa} d^{n+1} d^n = \sigma$. Similarly, $\phi d^{n+1} = \tau$. These follow from the observation that $d^n d^n$ is the unique n -simplex of $\tilde{\kappa}$ not containing the vertices n and $n+1$ in $\tilde{\kappa}$. For $0 \leq i \leq n-1$, by Proposition 19.1.2.3, we have $\phi d^i = \tilde{\kappa} d^n d^i = \tilde{\kappa} d^i d^{n-1}$, which in turn is $\kappa d^i d^{n-1} = \psi d^i s^{n-1} d^{n-1} = \psi d^i = \sigma d^i s^{n-1}$, as needed. This shows symmetry.

Finally, we wish to show transitivity. Let $\psi, \psi' \in X_{n+1}$ be homotopies $\sigma \sim \tau$ and $\tau \sim \eta$ respectively for some $\sigma, \tau, \eta \in X_n$. We wish to construct a homotopy $\psi'' \in X_{n+1}$ between σ and η . Indeed, we obtain a horn $\kappa : \Lambda_{n+2}^{n+2} \rightarrow X$ whose n^{th} and $n+1^{\text{th}}$ boundaries are ψ and ψ' respectively and the rest boundaries are required degeneracies. Filling this horn up by the Kan condition gives $\tilde{\kappa}$ and its $n+2^{\text{th}}$ -boundary is the required homotopy. \square

Consequently, we define homotopy groups of a Kan complex as follows.

Definition 19.1.6.9 (Homotopy groups of a Kan complex). Let X be a Kan complex and $x \in X_0$ be a base point. Then for $n \geq 1$ define

$$\pi_n(X, x_0) = \{\sigma \in X_n \mid \sigma|_{\partial \Delta^n} = c_{x_0}\} / \sim$$

where \sim is the homotopy rel ∂ . For $n = 0$, define

$$\pi_0(X) = X_0 / \sim$$

where

$$x \sim y \iff \exists \gamma \in X_1 \text{ s.t. } \gamma d^1 = x \text{ \& } \gamma d^0 = y.$$

We now show that $\pi_n(X, x)$ is indeed a group.

Construction 19.1.6.10 (Composition and group operation on $\pi_n(X, x)$). Let X be a Kan complex and $x \in X$ be a point in it (i.e a 0-simplex). Let $\sigma, \tau \in X_n$ be two n -simplices such that $\sigma|_{\partial \Delta^n} = \tau|_{\partial \Delta^n} = c_x$. We construct the composition $\sigma \cdot \tau$ of σ and τ as the following n -simplex. Construct the following horn $\kappa : \Lambda_n^{n+1} \rightarrow X$ whose $n-1^{\text{th}}$ -boundary is σ , $n+1^{\text{th}}$ -boundary is τ^1 and for

¹If one is thinking of paths, then it is important to note that $\sigma \cdot \tau$ is the one where τ is traversed first and then σ . One has to let go of the past notation because here simplices are not merely going to be paths, homotopies and so on, but rather more general objects like arrows of a category, 2-arrows and so on, so to be consistent with the notion of composition, it is best that we change the order in which we concatenate paths.

$0 \leq i \leq n-2$, $\kappa d^i = c_x$. It follows from horn-filling condition of X that we get an $n+1$ -simplex $\psi : \Delta^{n+1} \rightarrow X$ extending the horn κ . Consequently, we define the *concatenation* $\sigma \cdot \tau$ as the n^{th} -boundary of ψ , that is,

$$\sigma \cdot \tau = \psi d^n.$$

We claim that the operation

$$\begin{aligned} \cdot : \pi_n(X, x) \times \pi_n(X, x) &\longrightarrow \pi_n(X, x) \\ ([\alpha], [\beta]) &\longmapsto [\alpha \cdot \beta] \end{aligned}$$

is a well-defined function, that is, $[\alpha \cdot \beta]$ only depends on $[\alpha]$ and $[\beta]$.

Indeed, suppose $[\alpha] = [\alpha']$ and $[\beta] = [\beta']$ in $\pi_n(X, x)$. Then, we have a homotopy rel ∂ denoted $\psi \in X_{n+1}$ from α to α' and $\chi \in X_{n+1}$ from β to β' . We wish to construct $\phi \in X_{n+1}$ which is a homotopy rel ∂ from $\alpha \cdot \beta$ to $\alpha' \cdot \beta'$. As usual, we obtain this homotopy by constructing a higher horn and filling it by Kan condition.

To correctly denote the simplex to be constructed, we first observe that if $\delta \sim \epsilon$ is a homotopy rel ∂ of two n -simplices with $\delta|_{\partial\Delta^n} = \epsilon|_{\partial\Delta^n} = x$, then $\delta \cdot x \sim \epsilon$. Indeed, consider an $n+2$ -horn whose $n+2$ -boundary is the composition simplex $\delta \cdot x$, $n+1$ -boundary is ϵs_0 and all i -boundaries for $0 \leq i \leq n-1$ are degeneracies of δ . Filling this yields the n -boundary as the required homotopy.

We now show that if $\beta \sim \beta'$, then $\alpha \cdot \beta \sim \alpha \cdot \beta'$. This would complete the proof. Indeed, this is immediate by considering an $n+2$ -horn given by $\kappa : \Lambda_{n+1}^{n+2} \rightarrow X$ such that κd^{n+2} is the homotopy $\beta \sim \beta'$, κd^n is the composition $\alpha \cdot \beta$, κd^{n-1} is the composition $\alpha \cdot \beta'$ and the κd^i for $0 \leq i \leq n-2$ are all x .

Consequently, we get that \cdot is a well defined operation on $\pi_n(X, x)$. We will later show that \cdot makes $\pi_n(X, x)$ into a group. Moreover, we will show that $\pi_n(X, x) \cong \pi_n(|X|, x)$ and $\pi_n(\text{Sing}(Y), y) \cong \pi_n(Y, y_0)$!

19.1.7 The fundamental group of a Kan complex

Let X be a Kan complex and $x_0 \in X$ be a 0-simplex. The fundamental group $\pi_1(X, x_0)$ is explicitly given by the following

$$\begin{aligned} \pi_1(X, x_0) &= \{\sigma : \Delta^1 \rightarrow X \mid \sigma|_{\partial\Delta^1} = x_0\} / \sim \\ &= \{\sigma \in X_1 \mid d_0(\sigma) = d_1(\sigma) = x_0\} / \sim \end{aligned}$$

where $\sigma \sim \tau$ if and only if there exists a homotopy rel ∂ denoted $H : \Delta^2 \rightarrow X$, from σ to τ . That is, $d_1(H) = \sigma$, $d_2(H) = \tau$ and $d_0(H) = s_0(d_0(\sigma)) = s_0(x_0) = \text{id}_{x_0}$. This is a group where the operation is

$$\begin{aligned} \cdot : \pi_1(X, x_0) \times \pi_1(X, x_0) &\longrightarrow \pi_1(X, x_0) \\ ([\sigma], [\tau]) &\longmapsto [\sigma \cdot \tau] \end{aligned}$$

where $\sigma \cdot \tau$ is a 1-simplex which is their composition (Construction 19.1.6.10), obtained by the 1-boundary of the 2-simplex δ obtained by filling the horn

$$\kappa : \Lambda_1^2 \rightarrow X$$

whose $\partial_0 \Lambda_1^2 = \sigma$ and $\partial_1 \Lambda_1^2 = \tau$. More precisely, we define κ_1 as follows (this is sufficient by Example 19.1.4.2 and Lemma 19.1.4.3):

$$\begin{aligned}\kappa_1 : (\Lambda_1^2)_1 &\longrightarrow X_1 \\ \{0, 1\} &\longmapsto \sigma \\ \{1, 2\} &\longmapsto \tau \\ \{1, 1\} &\longmapsto s_0(d_0(\sigma)).\end{aligned}$$

In this section, following the usual terminology, we will write $\sigma \cdot \tau$ as $\tau * \sigma$ for $\sigma, \tau \in X_1$ with $d_0(\sigma) = d_1(\tau) = x_0$.

We now prove some basic results about $\pi_1(X, x_0)$. It is a good exercise to show that $\pi_1(X, x_0)$ is a group.

Theorem 19.1.7.1. *Let X be a Kan complex and $x_0 \in X_0$. Then $\pi_1(X, x_0)$ is a group.*

Proof. We first show that for any three $\alpha, \beta, \gamma \in X_1$ with their boundaries being x_0 , we have $(\alpha * \beta) * \gamma \sim \alpha * (\beta * \gamma)$. Indeed, let H be the witness of composition $\alpha * \beta$, K that of $\beta * \gamma$, L that of $(\alpha * \beta) * \gamma$ and P that of $\alpha * (\beta * \gamma)$. Now consider the following horn $\kappa : \Lambda_2^3 \rightarrow X$ given by following on non-degenerate 2-simplices:

$$\begin{aligned}\kappa_2 : (\Lambda_2^3)_2 &\longrightarrow X_2 \\ \{0, 1, 2\} &\longmapsto H \\ \{1, 2, 3\} &\longmapsto K \\ \{0, 2, 3\} &\longmapsto L.\end{aligned}$$

By Kan condition, the above horn is filled and its 2-boundary yields a 2-simplex $\chi \in X_2$ such that $d_0(\chi) = \beta * \gamma$, $d_1(\chi) = (\alpha * \beta) * \gamma$ and $d_2(\chi) = \alpha$. We now construct the required homotopy by filling another 3-horn. Indeed, consider a horn $\lambda : \Lambda_1^3 \rightarrow X$ such that $d_0(\lambda) = s_1(\beta * \gamma)$, $d_2(\lambda) = P$ and $d_3(\lambda) = \chi$. This fills to give its 1-boundary as the required homotopy.

The identity element being c_{x_0} and the existence of inverses are also immediate results of horn filling and is thus omitted. \square

A Kan complex is path-connected if $\pi_0(X) = 0$.

19.1.8 ∞ -categories

19.1.9 Theorem of Boardman-Vogt

19.2 Classical homotopical algebra

19.2.1 Model categories

We discuss now a general setup in which one can "do" homotopy theory.

Definition 19.2.1.1 (Model categories). Let \mathbf{C} be a category and $W, C, F \subseteq \mathbf{C}$ be subcategories of \mathbf{C} which are called weak equivalences (\approx), cofibrations (\rightarrow) and fibrations (\twoheadrightarrow) respectively. We call $W \cap C$ weak/acyclic cofibrations and $W \cap F$ weak/acyclic fibrations. Then, the tuple (\mathbf{C}, W, C, F) is a model category if it satisfies the following axioms:

1. The category \mathbf{C} has all finite limits and colimits.
2. Weak equivalences satisfies 2 out-of 3 property.
3. For any $f : X \rightarrow Y$ in \mathbf{C} , we have two factorizations

$$\begin{array}{ccc}
 X & \xrightarrow{f} & Y \\
 \swarrow \approx & & \nearrow \\
 & Z &
 \end{array}
 \qquad
 \begin{array}{ccc}
 X & \xrightarrow{f} & Y \\
 \searrow & & \swarrow \approx \\
 & Z &
 \end{array}$$

one a weak cofibration followed by fibration and another one a cofibration followed by a weak fibration.

4. We have

$$\begin{aligned}
 rlp(W \cap C) &= F \\
 C &= llp(W \cap F)
 \end{aligned}$$

where for a subcategory $S \subseteq \mathbf{C}$, the collection $rlp(S)$ denotes the collection of all maps $X \rightarrow Y$ in \mathbf{C} satisfying right lifting property wrt S , i.e., such that for *any* commutative square

$$\begin{array}{ccc}
 Z & \longrightarrow & X \\
 \downarrow \in S & \nearrow h & \downarrow f \\
 W & \longrightarrow & Y
 \end{array}$$

where left vertical arrow is in S , there exists a lift $h : W \rightarrow X$ as shown which makes all diagrams commute. Similarly, one defines $llp(S)$.

Definition 19.2.1.2 (Cofibrant/fibrant objects). An object X in a model category \mathbf{C} is cofibrant (fibrant) if the unique map from initial object $\emptyset \rightarrow X$ (to terminal object $X \rightarrow \text{pt.}$) is a cofibration (fibration).