

# MATH 635 - Connections

A Ressegetes Story

May 28, 2024

## Contents

<b>1</b>	<b>Covariant Derivative</b>	<b>2</b>
<b>2</b>	<b>Connection in a Moving Frame</b>	<b>2</b>
<b>3</b>	<b>Covariant Differentiation along a Curve</b>	<b>2</b>
<b>4</b>	<b>Levi-Civita Connection</b>	<b>2</b>
<b>5</b>	<b>Covariant Differentiation of Tensors</b>	<b>2</b>

Recall in Euclidean space, we have the notion of directional derivative: given a  $C^1$ -continuous function  $f$  and a vector  $h$ , the directional derivative at  $v$  in  $h$  is given by

$$\left. \frac{d}{dt} f(v + th) \right|_{t=0} = \lim_{t \rightarrow 0} \frac{1}{t} (f(v + th) - f(v)) \quad (1)$$

or, given a vector field  $Y$ , a tangent vector  $v$ , and viewing tangent vectors as derivations we have in coordinates

$$\bar{\nabla}_v Y = v(Y^1) \left. \frac{d}{dx^1} \right|_p + \cdots + v(Y^n) \left. \frac{d}{dx^n} \right|_p$$

The goal, then, is to extend this to arbitrary manifolds, without resorting to its ambient Euclidean space. This cannot directly extend to general Riemannian manifolds, as at different points their tangent spaces are in fact different spaces, where, for example, the vector field version of Eq. (1) involves subtracting vectors in different tangent spaces, which is not properly defined. This chapter introduces tools for handling this problem.

## 1 Covariant Derivative

## 2 Connection in a Moving Frame

## 3 Covariant Differentiation along a Curve

## 4 Levi-Civita Connection

## 5 Covariant Differentiation of Tensors