

11.1 Hair coat types in mammals

Chapter 11

The hair of mammals can be characterized partly by the length and partly by the physical structure, for instance straight or curly. The ratio between the protective guard hair and the underfur also plays a role for its physical appearance. Finally the colour of the hair can give camouflage or in cases where individuals with strong colour dominates the flock or individuals outside the flock competing for dominance. Especially for water dwelling animals or animals hunting for fish the fur is of great significance for the regulation of heat. The fur of for example seal, otter and mink also has to be very resistant to last through the winter. The hare and foxes also have a warm fur, but the coat of these animals do not need to be nearly as strong as the before mentioned. Most mammals grow their winter fur in October-November and shed it again for the summer in April-May.

Some of the physical characteristics of the hair are under simple Mendelian inheritance, for instance the characteristics below:

Hair types and genotypes

Dominant		Recessive	
Wire haired	W-	Straight haired	ww
Short haired	L-	Long haired	ll
Guard haired	M-	Mole	mm
Haired	Hr-	Hairless	hr/hr

Figure 11.1 shows examples of the differently haired types.

Figure 11.1.

The wire-haired Airedale Terrier



Long haired type in Dachshund.



The fur from a mole mink and a normal Standard mink from year 1965.



Hairless mouse.



[Next page](#)