

"Targeted metabolomic profiling of the plasma in G2019S LRRK2 knockin mice under a specialized diet".

Dataset Description:

G2019S LRRK2 knockin mice and their wild-type littermates were maintained on a normal chow diet (23.7% kcal from fat) for the first 11 months of life. At that time, animals were randomly assigned to either a high-fat diet group (60% kcal from fat) or a control diet group (10% kcal from fat). Mice remained on their respective diets for an additional five months. At 16 months of age, mice were anesthetized with an isoflurane vaporizer, and whole blood was collected from cardiac puncture. 50 μ L plasma from each mouse was split and processed using the Bligh-Dyer method. Targeted metabolomics peak picking and integration were conducted in Skyline (v25.1) using accurate mass MS1, MS2 fragmentation pattern matching, and retention time derived from analytical standards run through each chromatography method. Raw data files for all samples of a given experiment were imported and metabolite peaks were auto-integrated based standard verified m/z, precursor adducts, and retention times for all metabolites of interest. Each experimental group contained biological replicates (n = 6 per group).

Authors:

- Ma, Yue; [ORCID:0000-0002-1031-5175](https://orcid.org/0000-0002-1031-5175); Van Andel Institute
- Moore, Darren; [ORCID:0000-0003-2359-5679](https://orcid.org/0000-0003-2359-5679); Van Andel Institute

ASAP Team: Lee

Dataset Name: lee-mouse-ms-mb-plasma-g2019s-hf-diet, v0.1

Principal Investigator: Darren J. Moore Darren.Moore@vai.org

Dataset Submitter: Yue Ma yue.ma@vai.org

Publication DOI: NA

Grant IDs: ['ASAP-000592']

ASAP Lab: MOORE-LAB

ASAP Project: Are there bidirectional pathological interactions between senescence and LRRK2 models of PD

Project Description: Mutations in LRRK2 gene cause late-onset, autosomal dominant forms of familial PD that are clinically indistinguishable from sporadic PD. In addition, LRRK2 gene polymorphisms are associated with sporadic PD risk. The late disease-onset with these PD-linked genes suggest that interactions with age-related pathways are critical for manifesting or unmasking their pathogenic actions. Studies in experimental models further suggest that PD-linked LRRK2 mutations can induce abnormal mitochondrial function, lysosomal activity, autophagy, proteostasis and inflammatory processes, pathways that are dysregulated by cellular senescence. Our project focuses on determining whether: (1) PD-linked mutations in LRRK2 induce cellular senescence in the rodent brain, (2) increased senescence can exacerbate or unmask PD-like neurodegenerative phenotypes in LRRK2 mouse models, and (3) if removing senescent cells will provide neuroprotection in LRRK2 rodent models of PD.

Submission Date: 2026-04-30

This dataset is made available to researchers via the ASAP CRN Cloud: cloud.parkinsonsroadmap.org. Instructions for how to request access can be found in the [User Manual](#).

This research was funded by the Aligning Science Across Parkinson's Collaborative Research Network (ASAP CRN), through the Michael J. Fox Foundation for Parkinson's Research (MJFF).

This Zenodo deposit was created by the ASAP CRN Cloud staff on behalf of the dataset authors. It provides a citable reference for a CRN Cloud Dataset