


Applied Statistical Analysis I

Statistical inference review

Elena Karagianni, PhD Candidate

karagiae@tcd.ie

 September 24, 2025

Department of Political Science, Trinity College Dublin

Today's class

- Lecture Recap
- Exercises

Key concepts and terms

- **Unit of analysis:** The observation described by a set of data. For example, voters, parties, bills, elections, voting decisions etc.

Key concepts and terms

- **Unit of analysis:** The observation described by a set of data. For example, voters, parties, bills, elections, voting decisions etc.
- **Variable:** A characteristic or attribute of the unit of analysis that can vary across observations.
 - “a characteristic that can vary in value among subjects in a sample or population” (Agresti and Finlay 2009, 11)

Key concepts and terms

- **Unit of analysis:** The observation described by a set of data. For example, voters, parties, bills, elections, voting decisions etc.
- **Variable:** A characteristic or attribute of the unit of analysis that can vary across observations.
 - “a characteristic that can vary in value among subjects in a sample or population” (Agresti and Finlay 2009, 11)
- **Dependent variable (DV):** Also called the *outcome* or *response variable*;; Denoted as Y ; the phenomenon we aim to explain.

Key concepts and terms


- **Unit of analysis:** The observation described by a set of data. For example, voters, parties, bills, elections, voting decisions etc.
- **Variable:** A characteristic or attribute of the unit of analysis that can vary across observations.
 - “a characteristic that can vary in value among subjects in a sample or population” (Agresti and Finlay 2009, 11)
- **Dependent variable (DV):** Also called the *outcome* or *response variable*;; Denoted as Y ; the phenomenon we aim to explain.
- **Independent variable (IV):** Also called the *input*, *predictor*, or *covariate*; Denoted as X ; used to explain variation in the DV.

Key concepts and terms

- **Unit of analysis:** The observation described by a set of data. For example, voters, parties, bills, elections, voting decisions etc.
- **Variable:** A characteristic or attribute of the unit of analysis that can vary across observations.
 - “a characteristic that can vary in value among subjects in a sample or population” (Agresti and Finlay 2009, 11)
- **Dependent variable (DV):** Also called the *outcome* or *response variable*;; Denoted as Y; the phenomenon we aim to explain.
- **Independent variable (IV):** Also called the *input*, *predictor*, or *covariate*; Denoted as X; used to explain variation in the DV.
- What is variation? (Example: Age \rightarrow Income)
- *Necessary terms for regression analysis.*

Refers to the way variables are quantified. (e.g., economic wealth measured as GDP).

Measurement







Figure 1: Kellstedt and Whitten 2018, Chap. 5

- What is the relationship between population and sample?

Population and Sample

- **Population:** the total set of subjects of interest in a study
- **Sample:** the subset of the population on which the study collects data
- **Parameter:** numerical summary of the population
- **Statistic:** a numerical summary of the sample data



- **Descriptive statistics:** “summarize the information in a collection of data”

Agresti and Finlay 2009, 4.

- **Inferential statistics:** “provide predictions about a population, based on data from a sample of that population”


Agresti and Finlay 2009, 4.

Descriptive Statistics


Measures of Central Tendency and Variability

- **Central tendency**: mean, median, mode
- **Variability**: variance, standard deviation, range, IQR
- **Visualization**: boxplots


Descriptive Statistics: Central Tendency



mode



median



mean


- **Mode**: Most frequently occurring value of X . Some distributions can have more than one mode.
- **Median**: Value of X that falls in the middle position when the observations are ordered from smallest to largest.
 - Median = 50th percentile = 2nd quartile
- **Mean**: Most common measure of central tendency.

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n}$$

Descriptive Statistics: Comparing Measures of Central Tendency

- In a perfectly symmetric distribution, e.g., normal distribution: **mode = median = mean**
- Not true when the distribution is non-symmetric:
 - right-skewed distribution (positive skew): $\text{median} < \text{mean}$
 - left-skewed distribution (negative skew): $\text{median} > \text{mean}$
- Mean is sensitive to outliers, while the median is more robust.

Descriptive Statistics: Comparing Measures of Central Tendency



Descriptive Statistics: Variability

- **Sample Variance**: Average of the square deviations from the mean:

$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}$$

Why do we average by dividing by $n-1$? The sum of the deviations is always zero. Thus, the last deviation can be found once we know the other $n-1$. So we are not averaging n unrelated numbers. Only $n - 1$ squared deviations vary freely, these are called *degrees of freedom* of the variance.

- **(Sample) Standard Deviation**: Square-root of (sample) variance:

$$s = \sqrt{s^2} = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}}$$

- **Range**: Difference between largest and smallest measurement:

$$Range = x_{max} - x_{min}$$

- **Interquartile Range (IQR)**: Difference between upper and lower quartiles (range of the middle 50% of the distribution):

$$IQR = x_{Q3} - x_{Q1} \quad (1)$$

Probability

- What is a probability?
- What is a distribution?
- What is a probability distribution?

Probability: Basic terminology

- An **experiment** is a repeatable procedure for making an observation.
- An **outcome** is a possible results of such an experiment.
- The **sample space** (Ω) of an experiment is the set of all possible outcomes.
- An **event** is a subset of the sample space, i.e., any set of outcomes.
- The **probability** of an event is its long-run relative frequency.
 - If $\Pr(A) = 0.5$, i.e., probability of event A is 0.5, then event A will occur approximately half of the time when the experiment is repeated infinitely often.
 - If the experiment is repeated many (finite) times, then the approximation as relative frequency (proportion) is expected to improve as the number of repetitions increases.

$$P(A) = \frac{\text{Number of elements in A}}{\text{Number of all elements}}$$

Probability: Distribution




Figure 2: Example: Age of people in the room

It can take different shapes and, therefore, names: normal, binomial, t-distribution etc.

Probability: Probability Distribution

- Distributions of random variables are **probability distributions** if for all possible outcomes, it tells us the probabilities for these outcomes to occur.

Probability: Probability Distribution


- Distributions of random variables are **probability distributions** if for all possible outcomes, it tells us the probabilities for these outcomes to occur.
- Probability distributions are analogous to **frequency distributions**, except that they are based on **probability theory** rather than observations in sample data.

Probability: Probability Distribution


- Distributions of random variables are **probability distributions** if for all possible outcomes, it tells us the probabilities for these outcomes to occur.
- Probability distributions are analogous to **frequency distributions**, except that they are based on **probability theory** rather than observations in sample data.
- Definition by Agresti and Finlay (2009, 75): “lists the possible outcomes and their probabilities.”

Sampling Distribution

Recall the basic idea for empirical research:



Sampling Distribution



Sampling Distribution

- **Sampling Distribution:** “A sampling distribution of a statistic is the probability distribution that specifies probabilities for the possible values the statistic can take” (Agresti and Finlay, 2009, 87).
- For example: we have a population distribution with mean μ and variance σ^2 and we are interested in its mean.
- Repeatedly taking samples from that population and calculating the mean for each sample yields the **sampling distribution of the mean**.

Sampling Distribution

Why is this important?

- The corresponding probability theory “helps us predict how close a statistic falls to the parameter it estimates” Agresti and Finlay 2009, 87 → how close is \bar{x} to μ ?
- Usually only one sample/estimate → **Point estimate**: “is a single number that is the best guess for the parameter value” Agresti and Finlay 2009, 107
- “If we repeatedly took samples, then in the long run, the mean of the sample means would equal the population mean μ ”.
- “The standard error describes how much \bar{x} varies from sample to sample” → the SE is estimated based on the standard deviation:

$$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$$

Central Limit Theorem

What is the Central Limit Theorem?


Central Limit Theorem

What is the Central Limit Theorem? → The sampling distribution of the statistic approaches a normal distribution with mean μ and variance σ^2/n as n increases.

- This holds *regardless* of the shape of the original population distribution.
- Basis for application of statistics to many 'natural' phenomena (which are the sum of many unobserved random events).
- How? Take a sample, calculate its mean. Do the same thing again and again.
The distribution of sample means will be normal even if the population distribution was not.
- If you repeatedly draw random samples from the same population, calculate the means and plot them, you get a histogram that approaches a bell-shaped curve.

Normal Distribution

- Continuous distribution that describes data clustered around the mean.
- Uniquely determined by its mean/median/mode μ and variance σ^2 .
- Important for the **Central Limit Theorem**.



Confidence Intervals

What are confidence intervals?

Confidence Intervals

What are confidence intervals?

“an interval of numbers around the point estimate that we believe contains the parameter value” → point estimate \pm margin of error (Agresti and Finlay 2009, 110)

Confidence Intervals

What are confidence intervals?

“an interval of numbers around the point estimate that we believe contains the parameter value” → point estimate \pm margin of error (Agresti and Finlay 2009, 110)

Let's explain this a bit more!

Confidence Intervals

- Our estimate of a population parameter varies across repeated samples, thus generating a *sampling distribution*.
- Instead of a point estimate, we should better get an **interval estimate** - a **range within the true parameter lies with some level of certainty**.
- We can construct **confidence intervals** using the standard error or the variance of our estimates.
- We call a CI a q% confidence interval if it is constructed that it **contains the true parameter at least q% of the time if we repeat the experiment a large number of times**.
- Check out this visualization: <https://rpsychologist.com/d3/ci/>
- Attention! This does not mean that there is a q% probability for the population parameter to lie inside the interval!