

Problem Set 1

Applied Stats/Quant Methods 1

Due: October 8, 2026

Instructions

- Please show your work! You may lose points by simply writing in the answer. If the problem requires you to execute commands in R, please include the code you used to get your answers. Please also include the .R file that contains your code. If you are not sure if work needs to be shown for a particular problem, please ask.
- Your homework should be submitted electronically on GitHub.
- This problem set is due before 23:59 on Thursday October 8, 2026. No late assignments will be accepted.

Question 1: Education

A school counselor was curious about the average of IQ of the students in her school and took a random sample of 25 students' IQ scores. The following is the data set:

```
1 y <- c(105, 69, 86, 100, 82, 111, 104, 110, 87, 108, 87, 90, 94, 113, 112, 98,  
      80, 97, 95, 111, 114, 89, 95, 126, 98)
```

1. Find a 90% confidence interval for the average student IQ in the school. (As a hint, you first need the mean and SD to create a CI.)

The data is ratio (continuous), the sample size is 25 thus a small sample size with an unknown standard deviation. For this reason a t -distribution is appropriate. The sampling method is randomization, and the school counselor is interested in estimating the population mean which could be either higher or lower than the sample mean. For a 90% confidence interval, the remaining 10% is divided equally between the two tails, with 5% in each tail.

```

1 sample_mean <- mean(y) #the mean is 98.44
2 sample_sd <- sd(y) #the standard deviation is 13.09287
3 sample_size <- length(y) #the length is 25
4
5 conf_level <- 0.90
6 alpha <- 1 - conf_level #confidence interval: 1 - 0.90 = 0.10
7 df <- sample_size - 1 #the degrees of freedom is 24
8
9 t_critical <- qt(1 - alpha / 2, df) # The sample size is 25 and the
   population standard deviation is unknown, so a t-test is appropriate.
10 margin_of_error <- t_critical * sample_sd / sqrt(sample_size) # The MOE
   is 4.480072
11
12 confidence_interval <- c(
13   lower = sample_mean - margin_of_error,
14   upper = sample_mean + margin_of_error
15 )
16
17 confidence_interval
18 t.test(y, conf.level = 0.90) # Check using R's built-in function

```

lower	upper
93.96	102.92

The 90% confidence interval is 93.96 to 102.92, thus we are 90% confident that the population mean IQ of students at the school is between the lower and upper limits of the confidence interval.

- Next, the school counselor was curious whether the average student IQ in her school is higher than the average IQ score (100) among all the schools in the country. Using the same sample, conduct the appropriate hypothesis test with $\alpha = 0.05$.

The assumptions above are still applicable except this is a right-tailed test. The Null Hypothesis is that average IQ in the school is less than or equal to the national average. The Alternative Hypothesis is that the average IQ in the school is higher than the national average.

$$H_0 : \mu \leq 100$$

$$H_1 : \mu > 100$$

```

1 iq_test <- t.test(y, mu = 100, alternative = "greater", conf.level =
   0.95)
2 iq_test # IQ needs to be greater, thus alternative needs to be greater.
3 ifelse(iq_test$p.value < 0.05,
4   "We reject H0",
5   "We fail to reject H0")

```

If the p-value ≤ 0.05 , reject H_0 ; otherwise, fail to reject H_0 . The p-value = 0.7215 thus at the 5% significance level, there is insufficient evidence to conclude that the average student IQ is greater than 100. We fail to reject the H_0 .

Question 2: Political Economy

Researchers are curious about what affects the amount of money communities spend on addressing homelessness. The following variables constitute our data set about social welfare expenditures in the USA.

State	50 states in US
Y	per capita expenditure on shelters/housing assistance in state
X1	per capita personal income in state
X2	Number of residents per 100,000 that are "financially insecure" in state
X3	Number of people per thousand residing in urban areas in state
Region	1=Northeast, 2= North Central, 3= South, 4=West

Explore the `expenditure` data set and import data into R.

- Please plot the relationships among Y , $X1$, $X2$, and $X3$? What are the correlations among them (you just need to describe the graph and the relationships among them)?

```
1 pairs(
2   expenditure[, c("Y", "X1", "X2", "X3")]
3 )
4
5 expenditure_cor <- cor(expenditure[, c("Y", "X1", "X2", "X3")])
6 expenditure_cor
7
8 expenditure_cor_no_diag <- expenditure_cor
9 diag(expenditure_cor_no_diag) <- NA
10 expenditure_cor_no_diag
11
12 expenditure_relationship <- matrix(
13   ifelse(
14     expenditure_cor_no_diag >= 0.7, "Strong Positive",
15     ifelse(
16       expenditure_cor_no_diag >= 0.3, "Moderate Positive",
17       ifelse(
18         expenditure_cor_no_diag > 0, "Weak Positive",
19         ifelse(
20           expenditure_cor_no_diag <= -0.7, "Strong Negative",
21           ifelse(
22             expenditure_cor_no_diag <= -0.3, "Moderate Negative",
23             ifelse(
24               expenditure_cor_no_diag < 0, "Weak Negative",
25               "No Relationship"
26             )
27           )
28         )
29       )
30     )
31   )
```

```

28     )
29   )
30 )
31 ),
32 nrow = nrow(expenditure_cor_no_diag),
33 dimnames = dimnames(expenditure_cor_no_diag)
34 )
35 # (Statistics Resources: Correlation, 2026)
36 expenditure_relationship

```

The scatterplot (Figure 1) matrix shows upward-sloping patterns between all pairs of variables, consistent with their positive correlations. The correlation shows X1 has the strongest relationship with Y, followed by X3 and X2 (Table 1 and 2).

- Please plot the relationship between Y and *Region*? On average, which region has the highest per capita expenditure on housing assistance?

```

1   levels = c(1, 2, 3, 4),
2   labels = c("Northeast", "North Central", "South", "West")
3 )
4
5 boxplot(
6   Y ~ RegionName,
7   data = expenditure,
8   xlab = "Region",
9   ylab = "Per Capita Expenditure on Housing Assistance",
10  main = "Per Capita Expenditure by Region"
11 )
12
13 regional_means <- aggregate(Y ~ RegionName, data = expenditure, mean)
14 regional_means
15 regional_means[which.max(regional_means$Y), ]

```

Figure 2 shows that the second boxplot (North Central) has the most outliers and the fourth boxplot (West) has the longest upper and lower whisker. The West (Region 4) has the highest average expenditure, with a average mean of 88.31 per capita. The North Central region (Region 2) has the second-highest average at 83.92, followed by the Northeast (Region 1) at 79.44. The South (Region 3) has the lowest average expenditure at 69.19 per capita (Table 3).

- Please plot the relationship between Y and $X1$? Describe this graph and the relationship. Reproduce the above graph including one more variable *Region* and display different regions with different types of symbols and colors.

```

1   ylab = "Per Capita Expenditure on Housing Assistance (Y)",
2   main = "Relationship Between Y and X1",
3   pch = 19,
4   col = "steelblue")

```

```

5
6 abline(lm(Y ~ X1, data = expenditure),
7         col = "red",
8         lwd = 2)
9
10 model <- lm(Y ~ X1, data = expenditure)
11 summary(model)$r.squared
12 coef(model)
13
14 cor_y_x1 <- cor(expenditure$Y, expenditure$X1)
15 round(cor_y_x1, 3)

```

The scatterplot shows a moderate positive relationship between X1 and Y: as per-capita personal income (X1) increases, per-capita expenditure on housing assistance (Y) generally increases. The correlation between the two variables is 0.532, indicating a moderate positive linear association (Table 2).

The abline's formula is $Y = 32.546 + 0.02458X_1$. Approximately 28.27% of the variation in Y is explained by X1 using this linear regression model.

```

1 plot(expenditure$X1, expenditure$Y,
2       xlab = "Per Capita Personal Income (X1)",
3       ylab = "Per Capita Expenditure on Housing Assistance (Y)",
4       main = "Relationship Between Y and X1 by Region",
5       pch = c(16, 17, 15, 18)[expenditure$Region],
6       col = c("red", "blue", "green", "purple")[expenditure$Region])
7
8 legend("topright",
9       inset = c(0.725, 0),
10      legend = c("Northeast", "North Central", "South", "West"),
11      pch = c(16, 17, 15, 18),
12      col = c("red", "blue", "green", "purple"),
13      title = "Region",
14      cex = 0.5,
15      xpd = TRUE)
16
17 par(xpd = NA)

```

There is a positive relationship, as per-capita personal income increases, housing-assistance expenditure generally tends to increase. This relationship is not very strong nor perfectly linear. Therefore income alone does not explain differences in housing-assistance expenditure and there are big differences (Figure 4).

Bibliography

Statistics Resources: Correlation. (2026, September 29). Retrieved from National University: <https://resources.nu.edu/statsresources/correlation> ““

Figures and Tables




Figure 1: Correlation Matrix of Y, X1, X2, and X3

Table 1: Correlation Matrix of the Dependent and Independent Variables

	Y	X_1	X_2	X_3
Y	NA	0.5317	0.4483	0.4637
X_1	0.5317	NA	0.2056	0.5953
X_2	0.4483	0.2056	NA	0.2210
X_3	0.4637	0.5953	0.2210	NA

Table 2: Classification of Correlation Strengths

	Y	X_1	X_2	X_3
Y	NA	Moderate Positive	Moderate Positive	Moderate Positive
X_1	Moderate Positive	NA	Weak Positive	Moderate Positive
X_2	Moderate Positive	Weak Positive	NA	Weak Positive
X_3	Moderate Positive	Moderate Positive	Weak Positive	NA

Table 3: Mean Per-Capita Expenditure on Shelters and Housing Assistance by Region

Region	Mean Y
Northeast	79.44444
North Central	83.91667
South	69.18750
West	88.30769




Figure 2: Average Per Capita Expenditure on Housing Assistance by Region




Figure 3: Scatter-plot of Y and X1




Figure 4: Scatter-plot of Y and X1 by Region