
Today
Line Search
Newton's Method
Conjugate Gradient

Announcements
Exam 2 Monday
Lab 3 due tomorrow

Fixed step size gradient descent is bad!



$$\vec{p}_k \leftarrow -\nabla f(\vec{x}_k)$$

$$\vec{x}_{k+1} = \vec{x}_k + \alpha_k \vec{p}_k$$

Solve $\underset{\alpha}{\text{minimize}}\ \phi(\alpha)$

with golden section search, bisection, etc.

In practice, exact line search is usually too expensive.


Better to find a "good enough" point ← sufficient decrease

Simple option: Armijo backtracking

Armijo condition: $\phi(\alpha) \le \phi(0) + \mu_1 \alpha \phi'(0)$

(μ_1: tuning param: usually 10^{-4})

($\phi'(0) = \nabla f(\vec{x}_k)^T \vec{p}_k$)




Armijo Backtracking Line Search

input : current point $\vec{x}_k$, descent direction $\vec{p}_k$, initial step α_{init} (1 for Newton), sufficient decrease factor μ_1 (e.g. 10^{-4}), backtracking factor ρ (e.g. 0.5)
output: step size α_k

$\phi(\alpha) \leftarrow f(\vec{x}_k + \alpha \vec{p}_k)$ // f along the line
$\phi'(0) \leftarrow \nabla f(\vec{x}_k)^T \vec{p}_k$ // slope at the start, < 0
$\alpha \leftarrow \alpha_{\text{init}}$
while $\phi(\alpha) > \phi(0) + \mu_1 \alpha \phi'(0)$ **do**
  $\alpha \leftarrow \rho \alpha$ // shrink the step
end
return $\alpha_k = \alpha$

Newton's Method



Taylor Series

$$f(\vec{x}_k + \vec{s}) = f(\vec{x}_k) + \nabla f(\vec{x}_k)^T \vec{s} + \frac{1}{2} \vec{s}^T H(\vec{x}_k) \vec{s} \ \ldots$$

$$\nabla_{\vec{s}} f(\vec{x}_k + \vec{s}) \approx \nabla f(\vec{x}_k) + H(\vec{x}_k) \vec{s}_k = 0$$

$$\vec{s} = -H(\vec{x}_k)^{-1} \nabla f(\vec{x}_k)$$

Newton update

$$\vec{x}_{k+1} \leftarrow \vec{x}_k - H(\vec{x}_k) \backslash \nabla f(\vec{x}_k)$$