

Definitions

Let V be a vector space over \mathbb{R} . The dual space of V is V^* , the set of linear functions $V \rightarrow \mathbb{R}$. Let $0 \leq k < \dim(V)$. The tensor product $(V^*)^{\otimes k}$ is the set of k -linear functions $V^k \rightarrow \mathbb{R}$. The exterior k -forms $\Lambda^k(V^*)$ are a subset of $(V^*)^{\otimes k}$, specifically the asymmetric functions.

Let M be a smooth manifold. Then $T_p M$ is the tangent plane of M at the point $p \in M$. Also, TM is informally used as a function $p \mapsto T_p M$. The dual space $(T_p M)^*$ is written $T_p^* M$. Similarly, $T^* M$ refers to $p \mapsto T_p^* M$. (Apparently TM is called the tangent bundle and $T^* M$ is called the cotangent bundle.) Note that $T_p \mathbb{R}^n$ is \mathbb{R}^n itself for any $p \in \mathbb{R}^n$.

Many of these linear algebraic objects can be parameterized by a point on a manifold. For example, vector spaces are generalized by smooth vector fields. For a smooth manifold M , define $\mathfrak{X}(M)$ as the set of smooth functions that take each point $p \in M$ to a vector in $T_p M$. Restricting to a single point p , we get a vector space $(T_p M)$.

Differential forms generalize exterior forms in the same way. For $0 \leq k < \dim(V)$, $\Omega^k(M)$ is the set of "smooth" functions that take each point $p \in M$ to a k -form in $\Lambda^k(T_p^* M)$. Restricting to a single point p , all we get a space of k -forms $(\Lambda^k(T_p^* M))$. The differential forms $\Omega^k(M)$ were defined in class as $\{\sum_I a_I dx_I \mid a_I \text{ smooth}\}$. This is essentially equivalent to the definition here, except "smooth" is in quotes since we never defined what it means for a function like this to be smooth, but it is the same idea. Note that $\Omega^0(M)$ is just $C^\infty(M, \mathbb{R})$, the set of all smooth functions $M \rightarrow \mathbb{R}$.

d

The symbol d has many meanings. Suppose $f : M \rightarrow N$ is a smooth function between smooth manifolds M and N . Then for $p \in M$, $df|_p : T_p M \rightarrow T_{f(p)} N$ is linear. Explicitly, with $v \in T_p M$, $df|_p(v) = \sum \frac{\partial f}{\partial x_i} v_i$.

$$d : \Omega^k \rightarrow \Omega^{k+1}$$