

ST5225: Statistical Analysis of Networks

Lecture 6: Matching Markets

WANG Wanjie
staww@nus.edu.sg

Department of Statistics and Applied Probability
National University of Singapore (NUS)

Saturday 10th March, 2018

- Review
- Graph Partition
- Matching Markets

- Graph Partition
 - Cluster by edge betweenness
 - Hierarchical clustering: Euclidean distance, other dissimilarities; different linkages
 - Modularity
 - Dendrogram
- Bipartite graph
 - Balanced bipartite graph
 - Perfect matching: connectivity and distinction
 - Optimal assignment: maximize the overall valuation

For today:

- Matching Markets: Optimal assignments and Market-clearing prices

The model we start with is called *bipartite matching problem*.

Bipartite graph

For a graph $G = (V, E)$, if the set of vertices V can be divided into two disjoint and independent sets V_1 and V_2 such that every edge connects a vertex in V_1 to one in V_2 , then we call G as a bipartite graph.

Remark.

- Bipartite graph usually has two types of nodes (dorms and students).
- V_1 and V_2 may have different number of nodes. If they have the same number of nodes, we call it as *balanced*
- Bipartite matching problem is concerning whether there is a *perfect matching* on balanced bipartite graph.

Perfect Matching

When there are an equal number of nodes on each side of a bipartite graph, a *perfect matching* is an assignment of nodes on the left to nodes on the right, in such a way that



- (Connection) each node is connected by an edge to the node it is assigned to, and
- (distinction) no two nodes on the left are assigned to the same node on the right.

Remark.

- For a perfect matching, each node in V_1 is connected to one and only one “friend” in V_2 , and vice versa.

Dorm assignment problem

- 5 dorms available for 5 students; each room for 1 single student
- Each student has a list of acceptable rooms; shown in graph



Question: Are there always be such an assignment?

Constricted sets

A set S is constricted if $|S| > |N(S)|$.

Matching Theorem

If a balance bipartite graph has no perfect matching, then it must contain a constricted set.


- Equivalence between *perfect matching* and *no constricted set*

Perfect matching \iff No constricted sets


Room1	Xin	Valuations 12, 2, 4
Room2	Yoram	8, 7, 6
Room3	Zoe	7, 5, 2

- The three numbers show the evaluation for the three rooms
- We call the assignment that can maximize the overall evaluation score as an optimal assignment


- A university may not have the same charge on the rooms, yet the market won't
- On a market, there are a set of sellers and a set of buyers. Each buyer has valuation on the product of the sellers, and each seller ask for a price.




- 3 houses available for 3 buyers
- Each buyer has his/her valuations for each house. Each seller has a price for his/her house
- Questions
 - Can we still find an “optimal” assignment here?
 - If we can change the price, can we find an even better assignment?








- 3 houses available for 3 buyers
- Each buyer has his/her valuations for each house. Each seller has a price for his/her house
- Questions
 - Can we still find an “optimal” assignment here?
 - If we can change the price, can we find an even better assignment?















- 3 houses available for 3 buyers
- Each buyer has his/her valuations for each house. Each seller has a price for his/her house
- Questions
 - Can we still find an “optimal” assignment here?
 - If we can change the price, can we find an even better assignment?




- Payoffs: For a buyer and a house, the buyer's payoff is valuation – price of this house
- Therefore, we have a graph with *payoffs* as the previous *valuations*
- Given the prices, we can find an *optimal assignment* with respect to the payoffs
- The optimal assignment could maximize the payoffs

Prices	Sellers	Buyers	Valuations	Sellers	Buyers	Payoffs
5			12, 4, 2			7, 2, 2
2			8, 7, 6			3, 5, 6
0			7, 5, 2			2, 3, 2


- For the result, note that each buyer is connected to the seller with largest *payoff*
- Preferred-seller graph: Connect each buyer with the sellers that maximizes the payoff of this buyer, and the corresponding graph is called preferred-seller graph, since the seller is the preferred seller of the connected buyers.
- If there is a perfect matching in the preferred-seller graph, then this perfect matching is the optimal assignment.

Prices	Sellers	Buyers	Valuations	Sellers	Buyers	Payoffs
5			12, 4, 2			7, 2, 2
2			8, 7, 6			3, 5, 6
0			7, 5, 2			2, 3, 2

- Note: every buyer gets the house that maximizes the payoffs
- We call such a set of prices *market-clearing*, since the prices setting cause each house to get bought by a different buyer—the house market is clear!
- It is possible there is no perfect matching



(c) Prices that Don't Clear the Market



(d) Market-Clearing Prices (Tie-Breaking Required)

- *Existence: for any set of buyer valuations, there exists a set of market-clearing prices*
 - For any set of buyers, the sellers can set the prices carefully, so that each buyer gets distinct house

- *Optimality: For any set of market-clearing prices, the optimal assignment has the maximum total valuation of any assignment of sellers to buyers*
 - For an assignment M

Total payoff of $M = \text{Total valuation of } M - \text{Sum of all prices}$
 - Note that sum of all prices don't change. The optimal assignment maximizes the total payoff, so that maximizes the total valuation.

- *Existence: for any set of buyer valuations, there exists a set of market-clearing prices*
 - For any set of buyers, the sellers can set the prices carefully, so that each buyer gets distinct house

- *Optimality: For any set of market-clearing prices, the optimal assignment has the maximum total valuation of any assignment of sellers to buyers*
 - For an assignment M

Total payoff of $M = \text{Total valuation of } M - \text{Sum of all prices}$
 - Note that sum of all prices don't change. The optimal assignment maximizes the total payoff, so that maximizes the total valuation.

How to construct a set of market-clearing prices?

- Idea: if one house is in high demand (more than 1 buyers have maximum payoff on this house), then the price increases by 1
- Note: we always set the house with smallest price to be with price 0. It does not affect the result, and it helps to scale the result
- In a real market, the “hot” house usually has a higher price, so that the final buyer is the buyer who likes it most

How to construct a set of market-clearing prices?

- Idea: if one house is in high demand (more than 1 buyers have maximum payoff on this house), then the price increases by 1
- Note: we always set the house with smallest price to be with price 0. It does not affect the result, and it helps to scale the result
- In a real market, the “hot” house usually has a higher price, so that the final buyer is the buyer who likes it most

How to construct a set of market-clearing prices?


- Idea: if one house is in high demand (more than 1 buyers have maximum payoff on this house), then the price increases by 1
- Note: we always set the house with smallest price to be with price 0. It does not affect the result, and it helps to scale the result
- In a real market, the “hot” house usually has a higher price, so that the final buyer is the buyer who likes it most

Say there are n sellers (items) and buyers



Start: each item has price 0; each buyer assigns a value to each item

- 1** Assume smallest value is 0; if not, scale the price so that the smallest is 0
- 2** Construct the preferred-seller graph and check if there is a perfect matching
- 3** If yes, done
- 4** If not, find a constricted set of buyers S
- 5** Each seller in $N(S)$ increases the price by 1
- 6** check if the smallest price is 0, if not, subtract the same amount of each price so that the smallest is 0
- 7** go back to Step 1.


Market-Clearing Prices: Example



(a) Start of first round



(b) Start of second round



Market-Clearing Prices: Example II

Prices	Sellers	Buyers	Valuations
0			5, 7, 1
0			2, 3, 1
0			5, 4, 4

- Bipartite graphs with two sets of nodes V_1 and V_2 ; each $v_i \in V_1$ has a price; each $v_j \in V_2$ has a valuation for every $v_i \in V_1$
 - Find payoffs of each $v_j \in V_2$ assigned to each $v_i \in V_1$
 - With the payoffs, there is *optimal assignment*
 - For each $v_j \in V_2$, there is a preferred seller in V_1 , which maximizes the payoff of v_j . We call this seller as *preferred seller*
 - If the preferred-seller graph has a *perfect matching*, then we call the price as market-clearing price
- Market-clearing prices must exist, and maximizes the total valuations
- Construction of market-clearing prices: begin with 0, and increase the prices of constricted set by 1.

- Power on networks
- An experiment and the results
- Nash bargaining solution
- Ultimatum game
- Stable outcomes
- Find natural stable outcomes


- In the buyer-seller network, the buyer has payoff as valuation minus price.
- The seller may want to increase the price, so that the payoff splits between the seller and the buyer
- If the seller increases the price only a little bit, then the buyer still transacts with this seller with a fewer payoff, and the seller gets some profit.
- If the seller increases the price too much, the buyer may give up, and no transaction is done.
- *Bargaining on network*
- Bargaining is related to *power*, the popularity of the seller/buyer in the network

We talk about the general cases, not restricted to buyer-seller networks.

- Intuitively, we may say these two nodes have the same power (related to this graph), since they are symmetric



- Node B in this graph has a larger power, since nodes A and C requires the connection with B to get in touch with D and E



- Power of distributing resources


Set up an experiment to see people's bargaining behavior on networks.

- 1** Fix a small network. Let each node represent each player. Players that are connected with an edge can exchange information/make offers.
- 2** Each edge carries a resource (e.g., \$100). This resource is split among adjacent nodes if there is a deal between nodes.
- 3** Each node can only be part of at most one deal (possible that no deal is done).
- 4** Players can freely negotiate for a fixed amount of time how the resource is split.
- 5** The experiment is repeated multiple times to get the final result

- For a 2-node network, end up with a fair split. Since A and B has the same power on the transaction, they agree to split the money equally.




- For a 3-node network, B has higher power, where it can exclude A or C in the transaction. The result shows that A and C are symmetric, while B receives the majority of the money in the transaction.



- In the textbook, more complicated experiments, say, 4-nodes, 5-nodes are also discussed.

To discuss the problem, we think of each edge. The resources outside this edge is summarized as “outside options”, as the following figure



- Assume the total resource to split on this edge is 1.
- Since A can get x from outside if not split with B. So the expectation of A's payoff is at least x
- Similarly, the expectation of B's payoff is at least y
- If $x + y > 1$, impossible to satisfy both A and B
- If $x + y \leq 1$, then there is surplus $s = 1 - x - y$
- Nash bargaining solution: both A and B will get half of the surplus, $(1 - x - y)/2$, which means the outcome is

$$x + \frac{1 - x - y}{2} = \frac{1 + x - y}{2} \text{ to A; } y + \frac{1 - x - y}{2} = \frac{1 - x + y}{2} \text{ to B.}$$

Interesting remarks:

- The resource is not only money, it can be friendship, collaboration, etc.
- When bargaining, people tries to communicate about the outside option, so that gets a reasonable split
- In real life, for the two endpoints, usually one has a “higher-status” and the other has a “lower-status”. For example, a sophomore and a graduate student with high grades.
- It has been found that, people with high-status tend to inflate the size of their outside option; people with lower-status tend to reduce the size of their outside option.
- What’s more, the reduced size of outside option will be further underestimated by the higher-status people.
- Do you have the same problem in your job search?

Setup: Two players A and B; no communication allowed; 1-time game

- 1** Person A is given \$100 and told to propose a division among A and B
- 2** If B accepts the \$100 are split accordingly; if B rejects, both A and B get nothing

If both A and B are rational,

- B should accept any positive offer, since that's better than nothing
- Since B will accept any positive offer, A, as the person who propose the division, can keep more money
- Solution: A offers \$99 to A, and \$1 to B
- The difference comes from the power of division

Setup: Two players A and B; no communication allowed; 1-time game

- 1** Person A is given \$100 and told to propose a division among A and B
- 2** If B accepts the \$100 are split accordingly; if B rejects, both A and B get nothing

If both A and B are rational,

- B should accept any positive offer, since that's better than nothing
- Since B will accept any positive offer, A, as the person who propose the division, can keep more money
- Solution: A offers \$99 to A, and \$1 to B
- The difference comes from the power of division

Setup: Two players A and B; no communication allowed; 1-time game

- 1 Person A is given \$100 and told to propose a division among A and B
- 2 If B accepts the \$100 are split accordingly; if B rejects, both A and B get nothing

Reality:

- In 1/3 of the experiments, A offers 1/2-1/2 split to B
- Moreover, when the offer is very unfair, B rejects to accept the offer – even though he will get nothing
- When the amount of money changes, the result is still the same
- Guess: It hurts B's feelings when offered very small proportion, which behaves as a negative payoff
- Solution: Give B more time to consider, then more people would accept the “unfair” offer


- We need some mathematically defined terms to describe the exchanges on graphs

Outcomes

An *outcome* of a network exchange on a given graph consists of:

- A matching on the set of nodes, specifying who exchanges with whom.
- A number associated with each node, called its *value*, indicating how much this node gets from its exchange. The sum of matched nodes should be 1.

Example.




- All discussions are based on the outcomes
- Is the outcome stable?

Stable outcome


Outcome where no player can make an offer to another player such that both are *better off*, is called a stable outcome.




(a) Not a stable outcome




(b) A stable outcome



(c) Not a stable outcome



(d) A stable outcome




(e) A stable outcome

- *Instability*: if there is an edge not part of the matching, such that the sum of the endpoint values is < 1 , then we call this outcome as *an unstable outcome*.
 - Say the endpoints of this edge is A and B , then obviously A and B would prefer to exchange, so that they are better off. The original outcome is not stable.
- If there is no such instabilities, then an outcome is stable.
 - If an outcome is unstable, then there always exists such an edge.
 - It helps us to figure out the stability of an outcome. Just check all the edges and the values of their endpoints, and we know whether it is stable or not

- *Instability*: if there is an edge not part of the matching, such that the sum of the endpoint values is < 1 , then we call this outcome as *an unstable outcome*.
 - Say the endpoints of this edge is A and B , then obviously A and B would prefer to exchange, so that they are better off. The original outcome is not stable.
- If there is no such instabilities, then an outcome is stable.
 - If an outcome is unstable, then there always exists such an edge.
 - It helps us to figure out the stability of an outcome. Just check all the edges and the values of their endpoints, and we know whether it is stable or not

Example (restrictions of outcomes):

For a triangle, there are no stable outcomes!




- If we have many stable outcomes, then we prefer some “natural” set of outcomes, for which we call *balanced*


Balanced outcomes

An outcome is called a balanced outcome if, for each edge in the matching, the split of money represents the Nash bargaining outcome for the two nodes involved, given the best outside option.

Example: a balanced outcome



- It is closer to what happens in reality



- Power on networks with experiments
- Nash bargaining solution for two nodes with outside options
- Ultimatum game and the results
- Stable outcomes and natural stable outcomes (Nash bargaining solution required here)