

1 — PHY 494: Makeup assignment 1 (15 points total)

Due Thursday, April 22, 2021, 11:59pm.

This is a **optional Makeup Assignment**. If you choose to hand it in by the deadline, it will be graded like a normal homework assignment. If its grade is better than your worst homework grade then it will replace that grade.

| |
|--|
| Submission is to your private GitHub repository . |
|--|

1.1 ODE with Scipy (15 points)

In this problem you should learn how to use existing functions in a library, namely in the `scipy` package (in particular, [scipy.integrate](#)). Part of the problem is to find and read the documentation for the function and figure out how to use it.

Use the Scipy function `scipy.integrate.solve_ivp()` to solve the following ordinary differential equation

$$-\frac{1}{2} \frac{d^2 \psi_n(x)}{dx^2} + \left[\frac{1}{2} x^2 - E_n \right] \psi_n(x) = 0 \quad (1)$$

$$\psi_n(0) = 1; \quad \frac{d\psi_n(0)}{dx} = 0 \quad (2)$$

$$E_n = n + \frac{1}{2}, \quad n = 0, 2, 4, 6, \dots \quad (3)$$

for the real function $\psi_n(x)$ ¹ and the three values $n = 0$, $n = 2$, and $n = 8$.

The code should contain the following functions:

1. **ode_qmhosc()** solves the ODE Eq. 1:

```
psi = ode_qmhosc(x, psi0, dpsidx0, n=n)
```

that takes as arguments

¹If you have done Quantum Mechanics 1 (PHY 314) then you should recognize it as the Schrödinger equation for the simple harmonic oscillator.

- all the values x at which the solution should be evaluated (the first value *must* be the one for which the initial conditions are given, i.e., $x = 0$ for this problem),
- as initial conditions the function value and first derivative at $x = 0$ (Eq. 2), and
- the value of n that determines E_n (Eq. 3).

It should return the function values $\psi_n(x)$ and the derivatives $\psi'_n(x)$ as a numpy array of shape `(2, len(x))` (which is what the “y” key in the return result of `solve_ivp()` gives you).

2. `f_qmhosc()` is needed to transform the ODE Eq. 1 into standard form so that it can be solved with `scipy.integrate.solve_ivp()`.

`f_qmhosc(t, y, E)`

should produce the *ODE standard form*² of the derivative vector \mathbf{f} when provided with

- the current values of \mathbf{y}
- the current value of t
- and the parameter E_n (see Eq. 1).

The starter code in `makeup01.py` has additional hints on how to create a function that is suitable for `scipy.integrate.solve_ivp()`.

- (a) Numerically solve Eq. 1 on a lattice with $\Delta x = 0.01$ for $0 \leq x < 6$ for
 - $n = 0$
 - $n = 2$

²Remember that the ODE standard form is

$$\frac{d\mathbf{y}}{dt} = \mathbf{f}$$

and in the case of Eq. 1, we would have $t = x$, $y_0 = \psi_n(x)$, and $y_1 = \psi'_n(x) \equiv \frac{d\psi_n}{dx}$.

- $n = 8$

Use the *rk45* solver (find out how to select it for the `solve_ivp()` function) and increase the default tolerance parameters for the solver to `rtol=1e-8, atol=1e-8`.

- Write code that implements all the listed requirements. **[12 points]**
 - Plot all your solutions in one figure and include the figure as a PNG named `qmhosc.png`. **[3 points]**
- (b) BONUS: Correct physical³ solutions for Eq. 1 are *symmetric* ($\psi(x) = \psi(-x)$) for even n and *antisymmetric* ($\psi(x) = -\psi(-x)$) for odd n . This implies that the *initial conditions* for $n = 1, 3, 5, \dots$ are different from the ones for even n (Eq. 2). Choose appropriate initial conditions⁴ and write a function

```
phi0, dphidx0 = initial_values_qmhosc(n)
```

that produces the correct initial values depending on the value of n .

Using your initial values, solve Eq. 1 for $n = 3$ and plot the solution (include a PNG named `qmhosc-odd.png`). **[bonus +4*]**

³Physical solutions for the wavefunction $\psi_n(x)$ have to be normalizable, i.e., they have to decay to zero for $x \rightarrow \pm\infty$. The numerical solutions only fulfill this requirement for small x and even for moderately large x they start to diverge. It would be better to use specialized algorithms that have the normalizability requirement built in.

⁴Choose $\psi(0) \geq 0$, $\frac{d\psi(0)}{dx} \geq 0$ and either use values of 0 or 1 because this is the convention employed in the tests.