

# LECTURE 8: PRECISION OF OLS AND HYPOTHESIS TESTING

ECON 480 - ECONOMETRICS - FALL 2018

---

Ryan Safner

September 26, 2018

The Precision of OLS

Hypothesis Testing About Regression

Digression:  $p$ -Values and the Philosophy of Science

Back to Our Hypothesis Test: The Test-Statistic


Reporting Regression Outputs with **stargazer**

## THE PRECISION OF OLS

---


$$\hat{\beta}_1 \sim N(E[\hat{\beta}_1], \sigma_{\hat{\beta}_1})$$

- We want to know:




$$\hat{\beta}_1 \sim N(E[\hat{\beta}_1], \sigma_{\hat{\beta}_1})$$

- We want to know:
  - $E[\hat{\beta}_1]$ ; what is the center of the distribution?




$$\hat{\beta}_1 \sim N(E[\hat{\beta}_1], \sigma_{\hat{\beta}_1})$$

- We want to know:
  - $E[\hat{\beta}_1]$ ; what is the center of the distribution?
  - $\sigma_{\hat{\beta}_1}$ ; how precise is our estimate?



## PRECISION: VARIANCE OR STANDARD ERROR


- How precise is our estimate  $\hat{\beta}_1$ ?



<sup>1</sup>The "standard **error**" is the analogue of standard *deviation* for a sample statistic's sampling distribution. Recall the sampling distribution is the distribution of a statistic, like  $\bar{X}$  or  $\hat{\beta}_1$  over many potential samples.

## PRECISION: VARIANCE OR STANDARD ERROR

- How precise is our estimate  $\hat{\beta}_1$ ?
- We can talk of the **variance** ( $\sigma_{\hat{\beta}_1}^2$ ) or the **standard error** ( $\sigma_{\hat{\beta}_1}$ ) of  $\hat{\beta}_1$ <sup>1</sup>



<sup>1</sup>The "standard **error**" is the analogue of standard *deviation* for a sample statistic's sampling distribution. Recall the sampling distribution is the distribution of a statistic, like  $\bar{X}$  or  $\hat{\beta}_1$  over many potential samples.



- The variance of  $\hat{\beta}_1$  is

$$\text{var}(\hat{\beta}_1) = \frac{(\text{SER})^2}{n \times \text{var}(X)}$$

where SER is the standard error of the regression (again):

$$\text{SER} = \frac{\text{SSE}}{n - 2}$$

- The **variance** of  $\hat{\beta}_1$  is

$$\text{var}(\hat{\beta}_1) = \frac{(\text{SER})^2}{n \times \text{var}(X)}$$

where SER is the standard error of the regression (again):

$$\text{SER} = \frac{\text{SSE}}{n - 2}$$

- The **standard error** of  $\hat{\beta}_1$  is simply the square root of the variance

$$\text{se}(\hat{\beta}_1) = \sqrt{\text{var}(\hat{\beta}_1)}$$

$$\text{var}(\hat{\beta}_1) = \frac{(\text{SER})^2}{n \times \text{var}(X)}$$

- Variance is affected by three things

$$\text{var}(\hat{\beta}_1) = \frac{(SER)^2}{n \times \text{var}(X)}$$

- Variance is affected by three things
  1. Goodness of fit of the model: *SER*

$$\text{var}(\hat{\beta}_1) = \frac{(\text{SER})^2}{n \times \text{var}(X)}$$

- Variance is affected by three things
  1. **Goodness of fit of the model:  $\text{SER}$** 
    - Larger  $\text{SER}$ , larger  $\text{var}(\hat{\beta}_1)$

$$\text{var}(\hat{\beta}_1) = \frac{(SER)^2}{n \times \text{var}(X)}$$

- Variance is affected by three things
  1. **Goodness of fit of the model:  $SER$** 
    - Larger  $SER$ , larger  $\text{var}(\hat{\beta}_1)$
  2. **Sample size,  $n$**

$$\text{var}(\hat{\beta}_1) = \frac{(SER)^2}{n \times \text{var}(X)}$$

- Variance is affected by three things
  1. **Goodness of fit of the model:  $SER$** 
    - Larger  $SER$ , larger  $\text{var}(\hat{\beta}_1)$
  2. **Sample size,  $n$** 
    - Larger  $n$ , lower  $\text{var}(\hat{\beta}_1)$

$$\text{var}(\hat{\beta}_1) = \frac{(SER)^2}{n \times \text{var}(X)}$$


- Variance is affected by three things
  1. **Goodness of fit of the model:  $SER$** 
    - Larger  $SER$ , larger  $\text{var}(\hat{\beta}_1)$
  2. **Sample size,  $n$** 
    - Larger  $n$ , lower  $\text{var}(\hat{\beta}_1)$
  3. **Variation in  $X$**



$$\text{var}(\hat{\beta}_1) = \frac{(\text{SER})^2}{n \times \text{var}(X)}$$


- Variance is affected by three things
  1. **Goodness of fit of the model:  $\text{SER}$** 
    - Larger  $\text{SER}$ , larger  $\text{var}(\hat{\beta}_1)$
  2. **Sample size,  $n$** 
    - Larger  $n$ , lower  $\text{var}(\hat{\beta}_1)$
  3. **Variation in  $X$** 
    - Larger  $\text{var}(X)$ , smaller  $\text{var}(\hat{\beta}_1)$

## THE RELATION BETWEEN VARIANCE OF $X$ AND VARIANCE OF $\hat{\beta}_1$



- Smaller  $\text{var}(X)$  (light dots only)  $\implies$  larger  $\text{var}(\hat{\beta}_1)$ : harder to determine precise slope!

## THE RELATION BETWEEN VARIANCE OF $X$ AND VARIANCE OF $\hat{\beta}_1$



- Smaller  $\text{var}(X)$  (light dots only)  $\implies$  larger  $\text{var}(\hat{\beta}_1)$ : harder to determine precise slope!
- Larger  $\text{var}(X)$  (all dots)  $\implies$  smaller  $\text{var}(\hat{\beta}_1)$ : easier to determine precise slope!

## HYPOTHESIS TESTING ABOUT REGRESSION

---

- We have used statistics (OLS) to **estimate** a relationship between  $X$  and  $Y$

- We have used statistics (OLS) to **estimate** a relationship between  $X$  and  $Y$ 
  - *Relationship is causal if  $X$  is exogenous*

- We have used statistics (OLS) to **estimate** a relationship between  $X$  and  $Y$ 
  - *Relationship is causal if  $X$  is exogenous*
- The “bread and butter” of inferential statistics is **testing a hypothesis** about (a) population(s) parameter(s)

- We have used statistics (OLS) to **estimate** a relationship between X and Y
  - *Relationship is causal if X is exogenous*
- The “bread and butter” of inferential statistics is **testing a hypothesis** about (a) population(s) parameter(s)
  - Does reducing class size actually improve test scores?



- We have used statistics (OLS) to **estimate** a relationship between  $X$  and  $Y$ 
  - *Relationship is causal if  $X$  is exogenous*
- The “bread and butter” of inferential statistics is **testing a hypothesis** about (a) population(s) parameter(s)
  - Does reducing class size actually improve test scores?
  - Is the gender wage gap between men and women really \$0.77?

- We have used statistics (OLS) to **estimate** a relationship between X and Y
  - *Relationship is causal if X is exogenous*
- The “bread and butter” of inferential statistics is **testing a hypothesis** about (a) population(s) parameter(s)
  - Does reducing class size actually improve test scores?
  - Is the gender wage gap between men and women really \$0.77?
  - What percent of American voters support legalizing marijuana?

- We have used statistics (OLS) to **estimate** a relationship between X and Y
  - *Relationship is causal if X is exogenous*
- The “bread and butter” of inferential statistics is **testing a hypothesis** about (a) population(s) parameter(s)
  - Does reducing class size actually improve test scores?
  - Is the gender wage gap between men and women really \$0.77?
  - What percent of American voters support legalizing marijuana?
- **All modern science is built upon statistical hypothesis testing, so understand it well!**

- Note, we can test a lot of hypotheses about a lot of population parameters, e.g.

- Note, we can test a lot of hypotheses about a lot of population parameters, e.g.
  - A population mean  $\mu$  (e.g. average height of adults)

- Note, we can test a lot of hypotheses about a lot of population parameters, e.g.
  - A population mean  $\mu$  (e.g. average height of adults)
  - A population proportion  $p$  (e.g. percent of voters who voted for Trump)

- Note, we can test a lot of hypotheses about a lot of population parameters, e.g.
  - A population mean  $\mu$  (e.g. average height of adults)
  - A population proportion  $p$  (e.g. percent of voters who voted for Trump)
  - A difference in population means  $\mu_A - \mu_B$  (e.g. difference in average wages of men vs. women)

- Note, we can test a lot of hypotheses about a lot of population parameters, e.g.
  - A population mean  $\mu$  (e.g. average height of adults)
  - A population proportion  $p$  (e.g. percent of voters who voted for Trump)
  - A difference in population means  $\mu_A - \mu_B$  (e.g. difference in average wages of men vs. women)
  - A difference in population proportions  $p_A - p_B$  (e.g. difference in percent of patients reporting symptoms of drug A vs B)



- Note, we can test a lot of hypotheses about a lot of population parameters, e.g.
  - A population mean  $\mu$  (e.g. average height of adults)
  - A population proportion  $p$  (e.g. percent of voters who voted for Trump)
  - A difference in population means  $\mu_A - \mu_B$  (e.g. difference in average wages of men vs. women)
  - A difference in population proportions  $p_A - p_B$  (e.g. difference in percent of patients reporting symptoms of drug A vs B)
  - See all the possibilities in glorious detail in the **handout** on Inferential Statistics

- Note, we can test a lot of hypotheses about a lot of population parameters, e.g.
  - A population mean  $\mu$  (e.g. average height of adults)
  - A population proportion  $p$  (e.g. percent of voters who voted for Trump)
  - A difference in population means  $\mu_A - \mu_B$  (e.g. difference in average wages of men vs. women)
  - A difference in population proportions  $p_A - p_B$  (e.g. difference in percent of patients reporting symptoms of drug A vs B)
  - See all the possibilities in glorious detail in the **handout** on Inferential Statistics
- We will focus on hypotheses about the population regression slope ( $\hat{\beta}_1$ ) between  $X$  and  $Y$

- Note, we can test a lot of hypotheses about a lot of population parameters, e.g.
  - A population mean  $\mu$  (e.g. average height of adults)
  - A population proportion  $p$  (e.g. percent of voters who voted for Trump)
  - A difference in population means  $\mu_A - \mu_B$  (e.g. difference in average wages of men vs. women)
  - A difference in population proportions  $p_A - p_B$  (e.g. difference in percent of patients reporting symptoms of drug A vs B)
  - See all the possibilities in glorious detail in the **handout** on Inferential Statistics
- We will focus on hypotheses about the population regression slope ( $\hat{\beta}_1$ ) between  $X$  and  $Y$ 
  - i.e. if/when we've done our model right, the **causal effect of  $X$  on  $Y$**

- All scientific inquiries begin with a null hypothesis ( $H_0$ ) that proposes a specific value of a population parameter

- All scientific inquiries begin with a **null hypothesis ( $H_0$ )** that proposes a specific value of a population parameter
  - Notation: add a subscript 0:  $\beta_{1,0}$  (or  $\mu_0$ ,  $p_0$ , etc)

- All scientific inquiries begin with a **null hypothesis ( $H_0$ )** that proposes a specific value of a population parameter
  - Notation: add a subscript 0:  $\beta_{1,0}$  (or  $\mu_0$ ,  $p_0$ , etc)
- We suggest an **alternative hypothesis ( $H_a$ )**, often the one we hope to verify

- All scientific inquiries begin with a **null hypothesis ( $H_0$ )** that proposes a specific value of a population parameter
  - Notation: add a subscript 0:  $\beta_{1,0}$  (or  $\mu_0, p_0$ , etc)
- We suggest an **alternative hypothesis ( $H_a$ )**, often the one we hope to verify
  - Note, can be multiple alternative hypotheses:  $H_1, H_2, \dots, H_n$

- All scientific inquiries begin with a **null hypothesis ( $H_0$ )** that proposes a specific value of a population parameter
  - Notation: add a subscript 0:  $\beta_{1,0}$  (or  $\mu_0, p_0$ , etc)
- We suggest an **alternative hypothesis ( $H_a$ )**, often the one we hope to verify
  - Note, can be multiple alternative hypotheses:  $H_1, H_2, \dots, H_n$
- Ask: “Does our data (sample) give us sufficient evidence to reject  $H_0$  in favor of  $H_a$ ?”



- All scientific inquiries begin with a **null hypothesis ( $H_0$ )** that proposes a specific value of a population parameter
  - Notation: add a subscript 0:  $\beta_{1,0}$  (or  $\mu_0, p_0$ , etc)
- We suggest an **alternative hypothesis ( $H_a$ )**, often the one we hope to verify
  - Note, can be multiple alternative hypotheses:  $H_1, H_2, \dots, H_n$
- Ask: **“Does our data (sample) give us sufficient evidence to reject  $H_0$  in favor of  $H_a$ ?”**
  - Note: the test is *always* about  $H_0$ ! See if we have sufficient evidence to reject the status quo

- Null hypothesis assigns a value (or a range) to a population parameter

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**
  - Note: always an equality!

## NULL AND ALTERNATIVE HYPOTHESES II

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**
  - Note: always an equality!
- Alternative hypothesis must mathematically *contradict* the null hypothesis

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**
  - Note: always an equality!
- Alternative hypothesis must mathematically *contradict* the null hypothesis
  - e.g.  $\beta_1 \neq 2$  or  $\beta_1 > 20$  or  $\beta_1 \neq 0$

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**
  - Note: always an equality!
- Alternative hypothesis must mathematically *contradict* the null hypothesis
  - e.g.  $\beta_1 \neq 2$  or  $\beta_1 > 20$  or  $\beta_1 \neq 0$
  - Note: always an inequality!



## NULL AND ALTERNATIVE HYPOTHESES II

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**
  - Note: always an equality!
- Alternative hypothesis must mathematically *contradict* the null hypothesis
  - e.g.  $\beta_1 \neq 2$  or  $\beta_1 > 20$  or  $\beta_1 \neq 0$
  - Note: always an inequality!
- Alternative hypotheses come in two forms:

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**
  - Note: always an equality!
- Alternative hypothesis must mathematically *contradict* the null hypothesis
  - e.g.  $\beta_1 \neq 2$  or  $\beta_1 > 20$  or  $\beta_1 \neq 0$
  - Note: always an inequality!
- Alternative hypotheses come in two forms:
  1. **One-sided alternative:**  $\beta_1 > H_0$  or  $\beta_1 < H_0$

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**
  - Note: always an equality!
- Alternative hypothesis must mathematically *contradict* the null hypothesis
  - e.g.  $\beta_1 \neq 2$  or  $\beta_1 > 20$  or  $\beta_1 \neq 0$
  - Note: always an inequality!
- Alternative hypotheses come in two forms:
  1. **One-sided alternative:**  $\beta_1 > H_0$  or  $\beta_1 < H_0$
  2. **Two-sided alternative:**  $\beta_1 \neq H_0$

- Null hypothesis assigns a value (or a range) to a population parameter
  - e.g.  $\beta_1 = 2$  or  $\beta_1 \leq 20$
  - **Most common null hypothesis is  $\beta_1 = 0 \implies X$  has no effect on  $Y$  (no slope for a line)**
  - Note: always an equality!
- Alternative hypothesis must mathematically *contradict* the null hypothesis
  - e.g.  $\beta_1 \neq 2$  or  $\beta_1 > 20$  or  $\beta_1 \neq 0$
  - Note: always an inequality!
- Alternative hypotheses come in two forms:
  1. **One-sided alternative:**  $\beta_1 > H_0$  or  $\beta_1 < H_0$
  2. **Two-sided alternative:**  $\beta_1 \neq H_0$ 
    - Note this means either  $\beta_1 < H_0$  or  $\beta_1 > H_0$

- All statistical hypothesis tests have the following components:

- All statistical hypothesis tests have the following components:

1. A null hypothesis,  $H_0$

- All statistical hypothesis tests have the following components:
  1. A null hypothesis,  $H_0$
  2. An alternative hypothesis,  $H_a$

- All statistical hypothesis tests have the following components:
  1. A **null hypothesis**,  $H_0$
  2. An **alternative hypothesis**,  $H_a$
  3. A **test statistic** to determine if we reject  $H_0$  when the statistic reaches a “critical value”



- All statistical hypothesis tests have the following components:
  1. A **null hypothesis**,  $H_0$
  2. An **alternative hypothesis**,  $H_a$
  3. A **test statistic** to determine if we reject  $H_0$  when the statistic reaches a “critical value”
    - Beyond the critical value is the “rejection region”, sufficient evidence to reject  $H_0$

- All statistical hypothesis tests have the following components:
  1. A **null hypothesis**,  $H_0$
  2. An **alternative hypothesis**,  $H_a$
  3. A **test statistic** to determine if we reject  $H_0$  when the statistic reaches a “critical value”
    - Beyond the critical value is the “rejection region”, sufficient evidence to reject  $H_0$
  4. A **conclusion** whether or not to reject  $H_0$  in favor of  $H_a$

- Any sample statistic (e.g.  $\hat{\beta}_1$ ) will rarely be exactly equal to the hypothesized population parameter (e.g.  $\beta_1$ )

- Any sample statistic (e.g.  $\hat{\beta}_1$ ) will rarely be exactly equal to the hypothesized population parameter (e.g.  $\beta_1$ )
- Difference between observed statistic and true parameter could be because:

- Any sample statistic (e.g.  $\hat{\beta}_1$ ) will rarely be exactly equal to the hypothesized population parameter (e.g.  $\beta_1$ )
- Difference between observed statistic and true parameter could be because:
  1. Parameter is *not* the hypothesized value ( $H_0$  is false)

- Any sample statistic (e.g.  $\hat{\beta}_1$ ) will rarely be exactly equal to the hypothesized population parameter (e.g.  $\beta_1$ )
- Difference between observed statistic and true parameter could be because:
  1. Parameter is *not* the hypothesized value ( $H_0$  is false)
  2. Parameter is truly the hypothesized value ( $H_0$  is true) but *sampling variability* gave us a different estimate

- Any sample statistic (e.g.  $\hat{\beta}_1$ ) will rarely be exactly equal to the hypothesized population parameter (e.g.  $\beta_1$ )
- Difference between observed statistic and true parameter could be because:
  1. Parameter is *not* the hypothesized value ( $H_0$  is false)
  2. Parameter is truly the hypothesized value ( $H_0$  is true) but *sampling variability* gave us a different estimate
- We cannot distinguish between these two possibilities with any certainty

- We can interpret our estimates probabilistically as committing one of two types of error:



- We can interpret our estimates probabilistically as committing one of two types of error:
  1. **Type I error (false positive)**: rejecting  $H_0$  when it is in fact true

- We can interpret our estimates probabilistically as committing one of two types of error:
  1. **Type I error (false positive):** rejecting  $H_0$  when it is in fact true
    - Believing we found an important result when there is truly no relationship

- We can interpret our estimates probabilistically as committing one of two types of error:
  1. **Type I error (false positive)**: rejecting  $H_0$  when it is in fact true
    - Believing we found an important result when there is truly no relationship
  2. **Type II error (false negative)**: failing to reject  $H_0$  when it is in fact false

- We can interpret our estimates probabilistically as committing one of two types of error:
  1. **Type I error (false positive)**: rejecting  $H_0$  when it is in fact true
    - Believing we found an important result when there is truly no relationship
  2. **Type II error (false negative)**: failing to reject  $H_0$  when it is in fact false
    - Believing we found nothing when there was truly a relationship to find

## TYPE I AND TYPE II ERRORS III

	$H_0$ is True	$H_0$ is False
Reject $H_0$	Type I Error False Positive	Correct Outcome True Positive
Don't Reject $H_0$	Correct Outcome True Negative	Type II Error False Negative

## TYPE I AND TYPE II ERRORS IV

	Defendant is Innocent	Defendant is Guilty
Convict "I think he's guilty"	Type I Error False Positive	Correct Outcome True Positive
Don't Convict "I think he's innocent"	Correct Outcome True Negative	Type II Error False Negative

- Depending on context, committing one type of error may be more serious than the other

## TYPE I AND TYPE II ERRORS IV

	Defendant is Innocent	Defendant is Guilty
Convict "I think he's guilty"	Type I Error False Positive	Correct Outcome True Positive
Don't Convict "I think he's innocent"	Correct Outcome True Negative	Type II Error False Negative

- Depending on context, committing one type of error may be more serious than the other
- Common law *presumes* the defendant is innocent and a jury judges whether the evidence presented against the defendant would be plausible *if the defendant were in fact innocent*

### Example

For each of the following scenarios, identify the Type I error, Type II error,  $\alpha$  and  $\beta$ , and which error is of greater consequence?



### Example

For each of the following scenarios, identify the Type I error, Type II error,  $\alpha$  and  $\beta$ , and which error is of greater consequence?

- $H_0$  : the victim of an automobile accident is alive when he arrives at the Emergency Room

### Example

For each of the following scenarios, identify the Type I error, Type II error,  $\alpha$  and  $\beta$ , and which error is of greater consequence?

- $H_0$  : the victim of an automobile accident is alive when he arrives at the Emergency Room
- $H_0$ : a rock climber's equipment is safe

### Example

For each of the following scenarios, identify the Type I error, Type II error,  $\alpha$  and  $\beta$ , and which error is of greater consequence?

- $H_0$  : the victim of an automobile accident is alive when he arrives at the Emergency Room
- $H_0$ : a rock climber's equipment is safe
- $H_0$ : a woman is not pregnant

### Example

For each of the following scenarios, identify the Type I error, Type II error,  $\alpha$  and  $\beta$ , and which error is of greater consequence?

- $H_0$  : the victim of an automobile accident is alive when he arrives at the Emergency Room
- $H_0$ : a rock climber's equipment is safe
- $H_0$ : a woman is not pregnant
- $H_0$ : a highway project will cost no more than \$10 million

### Example

For each of the following scenarios, identify the Type I error, Type II error,  $\alpha$  and  $\beta$ , and which error is of greater consequence?

- $H_0$  : the victim of an automobile accident is alive when he arrives at the Emergency Room
- $H_0$ : a rock climber's equipment is safe
- $H_0$ : a woman is not pregnant
- $H_0$ : a highway project will cost no more than \$10 million
- $H_0$ : an experimental cancer drug has a cure rate of at least 75%

- The **significance level,  $\alpha$** , is the probability of a **Type I error**

$$\alpha = P(\text{Reject } H_0 | H_0 \text{ is true})$$

$$\beta = P(\text{Don't reject } H_0 | H_0 \text{ is false})$$

- The **significance level**,  $\alpha$ , is the probability of a **Type I error**

$$\alpha = P(\text{Reject } H_0 | H_0 \text{ is true})$$

- The **confidence level** is defined as  $1 - \alpha$

$$\beta = P(\text{Don't reject } H_0 | H_0 \text{ is false})$$

- The **significance level,  $\alpha$** , is the probability of a **Type I error**

$$\alpha = P(\text{Reject } H_0 | H_0 \text{ is true})$$

- The **confidence level** is defined as  $1 - \alpha$
- We often specify in advance an  $\alpha$ -level (0.10, 0.05, 0.01) with associated confidence level (90%, 95%, 99%)

$$\beta = P(\text{Don't reject } H_0 | H_0 \text{ is false})$$



- The **significance level**,  $\alpha$ , is the probability of a **Type I error**

$$\alpha = P(\text{Reject } H_0 | H_0 \text{ is true})$$

- The **confidence level** is defined as  $1 - \alpha$
- We often specify in advance an  $\alpha$ -level (0.10, 0.05, 0.01) with associated confidence level (90%, 95%, 99%)
- The probability of a **Type II error** is defined as  $\beta$ :

$$\beta = P(\text{Don't reject } H_0 | H_0 \text{ is false})$$

	$H_0$ is True	$H_0$ is False
Reject $H_0$	Type I Error $\alpha$	Correct Outcome $(1 - \beta)$
Don't Reject $H_0$	Correct Outcome $(1 - \alpha)$	Type II Error $\beta$

- The statistical **power of the test** is  $1 - \beta$ , the probability of correctly rejecting  $H_0$  when  $H_0$  is in fact false (e.g. not convicting an innocent person)

$$\text{Power} = 1 - \beta = P(\text{Reject } H_0 | H_0 \text{ is false})$$

- The statistical **power of the test** is  $1 - \beta$ , the probability of correctly rejecting  $H_0$  when  $H_0$  is in fact false (e.g. not convicting an innocent person)

$$\text{Power} = 1 - \beta = P(\text{Reject } H_0 | H_0 \text{ is false})$$

- The  **$p$ -value** or **significance probability** is the probability that, given the null hypothesis is true, the test statistic from a random sample will be at least as extreme as the test statistic of our sample

- The statistical **power of the test** is  $1 - \beta$ , the probability of correctly rejecting  $H_0$  when  $H_0$  is in fact false (e.g. not convicting an innocent person)

$$\text{Power} = 1 - \beta = P(\text{Reject } H_0 | H_0 \text{ is false})$$

- The  **$p$ -value** or **significance probability** is the probability that, given the null hypothesis is true, the test statistic from a random sample will be at least as extreme as the test statistic of our sample
- If  $p < \alpha$ , the results are “**statistically significant**”

- After running our test, we need to make a *decision* between the competing hypotheses

- After running our test, we need to make a *decision* between the competing hypotheses
- Compare  $p$ -value with *pre-determined*  $\alpha$  (commonly,  $\alpha = 0.05$ , 95% confidence level)

- After running our test, we need to make a *decision* between the competing hypotheses
- Compare  $p$ -value with *pre-determined*  $\alpha$  (commonly,  $\alpha = 0.05$ , 95% confidence level)
  - If  $p < \alpha$ : **statistically significant** evidence sufficient to *reject*  $H_0$  in favor of  $H_a$



- After running our test, we need to make a *decision* between the competing hypotheses
- Compare  $p$ -value with *pre-determined*  $\alpha$  (commonly,  $\alpha = 0.05$ , 95% confidence level)
  - If  $p < \alpha$ : **statistically significant** evidence sufficient to *reject*  $H_0$  in favor of  $H_a$
  - If  $p \geq \alpha$ : *insufficient* evidence to reject  $H_0$


- After running our test, we need to make a *decision* between the competing hypotheses
- Compare  $p$ -value with *pre-determined*  $\alpha$  (commonly,  $\alpha = 0.05$ , 95% confidence level)
  - If  $p < \alpha$ : **statistically significant** evidence sufficient to *reject*  $H_0$  in favor of  $H_a$
  - If  $p \geq \alpha$ : *insufficient* evidence to reject  $H_0$ 
    - Note this does **not** mean  $H_0$  is true! We merely have *failed to reject*  $H_0$

DIGRESSION:  $p$ -VALUES AND THE  
PHILOSOPHY OF SCIENCE

---

“The null hypothesis is never proved or established, but is possibly disproved, in the course of experimentation. Every experiment may be said to exist only in order to give the facts a chance of disproving the null hypothesis.”

(1931). *The Design of Experiments*



Sir Ronald A. Fisher

(1890-1962)

- Modern philosophy of science is largely based off of hypothesis testing and **falsifiability**, which form the "Scientific Method"



- Modern philosophy of science is largely based off of hypothesis testing and **falsifiability**, which form the "Scientific Method"
- For something to be "scientific", it must be *falsifiable*, or at least *testable*



- Modern philosophy of science is largely based off of hypothesis testing and **falsifiability**, which form the "Scientific Method"
- For something to be "scientific", it must be *falsifiable*, or at least *testable*
- Hypotheses can be corroborated with evidence, but always *tentative* until falsified by data in suggesting an alternative hypothesis



- Modern philosophy of science is largely based off of hypothesis testing and **falsifiability**, which form the "Scientific Method"
- For something to be "scientific", it must be *falsifiable*, or at least *testable*
- Hypotheses can be corroborated with evidence, but always *tentative* until falsified by data in suggesting an alternative hypothesis
  - e.g. "**All swans are white**" is a hypothesis rejected upon discovery of a single black swan





- Modern philosophy of science is largely based off of hypothesis testing and **falsifiability**, which form the "Scientific Method"
- For something to be "scientific", it must be *falsifiable*, or at least *testable*
- Hypotheses can be corroborated with evidence, but always *tentative* until falsified by data in suggesting an alternative hypothesis
  - e.g. "**All swans are white**" is a hypothesis rejected upon discovery of a single black swan
- Note: economics is a very different kind of "science" with a different methodology!



### Caution

It is easy to misinterpret what statistical significance and  $p$ -values mean. **THE FOLLOWING ARE FALSE:**

### Caution

It is easy to misinterpret what statistical significance and  $p$ -values mean. **THE FOLLOWING ARE FALSE:**

- $p$  is the probability that the alternative hypothesis is true (We can never *prove* an alternative hypothesis, only tentatively reject a null hypothesis)

### Caution

It is easy to misinterpret what statistical significance and  $p$ -values mean. **THE FOLLOWING ARE FALSE:**

- $p$  is the probability that the alternative hypothesis is true (We can never *prove* an alternative hypothesis, only tentatively reject a null hypothesis)
- $p$  is the probability that the null hypothesis is false (We are not proving the null hypothesis false, only saying that it is very unlikely that under the null hypothesis, we obtain an event as rare as our sample)

### Caution

It is easy to misinterpret what statistical significance and  $p$ -values mean. **THE FOLLOWING ARE FALSE:**

- $p$  is the probability that the alternative hypothesis is true (We can never *prove* an alternative hypothesis, only tentatively reject a null hypothesis)
- $p$  is the probability that the null hypothesis is false (We are not proving the null hypothesis false, only saying that it is very unlikely that under the null hypothesis, we obtain an event as rare as our sample)
- $p$  is the probability that the observed effects were produced purely by random chance ( $p$  is computed under a specific model (assuming  $H_0$  is true))

### Caution

It is easy to misinterpret what statistical significance and  $p$ -values mean. **THE FOLLOWING ARE FALSE:**

- $p$  is the probability that the alternative hypothesis is true (We can never *prove* an alternative hypothesis, only tentatively reject a null hypothesis)
- $p$  is the probability that the null hypothesis is false (We are not proving the null hypothesis false, only saying that it is very unlikely that under the null hypothesis, we obtain an event as rare as our sample)
- $p$  is the probability that the observed effects were produced purely by random chance ( $p$  is computed under a specific model (assuming  $H_0$  is true))
- $p$  tells us how significant our finding is ( $p$  tells us nothing about the *size* or the *real world significance* of any effect deemed “statistically significant”)

- Again,  $p$  is the probability that, assuming the null hypothesis is true, we obtain (by pure random chance) a test statistic at least as extreme as the one we estimated for our sample
  - This will make more sense in context, when we discuss the nature of our test statistics

- Again,  $p$  is the probability that, assuming the null hypothesis is true, we obtain (by pure random chance) a test statistic at least as extreme as the one we estimated for our sample
  - This will make more sense in context, when we discuss the nature of our test statistics
- Remember a low  $p$ -value means **either** that the null hypothesis is true and a highly improbable event has occurred or that the null hypothesis is false (we don't know which!)




## STATISTICAL SIGNIFICANCE AND $p$ -VALUES




SMBC 1623

# STATISTICAL SIGNIFICANCE AND $p$ -VALUES

HOW SCIENCE REPORTING WORKS:




SMBC 1623




XKCD 882

## STATISTICAL SIGNIFICANCE AND $p$ -VALUES III




XKCD 882

## STATISTICAL SIGNIFICANCE AND $p$ -VALUES IV




XKCD 882



XKCD 882


## STATISTICAL SIGNIFICANCE AND $p$ -VALUES VI

- Consider what “95% significance” or  $\alpha = 0.05$  means:



## STATISTICAL SIGNIFICANCE AND $p$ -VALUES VI

- Consider what “95% significance” or  $\alpha = 0.05$  means:
  - If we repeat a procedure 20 times, we should *expect* 1/20 (5%) to produce a fluke result!





“The widespread use of “statistical significance” (generally interpreted as  $p \leq 0.05$ ) as a license for making a claim of a scientific finding (or implied truth) leads to considerable distortion of the scientific process.”



Wasserstein, Ronald L. and Nicole A. Lazar, (2016). “The ASA’s Statement on  $p$ -Values: Context, Process, and Purpose” *The American Statistician* 30(2): 129-133.

Morning Mix

## How, and why, a journalist tricked news outlets into thinking chocolate makes you thin

By Sarah Kaplan May 26, 2015



### Most Read

- 1 Black man fatally shot by Tulsa police was unarmed, chief says, as 'disturbing' video is released
- 2 Scientists uncovered a skeleton from the ancient world's most famous -- and mysterious -- shipwreck
- 3 Father of suspected bomber Ahmad Rahami says he had called the FBI about him
- 4 'You can sleep tonight knowing the Klan is awake.' Filers like these are showing up on lawns across the U.S.
- 5 Aren't more white people than black people killed by police? Yes, but no.

### Our Online Games

Play right from this page



Washington Post: How, and why, a journalist tricked news outlets into thinking chocolate makes you thin

## BACK TO OUR HYPOTHESIS TEST: THE TEST-STATISTIC

---

- We next consider the population distribution **assuming  $H_0$  is true** and calculate a **test statistic**, which takes the following form:

$$\text{test statistic} = \frac{\text{sample statistic} - \text{hypothesized value}}{\text{standard error of the statistic}}$$

- We next consider the population distribution **assuming  $H_0$  is true** and calculate a **test statistic**, which takes the following form:

$$\text{test statistic} = \frac{\text{sample statistic} - \text{hypothesized value}}{\text{standard error of the statistic}}$$

- We then compare our test statistic against a **critical value** to determine if we can reject  $H_0$

- We next consider the population distribution **assuming  $H_0$  is true** and calculate a **test statistic**, which takes the following form:

$$\text{test statistic} = \frac{\text{sample statistic} - \text{hypothesized value}}{\text{standard error of the statistic}}$$

- We then compare our test statistic against a **critical value** to determine if we can reject  $H_0$
- Essentially: **test to see how likely a sample statistic at least as extreme as our discovery is if  $H_0$  were true**

- We are testing our estimated  $\hat{\beta}_1$  against a null hypothesis, e.g.  $\beta_{1,0} = 0$

- We are testing our estimated  $\hat{\beta}_1$  against a null hypothesis, e.g.  $\beta_{1,0} = 0$
- It would be nice if we could use normal distribution, our test statistic would just be Z-score:

$$Z = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$




## DISTRIBUTION OF $H_0$ II


- We are testing our estimated  $\hat{\beta}_1$  against a null hypothesis, e.g.  $\beta_{1,0} = 0$
- It would be nice if we could use normal distribution, our test statistic would just be Z-score:

$$Z = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$

- **p-value**: area in the tail(s) of the distr. of  $\hat{\beta}_1$  under  $H_0$  beyond our Z score



- The **critical value**  $Z^*$  is determined by our  $\alpha$  level (e.g. 0.05)




Critical values of  $Z^*$  with rejection regions in red

---

<sup>2</sup>As you can see, the empirical 68-95-99.7% rule is very close, but not perfect!

## DISTRIBUTION OF $H_0$ III

- The **critical value**  $Z^*$  is determined by our  $\alpha$  level (e.g. 0.05)
- For a 2-sided alternative and  $\alpha = 0.05$ ,  $Z^* = 1.96$ <sup>2</sup>




Critical values of  $Z^*$  with rejection regions in red

---

<sup>2</sup>As you can see, the empirical 68-95-99.7% rule is very close, but not perfect!

## DISTRIBUTION OF $H_0$ III

- The **critical value**  $Z^*$  is determined by our  $\alpha$  level (e.g. 0.05)
- For a 2-sided alternative and  $\alpha = 0.05$ ,  $Z^* = 1.96^2$
- Any Z-score beyond  $\pm 1.96$  is in **rejection region**, sufficient evidence to reject  $H_0$



Critical values of  $Z^*$  with rejection regions in red

---

<sup>2</sup>As you can see, the empirical 68-95-99.7% rule is very close, but not perfect!

- It would be nice if we *could* just use the normal distribution, and run a **Z-test**, as described above

- It would be nice if we *could* just use the normal distribution, and run a **Z-test**, as described above
- **Central Limit Theorem** lets us if  $n \geq 30$  and we know the population distribution  $\mu, \sigma$

- It would be nice if we *could* just use the normal distribution, and run a **Z-test**, as described above
- **Central Limit Theorem** lets us if  $n \geq 30$  and we know the population distribution  $\mu, \sigma$
- We almost never know them...

## STUDENT'S $t$ -DISTRIBUTION

- Worked at Guinness testing beer quality



William Sealy Gosset  
(1876-1937)



## STUDENT'S $t$ -DISTRIBUTION

- Worked at Guinness testing beer quality
- Using normal distributions with small sample sizes did not yield accurate estimates



William Sealy Gosset  
(1876-1937)

## STUDENT'S $t$ -DISTRIBUTION

- Worked at Guinness testing beer quality
- Using normal distributions with small sample sizes did not yield accurate estimates
- Developed a new distribution, using the pseudonym “Student,” to publish, the Student's  $t$ -distribution



William Sealy Gosset  
(1876-1937)

- Instead of  $Z$ -scores, we use the  $t$ -**score**, which has the same intuition (# of standard deviations above/below the mean)

$$t \sim t_{n-k-1}$$

- Instead of Z-scores, we use the  **$t$ -score**, which has the same intuition (# of standard deviations above/below the mean)

$$t \sim t_{n-k-1}$$

- $t$ -scores follow a **Student's  $t$ -distribution** with  $n - k - 1$  degrees of freedom

- Instead of Z-scores, we use the  **$t$ -score**, which has the same intuition (# of standard deviations above/below the mean)

$$t \sim t_{n-k-1}$$

- $t$ -scores follow a **Student's  $t$ -distribution** with  $n - k - 1$  degrees of freedom
  - $k$ : number of variables (e.g.  $X$ 's)

- Instead of Z-scores, we use the  **$t$ -score**, which has the same intuition (# of standard deviations above/below the mean)

$$t \sim t_{n-k-1}$$

- $t$ -scores follow a **Student's  $t$ -distribution** with  $n - k - 1$  degrees of freedom
  - $k$ : number of variables (e.g.  $X$ 's)
  - Formally, **degrees of freedom** ( $df$  or  $\nu$ ) are the number of independent values used for the calculation of a statistic minus the number of other statistics used as intermediate steps

- Instead of Z-scores, we use the  **$t$ -score**, which has the same intuition (# of standard deviations above/below the mean)

$$t \sim t_{n-k-1}$$

- $t$ -scores follow a **Student's  $t$ -distribution** with  $n - k - 1$  degrees of freedom
  - $k$ : number of variables (e.g.  $X$ 's)
  - Formally, **degrees of freedom** ( $df$  or  $\nu$ ) are the number of independent values used for the calculation of a statistic minus the number of other statistics used as intermediate steps
  - Here, we *first* had to calculate two statistics ( $\hat{\beta}_0, \hat{\beta}_1$ ) with our sample *before* estimating the sampling distribution of  $\hat{\beta}_1$  for hypothesis testing, thus  $df = n - 2$

- Instead of Z-scores, we use the **t-score**, which has the same intuition (# of standard deviations above/below the mean)

$$t \sim t_{n-k-1}$$

- t-scores follow a **Student's t-distribution** with  $n - k - 1$  degrees of freedom
  - $k$ : number of variables (e.g.  $X$ 's)
  - Formally, **degrees of freedom** ( $df$  or  $\nu$ ) are the number of independent values used for the calculation of a statistic minus the number of other statistics used as intermediate steps
  - Here, we *first* had to calculate two statistics ( $\hat{\beta}_0, \hat{\beta}_1$ ) with our sample *before* estimating the sampling distribution of  $\hat{\beta}_1$  for hypothesis testing, thus  $df = n - 2$
- t-distribution looks normal-ish (symmetric, unimodal, mean=0), but more mass in the tails




- Instead of Z-scores, we use the **t-score**, which has the same intuition (# of standard deviations above/below the mean)



$$t \sim t_{n-k-1}$$

- t-scores follow a **Student's t-distribution** with  $n - k - 1$  degrees of freedom
  - $k$ : number of variables (e.g.  $X$ 's)
  - Formally, **degrees of freedom** ( $df$  or  $\nu$ ) are the number of independent values used for the calculation of a statistic minus the number of other statistics used as intermediate steps
  - Here, we *first* had to calculate two statistics ( $\hat{\beta}_0, \hat{\beta}_1$ ) with our sample *before* estimating the sampling distribution of  $\hat{\beta}_1$  for hypothesis testing, thus  $df = n - 2$
- t-distribution looks normal-ish (symmetric, unimodal, mean=0), but more mass in the tails
- Exact shape of  $t$  depends on  $df$ : as  $\uparrow df$ ,  $t \rightarrow$  Normal distribution

## $t$ -DISTRIBUTIONS



# CALCULATING $t$ -SCORES: OLD-FASHIONED WAY

Two tail probability One tail probability		0.20 0.10	0.10 0.05	0.05 0.025
Table T				
Values of $t_{\alpha}$				
 <p>Two tails</p>	1	3.078	6.314	12.706
	2	1.886	2.920	4.303
	3	1.638	2.353	3.182
	4	1.533	2.132	2.776
	5	1.476	2.015	2.571
	6	1.440	1.943	2.447
	7	1.415	1.895	2.365
	8	1.397	1.860	2.306
	9	1.383	1.833	2.262
	10	1.372	1.812	2.228
 <p>One tail</p>	11	1.363	1.796	2.201
	12	1.356	1.782	2.179
	13	1.350	1.771	2.160
	14	1.345	1.761	2.145
	15	1.341	1.753	2.131
	16	1.337	1.746	2.120
	17	1.333	1.740	2.110
	18	1.330	1.734	2.101
	19	1.328	1.729	2.093
	$\vdots$	$\vdots$	$\vdots$	$\vdots$
	$\infty$	1.282	1.645	1.960
Confidence levels		80%	90%	95%

```
# use pt() command, needs t value and df  
pt(2,df=5) #probability of  $t > 2$  with 5 df
```

```
## [1] 0.9490303
```

```
pt(2,df=40) # probability of  $t > 2$  with 40 df
```

```
## [1] 0.9738388
```

```
pt(2, df=100) # probability of  $t > 2$  with 100 df
```

```
## [1] 0.9758939
```

```
pnorm(2, mean=0, sd=1) # compare to normal distribution!
```

- So our **test statistic** is a  **$t$ -score** (instead of  $Z$ -score)

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$

- So our **test statistic** is a  **$t$ -score** (instead of  $Z$ -score)

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$

- We then compare  $t$  to the critical value of  $t^*$  determined by our  $\alpha$ -level and the  $df$  for our  $t$ -distribution ( $n - k - 1$ )

- So our **test statistic** is a  **$t$ -score** (instead of  $Z$ -score)

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$

- We then compare  $t$  to the critical value of  $t^*$  determined by our  $\alpha$ -level and the  $df$  for our  $t$ -distribution ( $n - k - 1$ )
  - Note: there will be a unique critical value for every value of  $n - k - 1$ !

- So our **test statistic** is a  **$t$ -score** (instead of  $Z$ -score)

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$

- We then compare  $t$  to the critical value of  $t^*$  determined by our  $\alpha$ -level and the  $df$  for our  $t$ -distribution ( $n - k - 1$ )
  - Note: there will be a unique critical value for every value of  $n - k - 1$ !
  - R determines the critical  $t^*$  automatically with regression



- So our **test statistic** is a  **$t$ -score** (instead of  $Z$ -score)

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$

- We then compare  $t$  to the critical value of  $t^*$  determined by our  $\alpha$ -level and the  $df$  for our  $t$ -distribution ( $n - k - 1$ )
  - Note: there will be a unique critical value for every value of  $n - k - 1$ !
  - R determines the critical  $t^*$  automatically with regression
- $p\text{-value} = P(t < T)$




- So our **test statistic** is a  **$t$ -score** (instead of  $Z$ -score)

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$

- We then compare  $t$  to the critical value of  $t^*$  determined by our  $\alpha$ -level and the  $df$  for our  $t$ -distribution ( $n - k - 1$ )
  - Note: there will be a unique critical value for every value of  $n - k - 1$ !
  - R determines the critical  $t^*$  automatically with regression
- $p\text{-value} = P(t < T)$
- Reject  $H_0$  if  $p\text{-value} < \alpha$

## HYPOTHESIS TESTING WITH $t$ -DISTRIBUTION II

Depending on the desired alternative hypothesis:

Alternative	$p$ -value	PDF
$H_a : \beta_1 > \beta_{1,0}$	$P(T \geq t)$	
$H_a : \beta_1 < \beta_{1,0}$	$P(T \leq t)$	
$H_a : \beta_1 \neq \beta_{1,0}$	$2P(T \geq  t )$	

### Example

We have an estimated regression line:

$$\widehat{\text{Test Score}} = 689.93 - 2.28 \text{ STR}$$

(9.47)    (0.48)

- Regression reporting format: Coefficients with their (standard errors) beneath them

### Example

We have an estimated regression line:

$$\widehat{\text{Test Score}} = 689.93 - 2.28 \text{ STR}$$

(9.47)    (0.48)

- Regression reporting format: Coefficients with their (standard errors) beneath them
- Let  $H_0 : \beta_1 = 0$ ,  $H_1 : \beta_1 \neq 0$  (two-sided alternative)

### Example

We have an estimated regression line:

$$\widehat{\text{Test Score}} = 689.93 - 2.28 \text{ STR} \\ (9.47) \quad (0.48)$$

- Regression reporting format: Coefficients with their (standard errors) beneath them
- Let  $H_0 : \beta_1 = 0$ ,  $H_1 : \beta_1 \neq 0$  (two-sided alternative)
- $t$ -statistic:

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)}$$

### Example

We have an estimated regression line:

$$\widehat{\text{Test Score}} = 689.93 - 2.28 \text{ STR}$$
$$(9.47) \quad (0.48)$$

- Regression reporting format: Coefficients with their (standard errors) beneath them
- Let  $H_0 : \beta_1 = 0$ ,  $H_1 : \beta_1 \neq 0$  (two-sided alternative)
- $t$ -statistic:

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)} = \frac{-2.28 - 0}{0.48}$$

### Example

We have an estimated regression line:

$$\widehat{\text{Test Score}} = 689.93 - 2.28 \text{ STR}$$
$$(9.47) \quad (0.48)$$

- Regression reporting format: Coefficients with their (standard errors) beneath them
- Let  $H_0 : \beta_1 = 0$ ,  $H_1 : \beta_1 \neq 0$  (two-sided alternative)
- $t$ -statistic:

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)} = \frac{-2.28 - 0}{0.48} = -4.75$$



### Example

We have an estimated regression line:

$$\widehat{\text{Test Score}} = 689.93 - 2.28 \text{ STR}$$
$$(9.47) \quad (0.48)$$

- Regression reporting format: Coefficients with their (standard errors) beneath them
- Let  $H_0 : \beta_1 = 0$ ,  $H_1 : \beta_1 \neq 0$  (two-sided alternative)
- $t$ -statistic:

$$t = \frac{\hat{\beta}_1 - \beta_{1,0}}{SE(\hat{\beta}_1)} = \frac{-2.28 - 0}{0.48} = -4.75$$

```
# calculate p-value for t=-4.75, df=418
```

```
2*pt(-4.75,df=418) # x2 because we want both tails!
```

```
summary(school.regression)
```

```
##
## Call:
## lm(formula = testscr ~ str, data = CASchool)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -47.727 -14.251   0.483  12.822  48.540
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  698.9330     9.4675   73.825  < 2e-16 ***
## str          -2.2798     0.4798   -4.751 2.78e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 18.58 on 418 degrees of freedom
## Multiple R-squared:  0.05124,    Adjusted R-squared:  0.04897
## F-statistic: 22.58 on 1 and 418 DF,  p-value: 2.783e-06
```

- If  $|\hat{\beta}_k| > 2 \times SE(\hat{\beta}_k)$ , the estimate is significant

- If  $|\hat{\beta}_k| > 2 \times SE(\hat{\beta}_k)$ , the estimate is significant

$$\widehat{\text{Test Score}} = 689.93 - 2.28 \text{ STR} \\ (9.47) \quad (0.48)$$

- Since essentially  $t = \frac{\hat{\beta}_k}{SE(\hat{\beta}_k)}$  and we roughly want  $t \geq 2$  for 95% confidence level ( $\alpha=0.05$ )

- Random sampling variation implies that our estimated  $\hat{\beta}_1$  will vary slightly from sample to sample

- Random sampling variation implies that our estimated  $\hat{\beta}_1$  will vary slightly from sample to sample
- We can also calculate a **confidence interval** of  $\hat{\beta}_1$  values that contain the true slope  $\beta_0$  with a specified probability  $(1 - \alpha)$

- Random sampling variation implies that our estimated  $\hat{\beta}_1$  will vary slightly from sample to sample
- We can also calculate a **confidence interval** of  $\hat{\beta}_1$  values that contain the true slope  $\beta_0$  with a specified probability  $(1 - \alpha)$
- In general, a confidence interval takes the form:

(point estimate  $-$  margin of error, point estimate  $+$  margin of error)


- Recall the **empirical 68-95-99.7 rule**: approximately 95% of a normal distribution occurs within 2 standard deviations of the mean



## CONFIDENCE INTERVALS: INTUITION


- Recall the **empirical 68-95-99.7 rule**: approximately 95% of a normal distribution occurs within 2 standard deviations of the mean
- Thus, in 95% of samples, the true slope ( $\beta_1$ ) is likely to fall within about two standard errors of our estimated slope ( $\hat{\beta}_1$ )

$$(\hat{\beta}_1 - 2se(\hat{\beta}_1), \hat{\beta}_1 + 2se(\hat{\beta}_1))$$



## CONFIDENCE INTERVALS


- We need to find the  $t$ -score<sup>3</sup>,  $t_{\frac{\alpha}{2}}$  that puts an area equal to CL in the middle of the  $t$ -distribution  $t \sim t_{n-k-1}$



<sup>3</sup>If we knew the population distribution, and if  $n \geq 30$ , we could just use Z-score and normal distribution

## CONFIDENCE INTERVALS


- We need to find the  $t$ -score<sup>3</sup>,  $t_{\frac{\alpha}{2}}$  that puts an area equal to CL in the middle of the  $t$ -distribution  $t \sim t_{n-k-1}$
- The specific  $t_{\frac{\alpha}{2}}$  is (again) the **critical value** for a given confidence level



<sup>3</sup>If we knew the population distribution, and if  $n \geq 30$ , we could just use Z-score and normal distribution

## CONFIDENCE INTERVALS


- We need to find the  $t$ -score<sup>3</sup>,  $t_{\frac{\alpha}{2}}$  that puts an area equal to CL in the middle of the  $t$ -distribution  $t \sim t_{n-k-1}$
- The specific  $t_{\frac{\alpha}{2}}$  is (again) the **critical value** for a given confidence level
- $CL = 1 - \alpha$ , so  $\alpha$  is the area split equally across the two tails of the distribution ( $\frac{\alpha}{2}$ ) in each tail



<sup>3</sup>If we knew the population distribution, and if  $n \geq 30$ , we could just use Z-score and normal distribution

### Example

For  $\alpha = 0.05$  (or CL=95%) and large  $n$ ,  $t^* = Z^* = 1.96$



- The **margin of error (MOE)** is the critical value of  $t$  times the standard error of  $\hat{\beta}_1$

- The **margin of error (MOE)** is the critical value of  $t$  times the standard error of  $\hat{\beta}_1$

$$MOE = t_{\frac{\alpha}{2}} se(\hat{\beta}_1)$$

- For example, for 95% confidence and large  $n$ :

- The **margin of error (MOE)** is the critical value of  $t$  times the standard error of  $\hat{\beta}_1$

$$MOE = t_{\frac{\alpha}{2}} se(\hat{\beta}_1)$$

- For example, for 95% confidence and large  $n$ :

$$MOE = 1.96 se(\hat{\beta}_1)$$

- A confidence interval for  $\beta_1$  is then:



- The **margin of error (MOE)** is the critical value of  $t$  times the standard error of  $\hat{\beta}_1$

$$MOE = t_{\frac{\alpha}{2}} se(\hat{\beta}_1)$$

- For example, for 95% confidence and large  $n$ :

$$MOE = 1.96 se(\hat{\beta}_1)$$

- A confidence interval for  $\beta_1$  is then:

$$\left( \hat{\beta}_1 - t_{\frac{\alpha}{2}} se(\hat{\beta}_1), \quad \hat{\beta}_1 + t_{\frac{\alpha}{2}} se(\hat{\beta}_1) \right)$$

- “We estimate with  $[1 - \alpha]$  confidence that the true population mean is between [...] and [...].”

- For large  $n$ :

Confidence Level	Critical Value	Confidence Interval
90%	1.64	$\hat{\beta}_1 \pm 1.64se(\hat{\beta}_1)$
90%	1.96	$\hat{\beta}_1 \pm 1.96se(\hat{\beta}_1)$
90%	2.58	$\hat{\beta}_1 \pm 2.58se(\hat{\beta}_1)$

## CONFIDENCE INTERVALS IN R

- Generate confidence intervals with `confint()` command

```
confint(school.regression, level=0.90) # 90% confidence
```

```
##                5 %        95 %  
## (Intercept) 683.325725 714.540180  
## str         -3.070804  -1.488812
```

```
confint(school.regression, level=0.95) # 95% confidence
```

```
##                2.5 %       97.5 %  
## (Intercept) 680.32313 717.542779  
## str         -3.22298  -1.336637
```

```
confint(school.regression, level=0.99) # 99% confidence
```

```
##                0.5 %       99.5 %  
## (Intercept) 674.434473 723.431432  
## str         -3.521425  -1.038191
```

## REPORTING REGRESSION OUTPUTS WITH `stargazer`

---

- Regressions are reported with a **regression table** (especially when we have many models!)

- Regressions are reported with a **regression table** (especially when we have many models!)
- You can make this by hand, but it's much faster with a great package called **`stargazer`**

## REPORTING REGRESSION OUTPUTS WITH `stargazer`

- Regressions are reported with a **regression table** (especially when we have many models!)
- You can make this by hand, but it's much faster with a great package called **`stargazer`**

	Test Score
Class Size	-2.280*** (0.480)
Constant	698.933*** (9.467)
<i>N</i>	420
$R^2$	0.051
Residual Std. Error	18.581 (df = 418)

Notes:

\*\*\* Significant at the 1 percent level.

\*\* Significant at the 5 percent level.

\* Significant at the 10 percent level.

- Basic **stargazer** syntax is simple

```
library("stargazer") # load stargazer (you will need to install first!)  
stargazer(reg.name,type="type")
```



- Basic **stargazer** syntax is simple
  - First you need to load with `library()`, and install if the first time!

```
library("stargazer") # load stargazer (you will need to install first!)  
stargazer(reg.name,type="type")
```

- Basic **stargazer** syntax is simple
  - First you need to load with `library()`, and install if the first time!
  - First include the name of the regression object to print (e.g. `reg.name`)

```
library("stargazer") # load stargazer (you will need to install first!)  
stargazer(reg.name,type="type")
```

- Basic **stargazer** syntax is simple
  - First you need to load with `library()`, and install if the first time!
  - First include the name of the regression object to print (e.g. `reg.name`)
  - Then specify the type of output ("`html`", "`latex`" (for pdf), or "`text`")

```
library("stargazer") # load stargazer (you will need to install first!)  
stargazer(reg.name, type="type")
```

- `stargazer` allows for a lot of customization, e.g. the code for the above table:

```
library("stargazer")
stargazer(school.regression, type="latex", header=FALSE, style="qje",
          dep.var.labels = "Test Score", covariate.labels = c("Class Size"),
          title="Regression Results", float=FALSE, font.size = "scriptsize",
          omit.stat=c(adj.rsq,f))
```

- The raw output looks confusing, code that renders directly in html or pdf (via latex)

```
##
## \begingroup
## \scriptsize
## \begin{tabular}{@{\extracolsep{5pt}}lc}
## \[-1.8ex\]\hline
## \hline \[-1.8ex]
## \[-1.8ex] & Test Score \\\
## \hline \[-1.8ex]
## Class Size &  $-\$2.280^{***}$  \\\
## & (0.480) \\\
## & \\\
## Constant &  $698.933^{***}$  \\\
## & (9.467) \\\
## & \\\
## \textit{N} & 420 \\\
##  $R^2$  & 0.051 \\\
## Residual Std. Error & 18.581 (df = 418) \\\
## \hline
```

## stargazer: RENDERING DIRECTLY AS TEXT

```
stargazer(school.regression, type="text")
```

```
##
## =====
##                Dependent variable:
##                -----
##                testscr
## -----
## str                -2.280***
##                  (0.480)
##
## Constant           698.933***
##                  (9.467)
## -----
## Observations              420
## R2                       0.051
## Adjusted R2              0.049
## Residual Std. Error    18.581 (df = 418)
```