

intervention for P3 and P1 and a level trend at baseline and an increasing therapeutic trend during the intervention for P2. A stability envelope was applied to trend lines and showed that the data were stable at baseline for P3 but variable during the intervention, while data were stable in both phases for P1 and P2. P3's mean accuracy scores increased from 51.67% at baseline (range 55–45%) to 83.33% (range 100–65%) during the intervention, and maintenance was 100%. P1's mean accuracy scores increased from 36.8% (range 45–35%) at baseline to 80% (range 100–65%) during the intervention, and maintenance was 100%. P2's mean accuracy scores increased from 40.71% (range 45–35%) at baseline to 82% (range 85–65%) during the intervention, and maintenance was 75%.

Between-Condition Analysis: Evaluation of the behaviour change between conditions indicated improvements for the intervention relative to the baseline for all three participants. For P2, a change in performance across conditions meant a change from a level, deteriorating trend at baseline to an accelerating, improving trend during the intervention. For P3 and P1, performance was accelerating and improving across both conditions. All level change measures indicated a positive improving behaviour change across conditions for all participants. The relative level change, absolute level change, mean level change, and median level change were calculated to demonstrate between-condition effects: P3's scores were +20, +10, +30, and +31.13, respectively; P1's scores were +34.5, +30, +40, and +43.2, respectively; and P2's scores were +26, +30, +30, and +36.29, respectively. Figures 1–3 provide a visual display of the data. No overlapping data (Table 2) was present for any participants for ADL1 (PND = 100%), indicating that the intervention was very effective (Scruggs & Mastropieri, 1998). Tables displaying the full set of visual analysis data can be accessed via the Supplementary Materials, Tables S1–S3.




Figure 1. Cont.