

Introduction to Latent Variable Models


Piyush Rai

Probabilistic Machine Learning (CS772A)

Aug 29, 2017


A Simple Generative Model

- All observations $\{\mathbf{x}_1, \dots, \mathbf{x}_N\}$ generated from a distribution $p(\mathbf{x}|\theta)$



A Simple Generative Model


- All observations $\{\mathbf{x}_1, \dots, \mathbf{x}_N\}$ generated from a distribution $p(\mathbf{x}|\theta)$



- Unknowns: Parameters θ of the assumed data distribution $p(\mathbf{x}|\theta)$

A Simple Generative Model


- All observations $\{\mathbf{x}_1, \dots, \mathbf{x}_N\}$ generated from a distribution $p(\mathbf{x}|\theta)$



- Unknowns: Parameters θ of the assumed data distribution $p(\mathbf{x}|\theta)$
- Many ways to estimate the parameters (MLE, MAP, or Bayesian inference)


Generative Model with Latent Variables

- Assume each observation \mathbf{x}_n to be associated with a latent variable \mathbf{z}_n



Generative Model with Latent Variables


- Assume each observation \mathbf{x}_n to be associated with a latent variable \mathbf{z}_n



- In this “latent variable model” of data generation, data \mathbf{x} also depends some latent variable(s) \mathbf{z}
- \mathbf{z}_n is akin to a latent representation or “encoding” of \mathbf{x}_n ; controls what data “looks like”.

Generative Model with Latent Variables


- Assume each observation \mathbf{x}_n to be associated with a latent variable \mathbf{z}_n



- In this “latent variable model” of data generation, data \mathbf{x} also depends some latent variable(s) \mathbf{z}
- \mathbf{z}_n is akin to a latent representation or “encoding” of \mathbf{x}_n ; controls what data “looks like”. E.g.,
 - $\mathbf{z}_n \in \{1, \dots, K\}$ denotes the cluster \mathbf{x}_n belongs to

Generative Model with Latent Variables


- Assume each observation \mathbf{x}_n to be associated with a latent variable \mathbf{z}_n



- In this “latent variable model” of data generation, data \mathbf{x} also depends some latent variable(s) \mathbf{z}
- \mathbf{z}_n is akin to a latent representation or “encoding” of \mathbf{x}_n ; controls what data “looks like”. E.g.,
 - $\mathbf{z}_n \in \{1, \dots, K\}$ denotes the cluster \mathbf{x}_n belongs to
 - $\mathbf{z}_n \in \mathbb{R}^K$ denotes a low-dimensional latent representation or latent “code” for \mathbf{x}_n

Generative Model with Latent Variables


- Assume each observation \mathbf{x}_n to be associated with a latent variable \mathbf{z}_n



- In this “latent variable model” of data generation, data \mathbf{x} also depends some latent variable(s) \mathbf{z}
- \mathbf{z}_n is akin to a latent representation or “encoding” of \mathbf{x}_n ; controls what data “looks like”. E.g.,
 - $\mathbf{z}_n \in \{1, \dots, K\}$ denotes the cluster \mathbf{x}_n belongs to
 - $\mathbf{z}_n \in \mathbb{R}^K$ denotes a low-dimensional latent representation or latent “code” for \mathbf{x}_n
- Unknowns: $\{\mathbf{z}_1, \dots, \mathbf{z}_N\}$, and (θ, ϕ) .

Generative Model with Latent Variables


- Assume each observation \mathbf{x}_n to be associated with a latent variable \mathbf{z}_n



- In this “latent variable model” of data generation, data \mathbf{x} also depends some latent variable(s) \mathbf{z}
- \mathbf{z}_n is akin to a latent representation or “encoding” of \mathbf{x}_n ; controls what data “looks like”. E.g.,
 - $\mathbf{z}_n \in \{1, \dots, K\}$ denotes the cluster \mathbf{x}_n belongs to
 - $\mathbf{z}_n \in \mathbb{R}^K$ denotes a low-dimensional latent representation or latent “code” for \mathbf{x}_n
- Unknowns: $\{\mathbf{z}_1, \dots, \mathbf{z}_N\}$, and (θ, ϕ) . \mathbf{z}_n ’s called “local” variables; (θ, ϕ) called “global” variables

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n (o/w it is simply generative classification)

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n (o/w it is simply **generative classification**)
- Assume a latent random variable $\mathbf{z}_n \in \{1, \dots, K\}$ denotes which component generated \mathbf{x}_n

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n (o/w it is simply [generative classification](#))
- Assume a latent random variable $\mathbf{z}_n \in \{1, \dots, K\}$ denotes which component generated \mathbf{x}_n
- Here is a simple [generative story](#) for each \mathbf{x}_n , $n = 1, 2, \dots, N$

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n (o/w it is simply **generative classification**)
- Assume a latent random variable $\mathbf{z}_n \in \{1, \dots, K\}$ denotes which component generated \mathbf{x}_n
- Here is a simple **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$
 - First choose a mixture component $\mathbf{z}_n \in \{1, 2, \dots, K\}$ as $\mathbf{z}_n \sim \text{multinoulli}(\mathbf{z}|\phi)$

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n (o/w it is simply **generative classification**)
- Assume a latent random variable $\mathbf{z}_n \in \{1, \dots, K\}$ denotes which component generated \mathbf{x}_n
- Here is a simple **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$
 - First choose a mixture component $\mathbf{z}_n \in \{1, 2, \dots, K\}$ as $\mathbf{z}_n \sim \text{multinoulli}(\mathbf{z}|\phi)$
 - Now generate \mathbf{x}_n from that mixture component as $\mathbf{x}_n \sim p(\mathbf{x}|\theta_{\mathbf{z}_n})$

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n (o/w it is simply **generative classification**)
- Assume a latent random variable $\mathbf{z}_n \in \{1, \dots, K\}$ denotes which component generated \mathbf{x}_n
- Here is a simple **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$
 - First choose a mixture component $\mathbf{z}_n \in \{1, 2, \dots, K\}$ as $\mathbf{z}_n \sim \text{multinoulli}(\mathbf{z}|\phi)$
 - Now generate \mathbf{x}_n from that mixture component as $\mathbf{x}_n \sim p(\mathbf{x}|\theta_{\mathbf{z}_n})$
- Goal: Given data $\{\mathbf{x}_n\}_{n=1}^N$, learn the K distributions $(\theta_1, \dots, \theta_K)$ and latent variables $\mathbf{z}_1, \dots, \mathbf{z}_N$

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n (o/w it is simply **generative classification**)
- Assume a latent random variable $\mathbf{z}_n \in \{1, \dots, K\}$ denotes which component generated \mathbf{x}_n
- Here is a simple **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$
 - First choose a mixture component $\mathbf{z}_n \in \{1, 2, \dots, K\}$ as $\mathbf{z}_n \sim \text{multinoulli}(\mathbf{z}|\phi)$
 - Now generate \mathbf{x}_n from that mixture component as $\mathbf{x}_n \sim p(\mathbf{x}|\theta_{\mathbf{z}_n})$
- Goal: Given data $\{\mathbf{x}_n\}_{n=1}^N$, learn the K distributions $(\theta_1, \dots, \theta_K)$ and latent variables $\mathbf{z}_1, \dots, \mathbf{z}_N$
- If each $p(\mathbf{x}|\theta_k)$ is a Gaussian \Rightarrow **Gaussian Mixture Model**

A Motivating Example: Mixture Model


- Assume data $\{\mathbf{x}_n\}_{n=1}^N$ was generated from a mixture of K distributions $p(\mathbf{x}|\theta_1), \dots, p(\mathbf{x}|\theta_K)$



- Don't know mixture component generated each \mathbf{x}_n (o/w it is simply **generative classification**)
- Assume a latent random variable $\mathbf{z}_n \in \{1, \dots, K\}$ denotes which component generated \mathbf{x}_n
- Here is a simple **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$
 - First choose a mixture component $\mathbf{z}_n \in \{1, 2, \dots, K\}$ as $\mathbf{z}_n \sim \text{multinoulli}(\mathbf{z}|\phi)$
 - Now generate \mathbf{x}_n from that mixture component as $\mathbf{x}_n \sim p(\mathbf{x}|\theta_{\mathbf{z}_n})$
- Goal: Given data $\{\mathbf{x}_n\}_{n=1}^N$, learn the K distributions $(\theta_1, \dots, \theta_K)$ and latent variables $\mathbf{z}_1, \dots, \mathbf{z}_N$
- If each $p(\mathbf{x}|\theta_k)$ is a Gaussian \Rightarrow **Gaussian Mixture Model** (used for probabilistic or “soft” clustering)


Another Motivating Example: Latent Factor Model

- Assume data $\mathbf{x}_n \in \mathbb{R}^D$ generated from a low-dimensional latent factor $\mathbf{z}_n \in \mathbb{R}^K$



Another Motivating Example: Latent Factor Model


- Assume data $\mathbf{x}_n \in \mathbb{R}^D$ generated from a low-dimensional latent factor $\mathbf{z}_n \in \mathbb{R}^K$



- The z to x map can be a linear/nonlinear transformation

Another Motivating Example: Latent Factor Model


- Assume data $\mathbf{x}_n \in \mathbb{R}^D$ generated from a low-dimensional latent factor $\mathbf{z}_n \in \mathbb{R}^K$



- The \mathbf{z} to \mathbf{x} map can be a linear/nonlinear transformation
- Consider the following **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$

Another Motivating Example: Latent Factor Model


- Assume data $\mathbf{x}_n \in \mathbb{R}^D$ generated from a low-dimensional latent factor $\mathbf{z}_n \in \mathbb{R}^K$



- The \mathbf{z} to \mathbf{x} map can be a linear/nonlinear transformation
- Consider the following **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$
 - First generate \mathbf{z}_n from a K -dim distr. as $\mathbf{z}_n \sim p(\mathbf{z}|\phi)$

Another Motivating Example: Latent Factor Model


- Assume data $\mathbf{x}_n \in \mathbb{R}^D$ generated from a low-dimensional latent factor $\mathbf{z}_n \in \mathbb{R}^K$



- The z to x map can be a linear/nonlinear transformation
- Consider the following **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$
 - First generate \mathbf{z}_n from a K -dim distr. as $\mathbf{z}_n \sim p(\mathbf{z}|\phi)$
 - Now generate \mathbf{x}_n from a D -dim distr. as $\mathbf{x}_n \sim p(\mathbf{x}|\mathbf{z}_n, \theta)$

Another Motivating Example: Latent Factor Model


- Assume data $\mathbf{x}_n \in \mathbb{R}^D$ generated from a low-dimensional latent factor $\mathbf{z}_n \in \mathbb{R}^K$



- The \mathbf{z} to \mathbf{x} map can be a linear/nonlinear transformation
- Consider the following **generative story** for each \mathbf{x}_n , $n = 1, 2, \dots, N$
 - First generate \mathbf{z}_n from a K -dim distr. as $\mathbf{z}_n \sim p(\mathbf{z}|\phi)$
 - Now generate \mathbf{x}_n from a D -dim distr. as $\mathbf{x}_n \sim p(\mathbf{x}|\mathbf{z}_n, \theta)$
- If $p(\mathbf{z}|\phi)$ and $p(\mathbf{x}|\mathbf{z}, \theta)$ are Gaussians and \mathbf{z} to \mathbf{x} map linear \Rightarrow **factor analysis** or **probabilistic PCA**


Other Examples..

- State-space latent variable models (e.g., Hidden Markov Models, Kalman Filters)




Other Examples..

- State-space latent variable models (e.g., Hidden Markov Models, Kalman Filters)




- Latent variable models for “relational data” (e.g., ratings matrix, graph, etc.)




Other Examples..

- State-space latent variable models (e.g., Hidden Markov Models, Kalman Filters)




- Latent variable models for “relational data” (e.g., ratings matrix, graph, etc.)



Other Examples..


- Semi-supervised generative classification: Some training inputs can be unlabeled
 - These “missing” labels can be treated as latent variables and inferred



Latent Variable Models for Clustering and Density Estimation


Gaussian Mixture Model (GMM)

- A generative model for data clustering. Assume data generated from a mixture of K Gaussians



Gaussian Mixture Model (GMM)


- A generative model for data clustering. Assume data generated from a mixture of K Gaussians



- Let $\pi = (\pi_1, \pi_2, \dots, \pi_K)$ with $\sum_{k=1}^K \pi_k = 1$ denote the “mixing weights” of the K Gaussians

Gaussian Mixture Model (GMM)


- A generative model for data clustering. Assume data generated from a mixture of K Gaussians



- Let $\pi = (\pi_1, \pi_2, \dots, \pi_K)$ with $\sum_{k=1}^K \pi_k = 1$ denote the “mixing weights” of the K Gaussians
- Intuitively $\pi_k \in (0, 1)$ is the fraction of data “contributed” by the k -th Gaussian

Gaussian Mixture Model (GMM)

- A generative model for data clustering. Assume data generated from a mixture of K Gaussians




- Let $\pi = (\pi_1, \pi_2, \dots, \pi_K)$ with $\sum_{k=1}^K \pi_k = 1$ denote the “mixing weights” of the K Gaussians
- Intuitively $\pi_k \in (0, 1)$ is the fraction of data “contributed” by the k -th Gaussian
- π_k is the prior probability that \mathbf{x}_n comes from the k -th Gaussian (i.e., cluster id $\mathbf{z}_n = k$)

$$p(\mathbf{z}_n = k | \pi) = \pi_k$$

Gaussian Mixture Model (GMM)

- A generative model for data clustering. Assume data generated from a mixture of K Gaussians




- Let $\pi = (\pi_1, \pi_2, \dots, \pi_K)$ with $\sum_{k=1}^K \pi_k = 1$ denote the “mixing weights” of the K Gaussians
- Intuitively $\pi_k \in (0, 1)$ is the fraction of data “contributed” by the k -th Gaussian
- π_k is the prior probability that \mathbf{x}_n comes from the k -th Gaussian (i.e., cluster id $\mathbf{z}_n = k$)

$$p(\mathbf{z}_n = k | \pi) = \pi_k$$

- Same as a **multinoulli prior** on \mathbf{z} , i.e., $p(\mathbf{z}_n | \pi) = \prod_{k=1}^K \pi_k^{z_{nk}}$

Gaussian Mixture Model (GMM)

- A generative model for data clustering. Assume data generated from a mixture of K Gaussians




- Let $\pi = (\pi_1, \pi_2, \dots, \pi_K)$ with $\sum_{k=1}^K \pi_k = 1$ denote the “mixing weights” of the K Gaussians
- Intuitively $\pi_k \in (0, 1)$ is the fraction of data “contributed” by the k -th Gaussian
- π_k is the prior probability that \mathbf{x}_n comes from the k -th Gaussian (i.e., cluster id $\mathbf{z}_n = k$)

$$p(\mathbf{z}_n = k | \pi) = \pi_k$$

- Same as a **multinoulli prior** on \mathbf{z} , i.e., $p(\mathbf{z}_n | \pi) = \prod_{k=1}^K \pi_k^{z_{nk}}$ (note $z_{nk} = 1$ if $\mathbf{z}_n = k$; 0 otherwise)

Gaussian Mixture Model (GMM)


- A generative model for data clustering. Assume data generated from a mixture of K Gaussians



- Let $\pi = (\pi_1, \pi_2, \dots, \pi_K)$ with $\sum_{k=1}^K \pi_k = 1$ denote the “mixing weights” of the K Gaussians
- Intuitively $\pi_k \in (0, 1)$ is the fraction of data “contributed” by the k -th Gaussian
- π_k is the prior probability that \mathbf{x}_n comes from the k -th Gaussian (i.e., cluster id $\mathbf{z}_n = k$)
$$p(\mathbf{z}_n = k | \pi) = \pi_k$$
- Same as a **multinoulli prior** on \mathbf{z} , i.e., $p(\mathbf{z}_n | \pi) = \prod_{k=1}^K \pi_k^{z_{nk}}$ (note $z_{nk} = 1$ if $\mathbf{z}_n = k$; 0 otherwise)
- Given $\mathbf{z}_n = k$, we generate \mathbf{x}_n from the k -th Gaussian.

Gaussian Mixture Model (GMM)

- A generative model for data clustering. Assume data generated from a mixture of K Gaussians



- Let $\pi = (\pi_1, \pi_2, \dots, \pi_K)$ with $\sum_{k=1}^K \pi_k = 1$ denote the “mixing weights” of the K Gaussians
- Intuitively $\pi_k \in (0, 1)$ is the fraction of data “contributed” by the k -th Gaussian
- π_k is the prior probability that \mathbf{x}_n comes from the k -th Gaussian (i.e., cluster id $\mathbf{z}_n = k$)
$$p(\mathbf{z}_n = k | \pi) = \pi_k$$
- Same as a **multinoulli prior** on \mathbf{z} , i.e., $p(\mathbf{z}_n | \pi) = \prod_{k=1}^K \pi_k^{z_{nk}}$ (note $z_{nk} = 1$ if $\mathbf{z}_n = k$; 0 otherwise)
- Given $\mathbf{z}_n = k$, we generate \mathbf{x}_n from the k -th Gaussian. Thus

$$p(\mathbf{x}_n | \mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n | \mu_k, \Sigma_k)$$

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$
- What is the marginal distribution of \mathbf{x}_n , i.e., $p(\mathbf{x}_n)$?

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$
- What is the marginal distribution of \mathbf{x}_n , i.e., $p(\mathbf{x}_n)$?
 - To get that, we must sum over all possibilities of \mathbf{z}_n

$$p(\mathbf{x}_n) = \sum_{k=1}^K p(\mathbf{x}_n, \mathbf{z}_n = k)$$

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$
- What is the marginal distribution of \mathbf{x}_n , i.e., $p(\mathbf{x}_n)$?
 - To get that, we must sum over all possibilities of \mathbf{z}_n

$$p(\mathbf{x}_n) = \sum_{k=1}^K p(\mathbf{x}_n, \mathbf{z}_n = k) = \sum_{k=1}^K p(\mathbf{z}_n = k)p(\mathbf{x}_n|\mathbf{z}_n = k)$$

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$
- What is the marginal distribution of \mathbf{x}_n , i.e., $p(\mathbf{x}_n)$?

- To get that, we must sum over all possibilities of \mathbf{z}_n

$$p(\mathbf{x}_n) = \sum_{k=1}^K p(\mathbf{x}_n, \mathbf{z}_n = k) = \sum_{k=1}^K p(\mathbf{z}_n = k)p(\mathbf{x}_n|\mathbf{z}_n = k)$$

- Therefore the marginal distribution of \mathbf{x}_n will be

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$
- What is the marginal distribution of \mathbf{x}_n , i.e., $p(\mathbf{x}_n)$?
 - To get that, we must sum over all possibilities of \mathbf{z}_n

$$p(\mathbf{x}_n) = \sum_{k=1}^K p(\mathbf{x}_n, \mathbf{z}_n = k) = \sum_{k=1}^K p(\mathbf{z}_n = k)p(\mathbf{x}_n|\mathbf{z}_n = k)$$

- Therefore the marginal distribution of \mathbf{x}_n will be

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$
- What is the marginal distribution of \mathbf{x}_n , i.e., $p(\mathbf{x}_n)$?

- To get that, we must sum over all possibilities of \mathbf{z}_n

$$p(\mathbf{x}_n) = \sum_{k=1}^K p(\mathbf{x}_n, \mathbf{z}_n = k) = \sum_{k=1}^K p(\mathbf{z}_n = k)p(\mathbf{x}_n|\mathbf{z}_n = k)$$

- Therefore the marginal distribution of \mathbf{x}_n will be

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where Θ collectively denotes all the parameters of the GMM model $(\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$
- What is the marginal distribution of \mathbf{x}_n , i.e., $p(\mathbf{x}_n)$?

- To get that, we must sum over all possibilities of \mathbf{z}_n

$$p(\mathbf{x}_n) = \sum_{k=1}^K p(\mathbf{x}_n, \mathbf{z}_n = k) = \sum_{k=1}^K p(\mathbf{z}_n = k)p(\mathbf{x}_n|\mathbf{z}_n = k)$$

- Therefore the marginal distribution of \mathbf{x}_n will be

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where Θ collectively denotes all the parameters of the GMM model $(\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- Note that GMM also defines probability density (mixture of Gaussians) for the inputs. Therefore it can be used as both a **density estimation** model as well as a **clustering** model

Gaussian Mixture Model (GMM)

- Recall that $p(\mathbf{z}_n = k|\pi) = \pi_k$ and the conditional distribution $p(\mathbf{x}_n|\mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$
- What is the marginal distribution of \mathbf{x}_n , i.e., $p(\mathbf{x}_n)$?

- To get that, we must sum over all possibilities of \mathbf{z}_n

$$p(\mathbf{x}_n) = \sum_{k=1}^K p(\mathbf{x}_n, \mathbf{z}_n = k) = \sum_{k=1}^K p(\mathbf{z}_n = k)p(\mathbf{x}_n|\mathbf{z}_n = k)$$

- Therefore the marginal distribution of \mathbf{x}_n will be

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where Θ collectively denotes all the parameters of the GMM model $(\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- Note that GMM also defines probability density (mixture of Gaussians) for the inputs. Therefore it can be used as both a **density estimation** model as well as a **clustering** model
- Goal: Learn the GMM parameters $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$ (and cluster assignments $\{\mathbf{z}_1, \dots, \mathbf{z}_N\}$)

Gaussian Mixture Model (GMM)

- Recall the marginal distribution of \mathbf{x}_n

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

Gaussian Mixture Model (GMM)

- Recall the marginal distribution of \mathbf{x}_n

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

Gaussian Mixture Model (GMM)

- Recall the marginal distribution of \mathbf{x}_n

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- Suppose we want to do MLE for Θ . The MLE objective will be

$$\sum_{n=1}^N \log p(\mathbf{x}_n|\Theta) = \sum_{n=1}^N \log \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

- Doing MLE on this objective is tricky due to the “ $\log \sum$ ” term

Gaussian Mixture Model (GMM)

- Recall the marginal distribution of \mathbf{x}_n

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- Suppose we want to do MLE for Θ . The MLE objective will be

$$\sum_{n=1}^N \log p(\mathbf{x}_n|\Theta) = \sum_{n=1}^N \log \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

- Doing MLE on this objective is tricky due to the “ $\log \sum$ ” term
 - Parameters get coupled; no closed form solution (iterative methods needed, slow convergence)

Gaussian Mixture Model (GMM)

- Recall the marginal distribution of \mathbf{x}_n

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- Suppose we want to do MLE for Θ . The MLE objective will be

$$\sum_{n=1}^N \log p(\mathbf{x}_n|\Theta) = \sum_{n=1}^N \log \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

- Doing MLE on this objective is tricky due to the “ $\log \sum$ ” term
 - Parameters get coupled; no closed form solution (iterative methods needed, slow convergence)
 - Expectation Maximization (EM) helps solve such problems in a clean and efficient way

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**
- Key idea: Once \mathbf{z} is known, the MLE/MAP becomes very simple

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**
- Key idea: Once \mathbf{z} is known, the MLE/MAP becomes very simple
 - E.g., for GMM, once we know the “labels” \mathbf{z} , it reduces to estimating K Gaussians independently

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**
- Key idea: Once \mathbf{z} is known, the MLE/MAP becomes very simple
 - E.g., for GMM, once we know the “labels” \mathbf{z} , it reduces to estimating K Gaussians independently
- EM alternates between the following two steps until convergence

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**
- Key idea: Once \mathbf{z} is known, the MLE/MAP becomes very simple
 - E.g., for GMM, once we know the “labels” \mathbf{z} , it reduces to estimating K Gaussians independently
- EM alternates between the following two steps until convergence
 - Given the current estimate of Θ , **make a good “guess” for \mathbf{z} for each \mathbf{x}**

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**
- Key idea: Once \mathbf{z} is known, the MLE/MAP becomes very simple
 - E.g., for GMM, once we know the “labels” \mathbf{z} , it reduces to estimating K Gaussians independently
- EM alternates between the following two steps until convergence
 - Given the current estimate of Θ , **make a good “guess” for \mathbf{z}** for each \mathbf{x}
 - Use these guesses of \mathbf{z} to do **MLE/MAP on $p(\mathbf{x}, \mathbf{z}|\Theta)$** to “refine” Θ

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**
- Key idea: Once \mathbf{z} is known, the MLE/MAP becomes very simple
 - E.g., for GMM, once we know the “labels” \mathbf{z} , it reduces to estimating K Gaussians independently
- EM alternates between the following two steps until convergence
 - Given the current estimate of Θ , **make a good “guess” for \mathbf{z}** for each \mathbf{x}
 - Use these guesses of \mathbf{z} to do **MLE/MAP on $p(\mathbf{x}, \mathbf{z}|\Theta)$** to “refine” Θ
- **Important:** EM leads to very easy param. updates if $p(\mathbf{x}|\mathbf{z})$ and $p(\mathbf{z})$ are **exp. family distributions**

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**
- Key idea: Once \mathbf{z} is known, the MLE/MAP becomes very simple
 - E.g., for GMM, once we know the “labels” \mathbf{z} , it reduces to estimating K Gaussians independently
- EM alternates between the following two steps until convergence
 - Given the current estimate of Θ , **make a good “guess” for \mathbf{z}** for each \mathbf{x}
 - Use these guesses of \mathbf{z} to do **MLE/MAP on $p(\mathbf{x}, \mathbf{z}|\Theta)$** to “refine” Θ
- **Important:** EM leads to very easy param. updates if $p(\mathbf{x}|\mathbf{z})$ and $p(\mathbf{z})$ are **exp. family distributions**
- EM is guaranteed to converge to a local optima

Expectation Maximization (EM)

- A **general algorithm** for doing MLE/MAP in models that contain latent variables (e.g., GMM)
- The basic (rough) idea is to do MLE/MAP **not on $p(\mathbf{x}|\Theta)$** but **on $p(\mathbf{x}, \mathbf{z}|\Theta)$**
- EM thinks of not just \mathbf{x} as the data but **both \mathbf{x} and a good “guess” of \mathbf{z} as the data**
- Key idea: Once \mathbf{z} is known, the MLE/MAP becomes very simple
 - E.g., for GMM, once we know the “labels” \mathbf{z} , it reduces to estimating K Gaussians independently
- EM alternates between the following two steps until convergence
 - Given the current estimate of Θ , **make a good “guess” for \mathbf{z}** for each \mathbf{x}
 - Use these guesses of \mathbf{z} to do **MLE/MAP on $p(\mathbf{x}, \mathbf{z}|\Theta)$** to “refine” Θ
- **Important:** EM leads to very easy param. updates if $p(\mathbf{x}|\mathbf{z})$ and $p(\mathbf{z})$ are **exp. family distributions**
- EM is guaranteed to converge to a local optima
- More details in the next class..