

# Approximate Inference: Sampling Methods (2)


Piyush Rai

Probabilistic Machine Learning (CS772A)

Oct 3, 2017


# Sampling Methods: Recap

- Any probability distribution  $p(\mathbf{z})$  can be (approximately) represented using a set of samples



# Sampling Methods: Recap


- Any probability distribution  $p(\mathbf{z})$  can be (approximately) represented using a set of samples



- Samples can come from  $p(\mathbf{z})$  or some “proposal distribution” if  $p(\mathbf{z})$  is a “difficult” distribution

# Sampling Methods: Recap

- Any probability distribution  $p(\mathbf{z})$  can be (approximately) represented using a set of samples



- Samples can come from  $p(\mathbf{z})$  or some “proposal distribution” if  $p(\mathbf{z})$  is a “difficult” distribution
- Given a set of samples  $\{\mathbf{z}^{(\ell)}\}_{\ell=1}^L$ , the sample-based approximation of  $p(\mathbf{z})$  can be written as

$$p(\mathbf{z}) \approx \frac{1}{L} \sum_{\ell=1}^L \delta(\mathbf{z} = \mathbf{z}^{(\ell)}) \quad \text{or} \quad p(\mathbf{z}) \approx \frac{1}{L} \sum_{\ell=1}^L \delta_{\mathbf{z}^{(\ell)}}(\mathbf{z})$$

# Sampling Methods: Recap

- Looked at some basic methods for **generating samples** from a probability distribution
- Also, using the samples to **approximate difficult to compute expectations**

# Sampling Methods: Recap

- Looked at some basic methods for **generating samples** from a probability distribution
  - Transformation based methods
  - Rejection sampling
- Also, using the samples to **approximate difficult to compute expectations**

# Sampling Methods: Recap

- Looked at some basic methods for **generating samples** from a probability distribution
  - Transformation based methods
  - Rejection sampling
- Also, using the samples to **approximate difficult to compute expectations**

Monte Carlo Sampling:  $\mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)})$  where  $\{z^{(\ell)}\}_{\ell=1}^L \sim p(z)$

# Sampling Methods: Recap

- Looked at some basic methods for **generating samples** from a probability distribution
  - Transformation based methods
  - Rejection sampling
- Also, using the samples to **approximate difficult to compute expectations**

$$\text{Monte Carlo Sampling: } \mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)}) \quad \text{where } \{z^{(\ell)}\}_{\ell=1}^L \sim p(z)$$

$$\text{Important Sampling (1): } \mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)}) \frac{p(z^{(\ell)})}{q(z^{(\ell)})} \quad \text{where } \{z^{(\ell)}\}_{\ell=1}^L \sim q(z)$$



# Sampling Methods: Recap

- Looked at some basic methods for **generating samples** from a probability distribution
  - Transformation based methods
  - Rejection sampling
- Also, using the samples to **approximate difficult to compute expectations**

$$\text{Monte Carlo Sampling: } \mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)}) \quad \text{where } \{z^{(\ell)}\}_{\ell=1}^L \sim p(z)$$

$$\text{Important Sampling (1): } \mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)}) \frac{p(z^{(\ell)})}{q(z^{(\ell)})} \quad \text{where } \{z^{(\ell)}\}_{\ell=1}^L \sim q(z)$$

$$\text{Important Sampling (2): } \mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{Z_q}{Z_p} \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)}) \frac{\tilde{p}(z^{(\ell)})}{\tilde{q}(z^{(\ell)})} \quad \text{where } \{z^{(\ell)}\}_{\ell=1}^L \sim q(z)$$

# Sampling Methods: Recap

- Looked at some basic methods for **generating samples** from a probability distribution
  - Transformation based methods
  - Rejection sampling
- Also, using the samples to **approximate difficult to compute expectations**

$$\text{Monte Carlo Sampling: } \mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)}) \quad \text{where } \{z^{(\ell)}\}_{\ell=1}^L \sim p(z)$$

$$\text{Important Sampling (1): } \mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)}) \frac{p(z^{(\ell)})}{q(z^{(\ell)})} \quad \text{where } \{z^{(\ell)}\}_{\ell=1}^L \sim q(z)$$

$$\text{Important Sampling (2): } \mathbb{E}_{p(z)}[f(z)] = \int f(z)p(z)dz \approx \frac{Z_q}{Z_p} \frac{1}{L} \sum_{\ell=1}^L f(z^{(\ell)}) \frac{\tilde{p}(z^{(\ell)})}{\tilde{q}(z^{(\ell)})} \quad \text{where } \{z^{(\ell)}\}_{\ell=1}^L \sim q(z)$$

[Note: I.S. (1) assumes  $p(z)$  can be evaluated at any  $z$ , I.S. (2) assumes  $p(z) = \frac{\tilde{p}(z)}{Z_p}$  can only be evaluated up to a prop. constant]



# Limitations of Basic Sampling Methods

# Limitations of Basic Sampling Methods

- Transformation based methods: Usually limited to drawing from standard distributions



# Limitations of Basic Sampling Methods

- Transformation based methods: Usually limited to drawing from standard distributions
- Rejection Sampling and Importance Sampling: Require good proposal distributions



# Limitations of Basic Sampling Methods



- Transformation based methods: Usually limited to drawing from standard distributions
- Rejection Sampling and Importance Sampling: Require good proposal distributions



- Difficult to find good prop. distr. especially when  $\mathbf{z}$  is high-dim. (e.g., models with many params)

# Limitations of Basic Sampling Methods



- Transformation based methods: Usually limited to drawing from standard distributions
- Rejection Sampling and Importance Sampling: Require good proposal distributions



- Difficult to find good prop. distr. especially when  $\mathbf{z}$  is high-dim. (e.g., models with many params)
  - In high dimensions, most of the mass of  $p(\mathbf{z})$  is concentrated in a tiny region of the  $\mathbf{z}$  space

# Limitations of Basic Sampling Methods

- Transformation based methods: Usually limited to drawing from standard distributions
- Rejection Sampling and Importance Sampling: Require good proposal distributions



- Difficult to find good prop. distr. especially when  $\mathbf{z}$  is high-dim. (e.g., models with many params)
  - In high dimensions, most of the mass of  $p(\mathbf{z})$  is concentrated in a tiny region of the  $\mathbf{z}$  space
  - Difficult to *a priori* know what those regions are, thus difficult to come up with good proposal dist.



# Markov Chain Monte Carlo (MCMC) Methods

# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional

# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional
- Will again assume that we can evaluate  $p(\mathbf{z})$  at least up to a proportionality constant

# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional
- Will again assume that we can evaluate  $p(\mathbf{z})$  at least up to a proportionality constant
- Basic idea in MCMC: Use a **Markov Chain** to generate samples from  $p(\mathbf{z})$

$$\mathbf{z}^{(1)} \rightarrow \mathbf{z}^{(2)} \rightarrow \dots \rightarrow \mathbf{z}^{(L)}$$

# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional
- Will again assume that we can evaluate  $p(\mathbf{z})$  at least up to a proportionality constant
- Basic idea in MCMC: Use a **Markov Chain** to generate samples from  $p(\mathbf{z})$

$$\mathbf{z}^{(1)} \rightarrow \mathbf{z}^{(2)} \rightarrow \dots \rightarrow \mathbf{z}^{(L)}$$

- How: Given a current sample  $\mathbf{z}^{(\ell)}$  from the chain, generate the next sample  $\mathbf{z}^{(\ell+1)}$  as

# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional
- Will again assume that we can evaluate  $p(\mathbf{z})$  at least up to a proportionality constant
- Basic idea in MCMC: Use a **Markov Chain** to generate samples from  $p(\mathbf{z})$

$$\mathbf{z}^{(1)} \rightarrow \mathbf{z}^{(2)} \rightarrow \dots \rightarrow \mathbf{z}^{(L)}$$

- How: Given a current sample  $\mathbf{z}^{(\ell)}$  from the chain, generate the next sample  $\mathbf{z}^{(\ell+1)}$  as
  - Use a **proposal distribution**  $q(\mathbf{z}|\mathbf{z}^{(\ell)})$  to generate a candidate sample  $\mathbf{z}^*$

# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional
- Will again assume that we can evaluate  $p(\mathbf{z})$  at least up to a proportionality constant
- Basic idea in MCMC: Use a **Markov Chain** to generate samples from  $p(\mathbf{z})$

$$\mathbf{z}^{(1)} \rightarrow \mathbf{z}^{(2)} \rightarrow \dots \rightarrow \mathbf{z}^{(L)}$$

- How: Given a current sample  $\mathbf{z}^{(\ell)}$  from the chain, generate the next sample  $\mathbf{z}^{(\ell+1)}$  as
  - Use a **proposal distribution**  $q(\mathbf{z}|\mathbf{z}^{(\ell)})$  to generate a candidate sample  $\mathbf{z}^*$
  - Accept/reject  $\mathbf{z}^*$  as the next sample based on an acceptance criterion (will see later)

# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional
- Will again assume that we can evaluate  $p(\mathbf{z})$  at least up to a proportionality constant
- Basic idea in MCMC: Use a **Markov Chain** to generate samples from  $p(\mathbf{z})$

$$\mathbf{z}^{(1)} \rightarrow \mathbf{z}^{(2)} \rightarrow \dots \rightarrow \mathbf{z}^{(L)}$$

- How: Given a current sample  $\mathbf{z}^{(\ell)}$  from the chain, generate the next sample  $\mathbf{z}^{(\ell+1)}$  as
  - Use a **proposal distribution**  $q(\mathbf{z}|\mathbf{z}^{(\ell)})$  to generate a candidate sample  $\mathbf{z}^*$
  - Accept/reject  $\mathbf{z}^*$  as the next sample based on an acceptance criterion (will see later)
  - If accepted,  $\mathbf{z}^{(\ell+1)} = \mathbf{z}^*$ . If rejected,  $\mathbf{z}^{(\ell+1)} = \mathbf{z}^{(\ell)}$



# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional
- Will again assume that we can evaluate  $p(\mathbf{z})$  at least up to a proportionality constant
- Basic idea in MCMC: Use a **Markov Chain** to generate samples from  $p(\mathbf{z})$

$$\mathbf{z}^{(1)} \rightarrow \mathbf{z}^{(2)} \rightarrow \dots \rightarrow \mathbf{z}^{(L)}$$

- How: Given a current sample  $\mathbf{z}^{(\ell)}$  from the chain, generate the next sample  $\mathbf{z}^{(\ell+1)}$  as
  - Use a **proposal distribution**  $q(\mathbf{z}|\mathbf{z}^{(\ell)})$  to generate a candidate sample  $\mathbf{z}^*$
  - Accept/reject  $\mathbf{z}^*$  as the next sample based on an acceptance criterion (will see later)
  - If accepted,  $\mathbf{z}^{(\ell+1)} = \mathbf{z}^*$ . If rejected,  $\mathbf{z}^{(\ell+1)} = \mathbf{z}^{(\ell)}$
- If  $q(\mathbf{z}|\mathbf{z}^{(\ell)})$  has certain properties, the Markov chain's **stationary distribution** will be  $p(\mathbf{z})$


# Markov Chain Monte Carlo (MCMC)

- Goal: Generate samples from some **target distribution**  $p(\mathbf{z}) = \frac{\tilde{p}(\mathbf{z})}{Z}$ , where  $\mathbf{z}$  is high-dimensional
- Will again assume that we can evaluate  $p(\mathbf{z})$  at least up to a proportionality constant
- Basic idea in MCMC: Use a **Markov Chain** to generate samples from  $p(\mathbf{z})$

$$\mathbf{z}^{(1)} \rightarrow \mathbf{z}^{(2)} \rightarrow \dots \rightarrow \mathbf{z}^{(L)}$$

- How: Given a current sample  $\mathbf{z}^{(\ell)}$  from the chain, generate the next sample  $\mathbf{z}^{(\ell+1)}$  as
  - Use a **proposal distribution**  $q(\mathbf{z}|\mathbf{z}^{(\ell)})$  to generate a candidate sample  $\mathbf{z}^*$
  - Accept/reject  $\mathbf{z}^*$  as the next sample based on an acceptance criterion (will see later)
  - If accepted,  $\mathbf{z}^{(\ell+1)} = \mathbf{z}^*$ . If rejected,  $\mathbf{z}^{(\ell+1)} = \mathbf{z}^{(\ell)}$
- If  $q(\mathbf{z}|\mathbf{z}^{(\ell)})$  has certain properties, the Markov chain's **stationary distribution** will be  $p(\mathbf{z})$ 
  - Informally, stationary distribution means where the chain will eventually “reach”

# Markov Chain Monte Carlo (MCMC)



# Markov Chain

- Consider a sequence of random variables  $\mathbf{z}^{(1)}, \dots, \mathbf{z}^{(L)}$

# Markov Chain

- Consider a sequence of random variables  $\mathbf{z}^{(1)}, \dots, \mathbf{z}^{(L)}$
- A **first-order Markov Chain** assumes

$$p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(\ell)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)}) \quad \forall \ell$$

# Markov Chain

- Consider a sequence of random variables  $\mathbf{z}^{(1)}, \dots, \mathbf{z}^{(L)}$
- A **first-order Markov Chain** assumes

$$p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(\ell)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)}) \quad \forall \ell$$

- A first order Markov chain can be defined by the following

# Markov Chain

- Consider a sequence of random variables  $\mathbf{z}^{(1)}, \dots, \mathbf{z}^{(L)}$
- A **first-order Markov Chain** assumes

$$p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(\ell)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)}) \quad \forall \ell$$

- A first order Markov chain can be defined by the following
  - An **initial state distribution**  $p(\mathbf{z}^{(0)})$

# Markov Chain

- Consider a sequence of random variables  $\mathbf{z}^{(1)}, \dots, \mathbf{z}^{(L)}$
- A **first-order Markov Chain** assumes

$$p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(\ell)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)}) \quad \forall \ell$$

- A first order Markov chain can be defined by the following
  - An **initial state distribution**  $p(\mathbf{z}^{(0)})$
  - **Transition probabilities**  $T_\ell(\mathbf{z}^{(\ell)}, \mathbf{z}^{(\ell+1)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)})$



# Markov Chain

- Consider a sequence of random variables  $\mathbf{z}^{(1)}, \dots, \mathbf{z}^{(L)}$
- A **first-order Markov Chain** assumes

$$p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(\ell)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)}) \quad \forall \ell$$

- A first order Markov chain can be defined by the following
  - An **initial state distribution**  $p(\mathbf{z}^{(0)})$
  - **Transition probabilities**  $T_\ell(\mathbf{z}^{(\ell)}, \mathbf{z}^{(\ell+1)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)})$ : Distribution over the possible values of  $\mathbf{z}^{(\ell+1)}$

# Markov Chain


- Consider a sequence of random variables  $\mathbf{z}^{(1)}, \dots, \mathbf{z}^{(L)}$
- A **first-order Markov Chain** assumes

$$p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(\ell)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)}) \quad \forall \ell$$

- A first order Markov chain can be defined by the following
  - An **initial state distribution**  $p(\mathbf{z}^{(0)})$
  - Transition probabilities**  $T_\ell(\mathbf{z}^{(\ell)}, \mathbf{z}^{(\ell+1)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)})$ : Distribution over the possible values of  $\mathbf{z}^{(\ell+1)}$

Transition probabilities  
can be defined using a  
 $K \times K$  table if  $\mathbf{z}$  is a discrete  
r.v. with  $K$  possible values

	1	2	3	4	5
1	0.4	0.6	0.0	0.0	0.0
2	0.5	0.0	0.5	0.0	0.0
3	0.0	0.3	0.0	0.7	0.0
4	0.0	0.0	0.1	0.3	0.6
5	0.0	0.3	0.0	0.5	0.2



# Markov Chain


- Consider a sequence of random variables  $\mathbf{z}^{(1)}, \dots, \mathbf{z}^{(L)}$
- A **first-order Markov Chain** assumes

$$p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(\ell)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)}) \quad \forall \ell$$

- A first order Markov chain can be defined by the following
  - An **initial state distribution**  $p(\mathbf{z}^{(0)})$
  - Transition probabilities**  $T_\ell(\mathbf{z}^{(\ell)}, \mathbf{z}^{(\ell+1)}) = p(\mathbf{z}^{(\ell+1)} | \mathbf{z}^{(\ell)})$ : Distribution over the possible values of  $\mathbf{z}^{(\ell+1)}$

Transition probabilities  
can be defined using a  
 $K \times K$  table if  $\mathbf{z}$  is a discrete  
r.v. with  $K$  possible values

	1	2	3	4	5
1	0.4	0.6	0.0	0.0	0.0
2	0.5	0.0	0.5	0.0	0.0
3	0.0	0.3	0.0	0.7	0.0
4	0.0	0.0	0.1	0.3	0.6
5	0.0	0.3	0.0	0.5	0.2



- Homogeneous Markov Chain:** Transition probabilities  $T_\ell = T$  (same everywhere along the chain)

# Some Properties

- Consider the discrete case when  $\mathbf{z}$  has  $K$  possible states (and  $T$  is a matrix of size  $K \times K$ )

# Some Properties

- Consider the discrete case when  $\mathbf{z}$  has  $K$  possible states (and  $T$  is a matrix of size  $K \times K$ )
- Assume the graph representing the possible state transitions is **irreducible** and **aperiodic**

# Some Properties

- Consider the discrete case when  $\mathbf{z}$  has  $K$  possible states (and  $T$  is a matrix of size  $K \times K$ )
- Assume the graph representing the possible state transitions is **irreducible** and **aperiodic**
- **Ergodic Property:** Under the above assumption, for any choice of an **initial probability vector**  $\mathbf{v}$

$$\mathbf{v}T^m = \mathbf{p}^* \quad \text{as } m \rightarrow \infty$$

where the probability vector  $\mathbf{p}^*$  represents the **invariant or stationary distribution** of the chain

# Some Properties

- Consider the discrete case when  $\mathbf{z}$  has  $K$  possible states (and  $T$  is a matrix of size  $K \times K$ )
- Assume the graph representing the possible state transitions is **irreducible** and **aperiodic**
- **Ergodic Property:** Under the above assumption, for any choice of an **initial probability vector**  $\mathbf{v}$

$$\mathbf{v}T^m = \mathbf{p}^* \quad \text{as } m \rightarrow \infty$$

where the probability vector  $\mathbf{p}^*$  represents the **invariant or stationary distribution** of the chain

- Why do we need the graph to be irreducible and aperiodic?

# Some Properties

- Consider the discrete case when  $\mathbf{z}$  has  $K$  possible states (and  $T$  is a matrix of size  $K \times K$ )
- Assume the graph representing the possible state transitions is **irreducible** and **aperiodic**
- **Ergodic Property:** Under the above assumption, for any choice of an **initial probability vector**  $\mathbf{v}$

$$\mathbf{v}T^m = \mathbf{p}^* \quad \text{as } m \rightarrow \infty$$

where the probability vector  $\mathbf{p}^*$  represents the **invariant or stationary distribution** of the chain

- Why do we need the graph to be irreducible and aperiodic?
  - Irreducible: No disjoint sets of nodes. Can reach from any state to any state



# Some Properties

- Consider the discrete case when  $\mathbf{z}$  has  $K$  possible states (and  $T$  is a matrix of size  $K \times K$ )
- Assume the graph representing the possible state transitions is **irreducible** and **aperiodic**
- **Ergodic Property:** Under the above assumption, for any choice of an **initial probability vector**  $\mathbf{v}$

$$\mathbf{v}T^m = \mathbf{p}^* \quad \text{as } m \rightarrow \infty$$

where the probability vector  $\mathbf{p}^*$  represents the **invariant or stationary distribution** of the chain

- Why do we need the graph to be irreducible and aperiodic?
  - Irreducible: No disjoint sets of nodes. Can reach from any state to any state
  - Aperiodic: No cycles in the graph (otherwise would oscillate forever). Consider this example

$$\mathbf{v} = [1/5, 4/5] \quad T = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

.. multiplying  $\mathbf{v}$  by  $T$  repeatedly leads to oscillating values without convergence

# Some Properties

- Note that, for Ergodic case,  $\boldsymbol{p}^* T = \boldsymbol{p}^*$

# Some Properties

- Note that, for Ergodic case,  $\boldsymbol{p}^* T = \boldsymbol{p}^*$ . Therefore  $\boldsymbol{\pi}^*$  is the left eigenvector of  $T$  with eigenvalue 1

# Some Properties

- Note that, for Ergodic case,  $\boldsymbol{p}^* T = \boldsymbol{p}^*$ . Therefore  $\boldsymbol{\pi}^*$  is the left eigenvector of  $T$  with eigenvalue 1
- For the discrete-valued  $\boldsymbol{z}$  with  $K = 5$  possible states,  $\boldsymbol{p}^* = [p_1^*, \dots, p_5^*]$ , and we can write

$$\sum_{i=1}^5 p_i^* T_{ij} = p_j^*$$

# Some Properties

- Note that, for Ergodic case,  $\mathbf{p}^* T = \mathbf{p}^*$ . Therefore  $\pi^*$  is the left eigenvector of  $T$  with eigenvalue 1
- For the discrete-valued  $\mathbf{z}$  with  $K = 5$  possible states,  $\mathbf{p}^* = [p_1^*, \dots, p_5^*]$ , and we can write

$$\sum_{i=1}^5 p_i^* T_{ij} = p_j^*$$

- For the continuous  $\mathbf{z}$  case, we can equivalently write (for any two state values  $\mathbf{z}$  and  $\mathbf{z}'$ )

$$\int p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) d\mathbf{z}' = p^*(\mathbf{z})$$

# Some Properties

- Note that, for Ergodic case,  $\mathbf{p}^* T = \mathbf{p}^*$ . Therefore  $\pi^*$  is the left eigenvector of  $T$  with eigenvalue 1
- For the discrete-valued  $\mathbf{z}$  with  $K = 5$  possible states,  $\mathbf{p}^* = [p_1^*, \dots, p_5^*]$ , and we can write

$$\sum_{i=1}^5 p_i^* T_{ij} = p_j^*$$

- For the continuous  $\mathbf{z}$  case, we can equivalently write (for any two state values  $\mathbf{z}$  and  $\mathbf{z}'$ )

$$\int p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) d\mathbf{z}' = p^*(\mathbf{z})$$

- Suppose a Markov chain with transition probabilities  $T$  satisfies

$$p^*(\mathbf{z}) T(\mathbf{z}, \mathbf{z}') = p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) \quad (\text{Detailed Balance})$$

# Some Properties

- Note that, for Ergodic case,  $\mathbf{p}^* T = \mathbf{p}^*$ . Therefore  $\pi^*$  is the left eigenvector of  $T$  with eigenvalue 1
- For the discrete-valued  $\mathbf{z}$  with  $K = 5$  possible states,  $\mathbf{p}^* = [p_1^*, \dots, p_5^*]$ , and we can write

$$\sum_{i=1}^5 p_i^* T_{ij} = p_j^*$$

- For the continuous  $\mathbf{z}$  case, we can equivalently write (for any two state values  $\mathbf{z}$  and  $\mathbf{z}'$ )

$$\int p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) d\mathbf{z}' = p^*(\mathbf{z})$$

- Suppose a Markov chain with transition probabilities  $T$  satisfies

$$p^*(\mathbf{z}) T(\mathbf{z}, \mathbf{z}') = p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) \quad (\text{Detailed Balance})$$

- Integrating both sides w.r.t.  $\mathbf{z}'$  gives  $\int p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) d\mathbf{z}' = p^*(\mathbf{z})$  (i.e., the Ergodic property)
  - Thus a Markov chain with detailed balance will always converge to a stationary distribution  $p^*(\mathbf{z})$

# Some Properties

- Note that, for Ergodic case,  $\mathbf{p}^* T = \mathbf{p}^*$ . Therefore  $\pi^*$  is the left eigenvector of  $T$  with eigenvalue 1
- For the discrete-valued  $\mathbf{z}$  with  $K = 5$  possible states,  $\mathbf{p}^* = [p_1^*, \dots, p_5^*]$ , and we can write

$$\sum_{i=1}^5 p_i^* T_{ij} = p_j^*$$

- For the continuous  $\mathbf{z}$  case, we can equivalently write (for any two state values  $\mathbf{z}$  and  $\mathbf{z}'$ )

$$\int p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) d\mathbf{z}' = p^*(\mathbf{z})$$

- Suppose a Markov chain with transition probabilities  $T$  satisfies


$$p^*(\mathbf{z}) T(\mathbf{z}, \mathbf{z}') = p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) \quad (\text{Detailed Balance})$$

- Integrating both sides w.r.t.  $\mathbf{z}'$  gives  $\int p^*(\mathbf{z}') T(\mathbf{z}', \mathbf{z}) d\mathbf{z}' = p^*(\mathbf{z})$  (i.e., the Ergodic property)
  - Thus a Markov chain with detailed balance will always converge to a stationary distribution  $p^*(\mathbf{z})$
  - Homogeneous Markov Chains satisfy detailed balance/ergodic property under mild conditions




# MCMC: The Basic Scheme

- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



# MCMC: The Basic Scheme


- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



- But we usually require several samples to approximate the distribution. How do we get those?

# MCMC: The Basic Scheme


- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



- But we usually require several samples to approximate the distribution. How do we get those?
  - Start at an initial  $\mathbf{z}^{(0)}$ . Using a prop. dist.  $p(\mathbf{z}^{(\ell+1)}|\mathbf{z}^{(\ell)})$ , run the chain long enough, say  $T_1$  steps

# MCMC: The Basic Scheme


- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



- But we usually require several samples to approximate the distribution. How do we get those?
  - Start at an initial  $\mathbf{z}^{(0)}$ . Using a prop. dist.  $p(\mathbf{z}^{(\ell+1)}|\mathbf{z}^{(\ell)})$ , run the chain long enough, say  $T_1$  steps
  - Discard the first  $(T_1 - 1)$  samples (called “burn-in” samples) and take the last sample  $\mathbf{z}^{(T_1)}$

# MCMC: The Basic Scheme


- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



- But we usually require several samples to approximate the distribution. How do we get those?
  - Start at an initial  $\mathbf{z}^{(0)}$ . Using a prop. dist.  $p(\mathbf{z}^{(\ell+1)}|\mathbf{z}^{(\ell)})$ , run the chain long enough, say  $T_1$  steps
  - Discard the first  $(T_1 - 1)$  samples (called “burn-in” samples) and take the last sample  $\mathbf{z}^{(T_1)}$
  - Continue from  $\mathbf{z}^{(T_1)}$  up to  $T_2$  steps, discard intermediate samples, take the last sample  $\mathbf{z}^{(T_2)}$

# MCMC: The Basic Scheme


- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



- But we usually require several samples to approximate the distribution. How do we get those?
  - Start at an initial  $\mathbf{z}^{(0)}$ . Using a prop. dist.  $p(\mathbf{z}^{(\ell+1)}|\mathbf{z}^{(\ell)})$ , run the chain long enough, say  $T_1$  steps
  - Discard the first  $(T_1 - 1)$  samples (called “**burn-in**” samples) and take the last sample  $\mathbf{z}^{(T_1)}$
  - Continue from  $\mathbf{z}^{(T_1)}$  up to  $T_2$  steps, discard intermediate samples, take the last sample  $\mathbf{z}^{(T_2)}$ 
    - This helps ensure that  $\mathbf{z}^{(T_1)}$  and  $\mathbf{z}^{(T_2)}$  are **uncorrelated**

# MCMC: The Basic Scheme


- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



- But we usually require several samples to approximate the distribution. How do we get those?
  - Start at an initial  $\mathbf{z}^{(0)}$ . Using a prop. dist.  $p(\mathbf{z}^{(\ell+1)}|\mathbf{z}^{(\ell)})$ , run the chain long enough, say  $T_1$  steps
  - Discard the first  $(T_1 - 1)$  samples (called “burn-in” samples) and take the last sample  $\mathbf{z}^{(T_1)}$
  - Continue from  $\mathbf{z}^{(T_1)}$  up to  $T_2$  steps, discard intermediate samples, take the last sample  $\mathbf{z}^{(T_2)}$ 
    - This helps ensure that  $\mathbf{z}^{(T_1)}$  and  $\mathbf{z}^{(T_2)}$  are **uncorrelated**
  - Repeat the same for a total of  $S$  times

# MCMC: The Basic Scheme

- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution




- But we usually require several samples to approximate the distribution. How do we get those?
  - Start at an initial  $\mathbf{z}^{(0)}$ . Using a prop. dist.  $p(\mathbf{z}^{(\ell+1)}|\mathbf{z}^{(\ell)})$ , run the chain long enough, say  $T_1$  steps
  - Discard the first  $(T_1 - 1)$  samples (called “burn-in” samples) and take the last sample  $\mathbf{z}^{(T_1)}$
  - Continue from  $\mathbf{z}^{(T_1)}$  up to  $T_2$  steps, discard intermediate samples, take the last sample  $\mathbf{z}^{(T_2)}$ 
    - This helps ensure that  $\mathbf{z}^{(T_1)}$  and  $\mathbf{z}^{(T_2)}$  are **uncorrelated**
  - Repeat the same for a total of  $S$  times
  - In the end, we have  $S$  i.i.d. samples from  $p(\mathbf{z})$ , i.e.,  $\mathbf{z}^{(T_1)}, \mathbf{z}^{(T_2)}, \dots, \mathbf{z}^{(T_S)} \sim p(\mathbf{z})$



# MCMC: The Basic Scheme


- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



- But we usually require several samples to approximate the distribution. How do we get those?
  - Start at an initial  $\mathbf{z}^{(0)}$ . Using a prop. dist.  $p(\mathbf{z}^{(\ell+1)}|\mathbf{z}^{(\ell)})$ , run the chain long enough, say  $T_1$  steps
  - Discard the first  $(T_1 - 1)$  samples (called “burn-in” samples) and take the last sample  $\mathbf{z}^{(T_1)}$
  - Continue from  $\mathbf{z}^{(T_1)}$  up to  $T_2$  steps, discard intermediate samples, take the last sample  $\mathbf{z}^{(T_2)}$ 
    - This helps ensure that  $\mathbf{z}^{(T_1)}$  and  $\mathbf{z}^{(T_2)}$  are **uncorrelated**
  - Repeat the same for a total of  $S$  times
  - In the end, we have  $S$  i.i.d. samples from  $p(\mathbf{z})$ , i.e.,  $\mathbf{z}^{(T_1)}, \mathbf{z}^{(T_2)}, \dots, \mathbf{z}^{(T_S)} \sim p(\mathbf{z})$
  - Note: Good choices for  $T_1$  and  $T_i - T_{i-1}$  are usually based on heuristics

# MCMC: The Basic Scheme

- Running the MCMC chain **infinitely long** gives us ONE sample from the target distribution



- But we usually require several samples to approximate the distribution. How do we get those?
  - Start at an initial  $\mathbf{z}^{(0)}$ . Using a prop. dist.  $p(\mathbf{z}^{(\ell+1)}|\mathbf{z}^{(\ell)})$ , run the chain long enough, say  $T_1$  steps
  - Discard the first  $(T_1 - 1)$  samples (called “burn-in” samples) and take the last sample  $\mathbf{z}^{(T_1)}$
  - Continue from  $\mathbf{z}^{(T_1)}$  up to  $T_2$  steps, discard intermediate samples, take the last sample  $\mathbf{z}^{(T_2)}$ 
    - This helps ensure that  $\mathbf{z}^{(T_1)}$  and  $\mathbf{z}^{(T_2)}$  are **uncorrelated**
  - Repeat the same for a total of  $S$  times
  - In the end, we have  $S$  i.i.d. samples from  $p(\mathbf{z})$ , i.e.,  $\mathbf{z}^{(T_1)}, \mathbf{z}^{(T_2)}, \dots, \mathbf{z}^{(T_S)} \sim p(\mathbf{z})$
  - Note: Good choices for  $T_1$  and  $T_i - T_{i-1}$  are usually based on heuristics
  - Note: MCMC is an **approximate method** because we don't usually know what  $T_1$  is “long enough”

# Some MCMC Sampling Algorithms

# Metropolis-Hastings (MH) Sampling

- Assume a proposal distribution  $q(\mathbf{z}|\mathbf{z}^{(\tau)})$ , e.g.,  $\mathcal{N}(\mathbf{z}|\mathbf{z}^{(\tau)}, \sigma^2 \mathbf{I}_D)$

# Metropolis-Hastings (MH) Sampling

- Assume a proposal distribution  $q(\mathbf{z}|\mathbf{z}^{(\tau)})$ , e.g.,  $\mathcal{N}(\mathbf{z}|\mathbf{z}^{(\tau)}, \sigma^2 \mathbf{I}_D)$
- In each step, draw  $\mathbf{z}^* \sim q(\mathbf{z}|\mathbf{z}^{(\tau)})$  and accept the sample  $\mathbf{z}^*$  with probability

$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})} \right)$$

# Metropolis-Hastings (MH) Sampling

- Assume a proposal distribution  $q(\mathbf{z}|\mathbf{z}^{(\tau)})$ , e.g.,  $\mathcal{N}(\mathbf{z}|\mathbf{z}^{(\tau)}, \sigma^2 \mathbf{I}_D)$
- In each step, draw  $\mathbf{z}^* \sim q(\mathbf{z}|\mathbf{z}^{(\tau)})$  and accept the sample  $\mathbf{z}^*$  with probability

$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})} \right)$$

- The acceptance probability makes intuitive sense. Note the kind of  $\mathbf{z}^*$  would it favor/unfavor:

# Metropolis-Hastings (MH) Sampling

- Assume a proposal distribution  $q(\mathbf{z}|\mathbf{z}^{(\tau)})$ , e.g.,  $\mathcal{N}(\mathbf{z}|\mathbf{z}^{(\tau)}, \sigma^2 \mathbf{I}_D)$
- In each step, draw  $\mathbf{z}^* \sim q(\mathbf{z}|\mathbf{z}^{(\tau)})$  and accept the sample  $\mathbf{z}^*$  with probability

$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})} \right)$$

- The acceptance probability makes intuitive sense. Note the kind of  $\mathbf{z}^*$  would it favor/unfavor:
  - It favors accepting  $\mathbf{z}^*$  if  $\tilde{p}(\mathbf{z}^*)$  has a higher value than  $\tilde{p}(\mathbf{z}^{(\tau)})$

# Metropolis-Hastings (MH) Sampling

- Assume a proposal distribution  $q(\mathbf{z}|\mathbf{z}^{(\tau)})$ , e.g.,  $\mathcal{N}(\mathbf{z}|\mathbf{z}^{(\tau)}, \sigma^2 \mathbf{I}_D)$
- In each step, draw  $\mathbf{z}^* \sim q(\mathbf{z}|\mathbf{z}^{(\tau)})$  and accept the sample  $\mathbf{z}^*$  with probability

$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})} \right)$$

- The acceptance probability makes intuitive sense. Note the kind of  $\mathbf{z}^*$  would it favor/unfavor:
  - It favors accepting  $\mathbf{z}^*$  if  $\tilde{p}(\mathbf{z}^*)$  has a higher value than  $\tilde{p}(\mathbf{z}^{(\tau)})$
  - Unfavors  $\mathbf{z}^*$  if the proposal distribution  $q$  unduly favors it (i.e., if  $q(\mathbf{z}^*|\mathbf{z}^{(\tau)})$  is large)



# Metropolis-Hastings (MH) Sampling

- Assume a proposal distribution  $q(\mathbf{z}|\mathbf{z}^{(\tau)})$ , e.g.,  $\mathcal{N}(\mathbf{z}|\mathbf{z}^{(\tau)}, \sigma^2 \mathbf{I}_D)$
- In each step, draw  $\mathbf{z}^* \sim q(\mathbf{z}|\mathbf{z}^{(\tau)})$  and accept the sample  $\mathbf{z}^*$  with probability

$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})} \right)$$

- The acceptance probability makes intuitive sense. Note the kind of  $\mathbf{z}^*$  would it favor/unfavor:
  - It favors accepting  $\mathbf{z}^*$  if  $\tilde{p}(\mathbf{z}^*)$  has a higher value than  $\tilde{p}(\mathbf{z}^{(\tau)})$
  - Unfavors  $\mathbf{z}^*$  if the proposal distribution  $q$  unduly favors it (i.e., if  $q(\mathbf{z}^*|\mathbf{z}^{(\tau)})$  is large)
  - Favors  $\mathbf{z}^*$  if we can “reverse” to  $\mathbf{z}^{(\tau)}$  from  $\mathbf{z}^*$  (i.e., if  $q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)$  is large). Needed for good “mixing”

# Metropolis-Hastings (MH) Sampling

- Assume a proposal distribution  $q(\mathbf{z}|\mathbf{z}^{(\tau)})$ , e.g.,  $\mathcal{N}(\mathbf{z}|\mathbf{z}^{(\tau)}, \sigma^2 \mathbf{I}_D)$
- In each step, draw  $\mathbf{z}^* \sim q(\mathbf{z}|\mathbf{z}^{(\tau)})$  and accept the sample  $\mathbf{z}^*$  with probability

$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})} \right)$$

- The acceptance probability makes intuitive sense. Note the kind of  $\mathbf{z}^*$  would it favor/unfavor:
  - It favors accepting  $\mathbf{z}^*$  if  $\tilde{p}(\mathbf{z}^*)$  has a higher value than  $\tilde{p}(\mathbf{z}^{(\tau)})$
  - Unfavors  $\mathbf{z}^*$  if the proposal distribution  $q$  unduly favors it (i.e., if  $q(\mathbf{z}^*|\mathbf{z}^{(\tau)})$  is large)
  - Favors  $\mathbf{z}^*$  if we can “reverse” to  $\mathbf{z}^{(\tau)}$  from  $\mathbf{z}^*$  (i.e., if  $q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)$  is large). Needed for good “mixing”
- Transition probability of the Markov chain in MH sampling:  $T(\mathbf{z}^{(\tau)}, \mathbf{z}^*) = A(\mathbf{z}^*, \mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})$

# Metropolis-Hastings (MH) Sampling

- Assume a proposal distribution  $q(\mathbf{z}|\mathbf{z}^{(\tau)})$ , e.g.,  $\mathcal{N}(\mathbf{z}|\mathbf{z}^{(\tau)}, \sigma^2 \mathbf{I}_D)$
- In each step, draw  $\mathbf{z}^* \sim q(\mathbf{z}|\mathbf{z}^{(\tau)})$  and accept the sample  $\mathbf{z}^*$  with probability


$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})} \right)$$

- The acceptance probability makes intuitive sense. Note the kind of  $\mathbf{z}^*$  would it favor/unfavor:
  - It favors accepting  $\mathbf{z}^*$  if  $\tilde{p}(\mathbf{z}^*)$  has a higher value than  $\tilde{p}(\mathbf{z}^{(\tau)})$
  - Unfavors  $\mathbf{z}^*$  if the proposal distribution  $q$  unduly favors it (i.e., if  $q(\mathbf{z}^*|\mathbf{z}^{(\tau)})$  is large)
  - Favors  $\mathbf{z}^*$  if we can “reverse” to  $\mathbf{z}^{(\tau)}$  from  $\mathbf{z}^*$  (i.e., if  $q(\mathbf{z}^{(\tau)}|\mathbf{z}^*)$  is large). Needed for good “mixing”
- Transition probability of the Markov chain in MH sampling:  $T(\mathbf{z}^{(\tau)}, \mathbf{z}^*) = A(\mathbf{z}^*, \mathbf{z}^{(\tau)})q(\mathbf{z}^*|\mathbf{z}^{(\tau)})$
- **Exercise:** Show that  $T(\mathbf{z}, \mathbf{z}^{(\tau)})$  satisfies the detailed balance property

$$T(\mathbf{z}, \mathbf{z}^{(\tau)})p(\mathbf{z}) = T(\mathbf{z}^{(\tau)}, \mathbf{z})p(\mathbf{z}^{(\tau)})$$



# MH Sampling in Action: A Toy Example..

Target  $p(\mathbf{z}) = \mathcal{N}\left(\begin{bmatrix} 4 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 & 0.8 \\ 0.8 & 1 \end{bmatrix}\right)$ , Proposal  $q(\mathbf{z}^{(t)}|\mathbf{z}^{(t-1)}) = \mathcal{N}\left(\mathbf{z}^{(t-1)}, \begin{bmatrix} 0.01 & 0 \\ 0 & 0.01 \end{bmatrix}\right)$






# MH Sampling in Action: A Toy Example..

Target  $p(\mathbf{z}) = \mathcal{N}\left(\begin{bmatrix} 4 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 & 0.8 \\ 0.8 & 1 \end{bmatrix}\right)$ , Proposal  $q(\mathbf{z}^{(t)}|\mathbf{z}^{(t-1)}) = \mathcal{N}\left(\mathbf{z}^{(t-1)}, \begin{bmatrix} 0.01 & 0 \\ 0 & 0.01 \end{bmatrix}\right)$







# MH Sampling in Action: A Toy Example..

Target  $p(\mathbf{z}) = \mathcal{N}\left(\begin{bmatrix} 4 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 & 0.8 \\ 0.8 & 1 \end{bmatrix}\right)$ , Proposal  $q(\mathbf{z}^{(t)}|\mathbf{z}^{(t-1)}) = \mathcal{N}\left(\mathbf{z}^{(t-1)}, \begin{bmatrix} 0.01 & 0 \\ 0 & 0.01 \end{bmatrix}\right)$








# MH Sampling in Action: A Toy Example..

Target  $p(\mathbf{z}) = \mathcal{N}\left(\begin{bmatrix} 4 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 & 0.8 \\ 0.8 & 1 \end{bmatrix}\right)$ , Proposal  $q(\mathbf{z}^{(t)}|\mathbf{z}^{(t-1)}) = \mathcal{N}\left(\mathbf{z}^{(t-1)}, \begin{bmatrix} 0.01 & 0 \\ 0 & 0.01 \end{bmatrix}\right)$



# MH Sampling in Action: A Toy Example..

Target  $p(\mathbf{z}) = \mathcal{N}\left(\begin{bmatrix} 4 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 & 0.8 \\ 0.8 & 1 \end{bmatrix}\right)$ , Proposal  $q(\mathbf{z}^{(t)}|\mathbf{z}^{(t-1)}) = \mathcal{N}\left(\mathbf{z}^{(t-1)}, \begin{bmatrix} 0.01 & 0 \\ 0 & 0.01 \end{bmatrix}\right)$





# MH Sampling: Some Comments

- Special Case: If proposal distrib. is symmetric, we get [Metropolis Sampling](#) algorithm with


$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})} \right)$$

# MH Sampling: Some Comments

- Special Case: If proposal distrib. is symmetric, we get **Metropolis Sampling** algorithm with

$$A(\mathbf{z}^*, \mathbf{z}^{(\tau)}) = \min \left( 1, \frac{\tilde{p}(\mathbf{z}^*)}{\tilde{p}(\mathbf{z}^{(\tau)})} \right)$$

- Limitation: MH can have a very slow convergence



Generic proposals use

$$Q(x'; x) = \mathcal{N}(x, \sigma^2)$$

$\sigma$  large  $\rightarrow$  many rejections

$\sigma$  small  $\rightarrow$  slow diffusion:

$\sim (L/\sigma)^2$  iterations required

# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$

# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$
- However, suppose we can't sample from  $p(\mathbf{z})$  but can sample from each conditional  $p(z_i | \mathbf{z}_{-i})$

# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$
- However, suppose we can't sample from  $p(\mathbf{z})$  but can sample from each conditional  $p(z_i | \mathbf{z}_{-i})$ 
  - Can we do easily if we have a **locally conjugate model** (e.g., Gaussian matrix factorization)

# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$
- However, suppose we can't sample from  $p(\mathbf{z})$  but can sample from each conditional  $p(z_i | \mathbf{z}_{-i})$ 
  - Can we do easily if we have a **locally conjugate model** (e.g., Gaussian matrix factorization)
- Gibbs sampling uses the conditionals  $p(z_i | \mathbf{z}_{-i})$  as the proposal distribution

# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$
- However, suppose we can't sample from  $p(\mathbf{z})$  but can sample from each conditional  $p(z_i | \mathbf{z}_{-i})$ 
  - Can we do easily if we have a **locally conjugate model** (e.g., Gaussian matrix factorization)
- Gibbs sampling uses the conditionals  $p(z_i | \mathbf{z}_{-i})$  as the proposal distribution
- Gibbs sampling samples from these conditionals in a **cyclic order**

# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$
- However, suppose we can't sample from  $p(\mathbf{z})$  but can sample from each conditional  $p(z_i | \mathbf{z}_{-i})$ 
  - Can we do easily if we have a **locally conjugate model** (e.g., Gaussian matrix factorization)
- Gibbs sampling uses the conditionals  $p(z_i | \mathbf{z}_{-i})$  as the proposal distribution
- Gibbs sampling samples from these conditionals in a **cyclic order**
- Gibbs sampling is equivalent to Metropolis Hastings sampling with acceptance prob. = 1



# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$
- However, suppose we can't sample from  $p(\mathbf{z})$  but can sample from each conditional  $p(z_i | \mathbf{z}_{-i})$ 
  - Can we do easily if we have a **locally conjugate model** (e.g., Gaussian matrix factorization)
- Gibbs sampling uses the conditionals  $p(z_i | \mathbf{z}_{-i})$  as the proposal distribution
- Gibbs sampling samples from these conditionals in a **cyclic order**
- Gibbs sampling is equivalent to Metropolis Hastings sampling with acceptance prob. = 1

$$A(\mathbf{z}^*, \mathbf{z}) = \frac{p(\mathbf{z}^*)q(\mathbf{z} | \mathbf{z}^*)}{p(\mathbf{z})q(\mathbf{z}^* | \mathbf{z})}$$

# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$
- However, suppose we can't sample from  $p(\mathbf{z})$  but can sample from each conditional  $p(z_i | \mathbf{z}_{-i})$ 
  - Can we do it easily if we have a **locally conjugate model** (e.g., Gaussian matrix factorization)
- Gibbs sampling uses the conditionals  $p(z_i | \mathbf{z}_{-i})$  as the proposal distribution
- Gibbs sampling samples from these conditionals in a **cyclic order**
- Gibbs sampling is equivalent to Metropolis Hastings sampling with acceptance prob. = 1

$$A(\mathbf{z}^*, \mathbf{z}) = \frac{p(\mathbf{z}^*)q(\mathbf{z} | \mathbf{z}^*)}{p(\mathbf{z})q(\mathbf{z}^* | \mathbf{z})} = \frac{p(z_i^* | \mathbf{z}_{-i}^*)p(\mathbf{z}_{-i}^*)p(z_i | \mathbf{z}_{-i}^*)}{p(z_i | \mathbf{z}_{-i})p(\mathbf{z}_{-i})p(z_i^* | \mathbf{z}_{-i})}$$

# Gibbs Sampling (Geman & Geman, 1984)

- Suppose we wish to sample from a joint distribution  $p(\mathbf{z})$  where  $\mathbf{z} = (z_1, z_2, \dots, z_M)$
- However, suppose we can't sample from  $p(\mathbf{z})$  but can sample from each conditional  $p(z_i | \mathbf{z}_{-i})$ 
  - Can we do easily if we have a **locally conjugate model** (e.g., Gaussian matrix factorization)
- Gibbs sampling uses the conditionals  $p(z_i | \mathbf{z}_{-i})$  as the proposal distribution
- Gibbs sampling samples from these conditionals in a **cyclic order**
- Gibbs sampling is equivalent to Metropolis Hastings sampling with acceptance prob. = 1

$$A(\mathbf{z}^*, \mathbf{z}) = \frac{p(\mathbf{z}^*)q(\mathbf{z}|\mathbf{z}^*)}{p(\mathbf{z})q(\mathbf{z}^*|\mathbf{z})} = \frac{p(z_i^*|\mathbf{z}_{-i}^*)p(\mathbf{z}_{-i}^*)p(z_i|\mathbf{z}_{-i}^*)}{p(z_i|\mathbf{z}_{-i})p(\mathbf{z}_{-i})p(z_i^*|\mathbf{z}_{-i})} = 1$$

where we use the fact that  $\mathbf{z}_{-i}^* = \mathbf{z}_{-i}$

# Gibbs Sampling: Sketch of the Algorithm

$M$ : Total number of variables,  $T$ : number of Gibbs sampling steps

1. Initialize  $\{z_i : i = 1, \dots, M\}$
2. For  $\tau = 1, \dots, T$ :
  - Sample  $z_1^{(\tau+1)} \sim p(z_1 | z_2^{(\tau)}, z_3^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - Sample  $z_2^{(\tau+1)} \sim p(z_2 | z_1^{(\tau+1)}, z_3^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - $\vdots$
  - Sample  $z_j^{(\tau+1)} \sim p(z_j | z_1^{(\tau+1)}, \dots, z_{j-1}^{(\tau+1)}, z_{j+1}^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - $\vdots$
  - Sample  $z_M^{(\tau+1)} \sim p(z_M | z_1^{(\tau+1)}, z_2^{(\tau+1)}, \dots, z_{M-1}^{(\tau+1)})$ .

# Gibbs Sampling: Sketch of the Algorithm

$M$ : Total number of variables,  $T$ : number of Gibbs sampling steps

1. Initialize  $\{z_i : i = 1, \dots, M\}$
2. For  $\tau = 1, \dots, T$ :
  - Sample  $z_1^{(\tau+1)} \sim p(z_1 | z_2^{(\tau)}, z_3^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - Sample  $z_2^{(\tau+1)} \sim p(z_2 | z_1^{(\tau+1)}, z_3^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - $\vdots$
  - Sample  $z_j^{(\tau+1)} \sim p(z_j | z_1^{(\tau+1)}, \dots, z_{j-1}^{(\tau+1)}, z_{j+1}^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - $\vdots$
  - Sample  $z_M^{(\tau+1)} \sim p(z_M | z_1^{(\tau+1)}, z_2^{(\tau+1)}, \dots, z_{M-1}^{(\tau+1)})$ .

Note: When sampling each variable from its conditional posterior, we use the most recent values of all other variables (this is akin to a co-ordinate ascent like procedure)

# Gibbs Sampling: Sketch of the Algorithm

$M$ : Total number of variables,  $T$ : number of Gibbs sampling steps


1. Initialize  $\{z_i : i = 1, \dots, M\}$
2. For  $\tau = 1, \dots, T$ :
  - Sample  $z_1^{(\tau+1)} \sim p(z_1 | z_2^{(\tau)}, z_3^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - Sample  $z_2^{(\tau+1)} \sim p(z_2 | z_1^{(\tau+1)}, z_3^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - $\vdots$
  - Sample  $z_j^{(\tau+1)} \sim p(z_j | z_1^{(\tau+1)}, \dots, z_{j-1}^{(\tau+1)}, z_{j+1}^{(\tau)}, \dots, z_M^{(\tau)})$ .
  - $\vdots$
  - Sample  $z_M^{(\tau+1)} \sim p(z_M | z_1^{(\tau+1)}, z_2^{(\tau+1)}, \dots, z_{M-1}^{(\tau+1)})$ .

Note: When sampling each variable from its conditional posterior, we use the most recent values of all other variables (this is akin to a co-ordinate ascent like procedure)

Note: Order of updating the variables *usually* doesn't matter (but see “Scan Order in Gibbs Sampling: Models in Which it Matters and Bounds on How Much” from NIPS 2016)


# Gibbs Sampling: A Simple Example

Can sample from a 2-D Gaussian using 1-D Gaussians (recall that if the joint distribution is a 2-D Gaussian, conditionals will simply be 1-D Gaussians)



# Gibbs Sampling: A Simple Example


Can sample from a 2-D Gaussian using 1-D Gaussians (recall that if the joint distribution is a 2-D Gaussian, conditionals will simply be 1-D Gaussians)






# Gibbs Sampling: A Simple Example

Can sample from a 2-D Gaussian using 1-D Gaussians (recall that if the joint distribution is a 2-D Gaussian, conditionals will simply be 1-D Gaussians)



# Gibbs Sampling: A Simple Example

Can sample from a 2-D Gaussian using 1-D Gaussians (recall that if the joint distribution is a 2-D Gaussian, conditionals will simply be 1-D Gaussians)



# Next Class..

- More examples of Gibbs sampling
- Random-walk avoiding MCMC methods
- “Using” MCMC. Pros and Cons.
- Some recent advances in MCMC