

Mixture Models and GMM (Contd.)


Piyush Rai

Probabilistic Machine Learning (CS772A)

Aug 31, 2017


Recap: Gaussian Mixture Model

- A model for **data clustering** and **density estimation**
- Assumes data generated from a mixture of K Gaussians with mixing proportions π_1, \dots, π_K



Recap: Gaussian Mixture Model

- A model for **data clustering** and **density estimation**
- Assumes data generated from a mixture of K Gaussians with mixing proportions π_1, \dots, π_K




- The **prior probability** of observation \mathbf{x}_n generated from the k -th Gaussian

$$p(\mathbf{z}_n = k|\pi) = \pi_k$$

Recap: Gaussian Mixture Model

- A model for **data clustering** and **density estimation**
- Assumes data generated from a mixture of K Gaussians with mixing proportions π_1, \dots, π_K




- The **prior probability** of observation \mathbf{x}_n generated from the k -th Gaussian

$$p(\mathbf{z}_n = k|\pi) = \pi_k$$

- Same as a **multinoulli prior** on \mathbf{z} , i.e., $p(\mathbf{z}_n|\pi) = \prod_{k=1}^K \pi_k^{z_{nk}}$

Recap: Gaussian Mixture Model

- A model for **data clustering** and **density estimation**
- Assumes data generated from a mixture of K Gaussians with mixing proportions π_1, \dots, π_K




- The **prior probability** of observation \mathbf{x}_n generated from the k -th Gaussian

$$p(\mathbf{z}_n = k | \pi) = \pi_k$$

- Same as a **multinoulli prior** on \mathbf{z} , i.e., $p(\mathbf{z}_n | \pi) = \prod_{k=1}^K \pi_k^{z_{nk}}$ (note $z_{nk} = 1$ if $\mathbf{z}_n = k$; 0 otherwise)

Recap: Gaussian Mixture Model

- A model for **data clustering** and **density estimation**
- Assumes data generated from a mixture of K Gaussians with mixing proportions π_1, \dots, π_K




- The **prior probability** of observation \mathbf{x}_n generated from the k -th Gaussian

$$p(\mathbf{z}_n = k | \pi) = \pi_k$$

- Same as a **multinoulli prior** on \mathbf{z} , i.e., $p(\mathbf{z}_n | \pi) = \prod_{k=1}^K \pi_k^{z_{nk}}$ (note $z_{nk} = 1$ if $\mathbf{z}_n = k$; 0 otherwise)
- If $\mathbf{z}_n = k$, we generate \mathbf{x}_n from the k -th Gaussian.

Recap: Gaussian Mixture Model

- A model for **data clustering** and **density estimation**
- Assumes data generated from a mixture of K Gaussians with mixing proportions π_1, \dots, π_K



- The **prior probability** of observation \mathbf{x}_n generated from the k -th Gaussian

$$p(\mathbf{z}_n = k | \pi) = \pi_k$$

- Same as a **multinoulli prior** on \mathbf{z} , i.e., $p(\mathbf{z}_n | \pi) = \prod_{k=1}^K \pi_k^{z_{nk}}$ (note $z_{nk} = 1$ if $\mathbf{z}_n = k$; 0 otherwise)
- If $\mathbf{z}_n = k$, we generate \mathbf{x}_n from the k -th Gaussian. Thus

$$p(\mathbf{x}_n | \mathbf{z}_n = k) = \mathcal{N}(\mathbf{x}_n | \mu_k, \Sigma_k)$$

Recap: Gaussian Mixture Model (GMM)

- The marginal distribution of \mathbf{x}_n (requires summing over all possibilities of \mathbf{z}_n)

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

Recap: Gaussian Mixture Model (GMM)

- The marginal distribution of \mathbf{x}_n (requires summing over all possibilities of \mathbf{z}_n)

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

Recap: Gaussian Mixture Model (GMM)

- The marginal distribution of \mathbf{x}_n (requires summing over all possibilities of \mathbf{z}_n)

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- The MLE objective for the GMM will be

$$\sum_{n=1}^N \log p(\mathbf{x}_n|\Theta) = \sum_{n=1}^N \log \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

Recap: Gaussian Mixture Model (GMM)

- The marginal distribution of \mathbf{x}_n (requires summing over all possibilities of \mathbf{z}_n)

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- The MLE objective for the GMM will be

$$\sum_{n=1}^N \log p(\mathbf{x}_n|\Theta) = \sum_{n=1}^N \log \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

- Doing MLE on this objective is tricky due to the “ $\log \sum$ ” term

Recap: Gaussian Mixture Model (GMM)

- The marginal distribution of \mathbf{x}_n (requires summing over all possibilities of \mathbf{z}_n)

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- The MLE objective for the GMM will be

$$\sum_{n=1}^N \log p(\mathbf{x}_n|\Theta) = \sum_{n=1}^N \log \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

- Doing MLE on this objective is tricky due to the “ $\log \sum$ ” term
 - Parameters get coupled; no closed form solution (iterative methods needed, slow convergence)

Recap: Gaussian Mixture Model (GMM)

- The marginal distribution of \mathbf{x}_n (requires summing over all possibilities of \mathbf{z}_n)

$$p(\mathbf{x}_n|\Theta) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

where $\Theta = (\pi, \{\mu_k, \Sigma_k\}_{k=1}^K)$

- The MLE objective for the GMM will be

$$\sum_{n=1}^N \log p(\mathbf{x}_n|\Theta) = \sum_{n=1}^N \log \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n|\mu_k, \Sigma_k)$$

- Doing MLE on this objective is tricky due to the “ $\log \sum$ ” term
 - Parameters get coupled; no closed form solution (iterative methods needed, slow convergence)
- An alternating “guess, re-estimate, and repeat until converge” algorithm helps solve such problems in a clean, simple, and efficient way; basically, the [Expectation Maximization](#) (EM) algorithm

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the z_n for each x_n

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the \mathbf{z}_n for each \mathbf{x}_n
- Reason: If \mathbf{z}_n is known (suppose $\mathbf{z}_n = k$), then the summation over \mathbf{z}_n isn't required

$$\sum_{n=1}^N \log \cancel{\sum_{k=1}^K} \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the \mathbf{z}_n for each \mathbf{x}_n
- Reason: If \mathbf{z}_n is known (suppose $\mathbf{z}_n = k$), then the summation over \mathbf{z}_n isn't required

$$\sum_{n=1}^N \log \cancel{\sum_{k=1}^K} \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

- With \mathbf{z}_n known, can treat $(\mathbf{x}_n, \mathbf{z}_n)$ as our “data” and do **MLE on $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$** , instead of $p(\mathbf{x}_n | \Theta)$

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the \mathbf{z}_n for each \mathbf{x}_n
- Reason: If \mathbf{z}_n is known (suppose $\mathbf{z}_n = k$), then the summation over \mathbf{z}_n isn't required

$$\sum_{n=1}^N \log \cancel{\sum_{k=1}^K} \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

- With \mathbf{z}_n known, can treat $(\mathbf{x}_n, \mathbf{z}_n)$ as our “data” and do MLE on $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$, instead of $p(\mathbf{x}_n | \Theta)$
 - $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$ is known as “complete data likelihood” (CLL) - \mathbf{z}_n makes \mathbf{x}_n “complete”

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the \mathbf{z}_n for each \mathbf{x}_n
- Reason: If \mathbf{z}_n is known (suppose $\mathbf{z}_n = k$), then the summation over \mathbf{z}_n isn't required

$$\sum_{n=1}^N \log \cancel{\sum_{k=1}^K} \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

- With \mathbf{z}_n known, can treat $(\mathbf{x}_n, \mathbf{z}_n)$ as our “data” and do MLE on $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$, instead of $p(\mathbf{x}_n | \Theta)$
 - $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$ is known as “complete data likelihood” (CLL) - \mathbf{z}_n makes \mathbf{x}_n “complete”
 - $p(\mathbf{x}_n | \Theta)$ is known as “incomplete data likelihood” (ILL)

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the \mathbf{z}_n for each \mathbf{x}_n
- Reason: If \mathbf{z}_n is known (suppose $\mathbf{z}_n = k$), then the summation over \mathbf{z}_n isn't required

$$\sum_{n=1}^N \log \cancel{\sum_{k=1}^K} \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

- With \mathbf{z}_n known, can treat $(\mathbf{x}_n, \mathbf{z}_n)$ as our “data” and do MLE on $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$, instead of $p(\mathbf{x}_n | \Theta)$
 - $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$ is known as “complete data likelihood” (CLL) - \mathbf{z}_n makes \mathbf{x}_n “complete”
 - $p(\mathbf{x}_n | \Theta)$ is known as “incomplete data likelihood” (ILL)
- Notation: $\mathbf{z}_n = [z_{n1}, z_{n2}, \dots, z_{nK}]$ has a one-hot representation (note: only a single z_{nk} will be 1)

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the \mathbf{z}_n for each \mathbf{x}_n
- Reason: If \mathbf{z}_n is known (suppose $\mathbf{z}_n = k$), then the summation over \mathbf{z}_n isn't required

$$\sum_{n=1}^N \log \cancel{\sum_{k=1}^K} \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

- With \mathbf{z}_n known, can treat $(\mathbf{x}_n, \mathbf{z}_n)$ as our “data” and do MLE on $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$, instead of $p(\mathbf{x}_n | \Theta)$
 - $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$ is known as “complete data likelihood” (CLL) - \mathbf{z}_n makes \mathbf{x}_n “complete”
 - $p(\mathbf{x}_n | \Theta)$ is known as “incomplete data likelihood” (ILL)
- Notation: $\mathbf{z}_n = [z_{n1}, z_{n2}, \dots, z_{nK}]$ has a one-hot representation (note: only a single z_{nk} will be 1)
- Denoting all the GMM parameters by Θ , the complete data log-likelihood will be

$$\text{CLL}(\Theta) = \sum_{n=1}^N \log p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$$

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the \mathbf{z}_n for each \mathbf{x}_n
- Reason: If \mathbf{z}_n is known (suppose $\mathbf{z}_n = k$), then the summation over \mathbf{z}_n isn't required

$$\sum_{n=1}^N \log \cancel{\sum_{k=1}^K} \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

- With \mathbf{z}_n known, can treat $(\mathbf{x}_n, \mathbf{z}_n)$ as our “data” and do MLE on $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$, instead of $p(\mathbf{x}_n | \Theta)$
 - $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$ is known as “complete data likelihood” (CLL) - \mathbf{z}_n makes \mathbf{x}_n “complete”
 - $p(\mathbf{x}_n | \Theta)$ is known as “incomplete data likelihood” (ILL)
- Notation: $\mathbf{z}_n = [z_{n1}, z_{n2}, \dots, z_{nK}]$ has a one-hot representation (note: only a single z_{nk} will be 1)
- Denoting all the GMM parameters by Θ , the complete data log-likelihood will be

$$\text{CLL}(\Theta) = \sum_{n=1}^N \log p(\mathbf{x}_n, \mathbf{z}_n | \Theta) = \sum_{n=1}^N \log p(\mathbf{x}_n | \mathbf{z}_n) p(\mathbf{z}_n)$$

GMM Parameter Estimation

- MLE for GMM (and LVMs in general) becomes simple if we knew (or “guess”) the \mathbf{z}_n for each \mathbf{x}_n
- Reason: If \mathbf{z}_n is known (suppose $\mathbf{z}_n = k$), then the summation over \mathbf{z}_n isn't required

$$\sum_{n=1}^N \log \cancel{\sum_{k=1}^K} \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

- With \mathbf{z}_n known, can treat $(\mathbf{x}_n, \mathbf{z}_n)$ as our “data” and do MLE on $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$, instead of $p(\mathbf{x}_n | \Theta)$
 - $p(\mathbf{x}_n, \mathbf{z}_n | \Theta)$ is known as “complete data likelihood” (CLL) - \mathbf{z}_n makes \mathbf{x}_n “complete”
 - $p(\mathbf{x}_n | \Theta)$ is known as “incomplete data likelihood” (ILL)
- Notation: $\mathbf{z}_n = [z_{n1}, z_{n2}, \dots, z_{nK}]$ has a one-hot representation (note: only a single z_{nk} will be 1)
- Denoting all the GMM parameters by Θ , the complete data log-likelihood will be

$$\text{CLL}(\Theta) = \sum_{n=1}^N \log p(\mathbf{x}_n, \mathbf{z}_n | \Theta) = \sum_{n=1}^N \log p(\mathbf{x}_n | \mathbf{z}_n) p(\mathbf{z}_n) = \sum_{n=1}^N \log \prod_{k=1}^K [p(\mathbf{x}_n | \mathbf{z}_n = k) p(\mathbf{z}_n = k)]^{z_{nk}}$$

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k)$$

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.).

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, **we need the z_{nk} 's (each a binary r.v.)**. This is where we'll make a guess.

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.). This is where we'll make a guess.
 - Idea: Use posterior expectation of z_{nk} as our guess

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.). This is where we'll make a guess.
 - Idea: Use posterior expectation of z_{nk} as our guess (seems reasonable; will see justification shortly)

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.). This is where we'll make a guess.
 - Idea: Use posterior expectation of z_{nk} as our guess (seems reasonable; will see justification shortly)

$$\mathbb{E}[z_{nk}] = 0 \times p(z_{nk} = 0 | \mathbf{x}_n) + 1 \times p(z_{nk} = 1 | \mathbf{x}_n)$$

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.). This is where we'll make a guess.
 - Idea: Use posterior expectation of z_{nk} as our guess (seems reasonable; will see justification shortly)

$$\begin{aligned} \mathbb{E}[z_{nk}] &= 0 \times p(z_{nk} = 0 | \mathbf{x}_n) + 1 \times p(z_{nk} = 1 | \mathbf{x}_n) \\ &= p(z_{nk} = 1 | \mathbf{x}_n) \end{aligned}$$

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.). This is where we'll make a guess.
 - Idea: Use posterior expectation of z_{nk} as our guess (seems reasonable; will see justification shortly)

$$\begin{aligned} \mathbb{E}[z_{nk}] &= 0 \times p(z_{nk} = 0 | \mathbf{x}_n) + 1 \times p(z_{nk} = 1 | \mathbf{x}_n) \\ &= p(z_{nk} = 1 | \mathbf{x}_n) \\ &\propto p(z_{nk} = 1) p(\mathbf{x}_n | z_{nk} = 1) \quad (\text{from Bayes Rule}) \end{aligned}$$

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.). This is where we'll make a guess.
 - Idea: Use posterior expectation of z_{nk} as our guess (seems reasonable; will see justification shortly)

$$\mathbb{E}[z_{nk}] = 0 \times p(z_{nk} = 0 | \mathbf{x}_n) + 1 \times p(z_{nk} = 1 | \mathbf{x}_n)$$

$$= p(z_{nk} = 1 | \mathbf{x}_n)$$

$$\propto p(z_{nk} = 1) p(\mathbf{x}_n | z_{nk} = 1) \quad (\text{from Bayes Rule})$$

$$\text{Thus } \mathbb{E}[z_{nk}] \propto \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \quad (\text{Posterior prob. that } \mathbf{x}_n \text{ is generated by } k\text{-th Gaussian})$$

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.). This is where we'll make a guess.
 - Idea: Use posterior expectation of z_{nk} as our guess (seems reasonable; will see justification shortly)

$$\mathbb{E}[z_{nk}] = 0 \times p(z_{nk} = 0 | \mathbf{x}_n) + 1 \times p(z_{nk} = 1 | \mathbf{x}_n)$$

$$= p(z_{nk} = 1 | \mathbf{x}_n)$$

$$\propto p(z_{nk} = 1) p(\mathbf{x}_n | z_{nk} = 1) \quad (\text{from Bayes Rule})$$

$$\text{Thus } \mathbb{E}[z_{nk}] \propto \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \quad (\text{Posterior prob. that } \mathbf{x}_n \text{ is generated by } k\text{-th Gaussian})$$

- Note: We can finally normalize $\mathbb{E}[z_{nk}]$ as $\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$ since $\sum_{k=1}^K \mathbb{E}[z_{nk}] = 1$

GMM Parameter Estimation

- The CLL gets further simplified to

$$\text{CLL}(\Theta) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} \log p(\mathbf{z}_n = k) p(\mathbf{x}_n | \mathbf{z}_n = k) = \sum_{n=1}^N \sum_{k=1}^K z_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Simple algebraic form. MLE easy by taking (partial) derivatives w.r.t. each parameter
- However, to do so, we need the z_{nk} 's (each a binary r.v.). This is where we'll make a guess.
 - Idea: Use posterior expectation of z_{nk} as our guess (seems reasonable; will see justification shortly)

$$\mathbb{E}[z_{nk}] = 0 \times p(z_{nk} = 0 | \mathbf{x}_n) + 1 \times p(z_{nk} = 1 | \mathbf{x}_n)$$

$$= p(z_{nk} = 1 | \mathbf{x}_n)$$

$$\propto p(z_{nk} = 1) p(\mathbf{x}_n | z_{nk} = 1) \quad (\text{from Bayes Rule})$$

$$\text{Thus } \mathbb{E}[z_{nk}] \propto \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \quad (\text{Posterior prob. that } \mathbf{x}_n \text{ is generated by } k\text{-th Gaussian})$$

- Note: We can finally normalize $\mathbb{E}[z_{nk}]$ as $\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$ since $\sum_{k=1}^K \mathbb{E}[z_{nk}] = 1$
- **Wait!** Computing $\mathbb{E}[z_{nk}]$ requires knowing $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ (chicken-and-egg problem ☺)

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an [alternating approach](#)

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an [alternating approach](#)

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known.

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}]$$

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \mu_k, \Sigma_k)}{\sum_{\ell=1}^K \pi_{\ell} \mathcal{N}(\mathbf{x}_n | \mu_{\ell}, \Sigma_{\ell})}$$

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \mu_k, \Sigma_k)}{\sum_{\ell=1}^K \pi_{\ell} \mathcal{N}(\mathbf{x}_n | \mu_{\ell}, \Sigma_{\ell})} \quad (\text{for } n = 1, \dots, N, k = 1, \dots, K)$$

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_{\ell} \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_{\ell}, \boldsymbol{\Sigma}_{\ell})} \quad (\text{for } n = 1, \dots, N, k = 1, \dots, K)$$

- 2 Using $\mathbb{E}[z_{nk}]$ computed in step 1, **maximize** the expected CLL objective w.r.t. $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_{\ell} \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_{\ell}, \boldsymbol{\Sigma}_{\ell})} \quad (\text{for } n = 1, \dots, N, k = 1, \dots, K)$$

- 2 Using $\mathbb{E}[z_{nk}]$ computed in step 1, **maximize** the expected CLL objective w.r.t. $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \mathbb{E}[z_{nk}] [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_{\ell} \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_{\ell}, \boldsymbol{\Sigma}_{\ell})} \quad (\text{for } n = 1, \dots, N, k = 1, \dots, K)$$

- 2 Using $\mathbb{E}[z_{nk}]$ computed in step 1, **maximize** the expected CLL objective w.r.t. $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \mathbb{E}[z_{nk}] [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

.. this will give ML estimates for parameters $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$. Details on the next slides.

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_{\ell} \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_{\ell}, \boldsymbol{\Sigma}_{\ell})} \quad (\text{for } n = 1, \dots, N, k = 1, \dots, K)$$

- 2 Using $\mathbb{E}[z_{nk}]$ computed in step 1, **maximize** the expected CLL objective w.r.t. $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \mathbb{E}[z_{nk}] [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

.. this will give ML estimates for parameters $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$. Details on the next slides.

The algorithm alternates between step 1 and 2 until convergence

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_{\ell} \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_{\ell}, \boldsymbol{\Sigma}_{\ell})} \quad (\text{for } n = 1, \dots, N, k = 1, \dots, K)$$

- 2 Using $\mathbb{E}[z_{nk}]$ computed in step 1, **maximize** the expected CLL objective w.r.t. $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \mathbb{E}[z_{nk}] [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

.. this will give ML estimates for parameters $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$. Details on the next slides.

The algorithm alternates between step 1 and 2 until convergence

This is **an example** of the more general **Expectation Maximization** (EM) algorithm.

GMM Parameter Estimation: The Alternating Approach

Can solve the chicken-and-egg problem by taking an **alternating approach**

- 1 Assume $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$ is known. Estimate the required **expectations** of the latent variables

$$\mathbb{E}[z_{nk}] = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_{\ell} \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_{\ell}, \boldsymbol{\Sigma}_{\ell})} \quad (\text{for } n = 1, \dots, N, k = 1, \dots, K)$$

- 2 Using $\mathbb{E}[z_{nk}]$ computed in step 1, **maximize** the expected CLL objective w.r.t. $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \mathbb{E}[z_{nk}] [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

.. this will give ML estimates for parameters $\Theta = \{\pi_k, \mu_k, \Sigma_k\}$. Details on the next slides.

The algorithm alternates between step 1 and 2 until convergence

This is an **example** of the more general **Expectation Maximization** (EM) algorithm. EM can be used for MLE/MAP in **probabilistic models that contain latent variables** making standard MLE/MAP hard

GMM Parameter Estimation

- Given $\mathbb{E}[z_{nk}] = \gamma_{nk} = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$, the expected complete data log-lik.

GMM Parameter Estimation

- Given $\mathbb{E}[z_{nk}] = \gamma_{nk} = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$, the expected complete data log-lik.

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \gamma_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

GMM Parameter Estimation

- Given $\mathbb{E}[z_{nk}] = \gamma_{nk} = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$, the expected complete data log-lik.

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \gamma_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Taking derivatives w.r.t. $\boldsymbol{\mu}_k$ and $\boldsymbol{\Sigma}_k$, $\forall k = 1, \dots, K$

GMM Parameter Estimation

- Given $\mathbb{E}[z_{nk}] = \gamma_{nk} = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$, the expected complete data log-lik.

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \gamma_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Taking derivatives w.r.t. $\boldsymbol{\mu}_k$ and $\boldsymbol{\Sigma}_k$, $\forall k = 1, \dots, K$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\mu}_k} = \frac{\partial}{\partial \boldsymbol{\mu}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\mu}_k \text{ can be ignored})$$

GMM Parameter Estimation

- Given $\mathbb{E}[z_{nk}] = \gamma_{nk} = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$, the expected complete data log-lik.

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \gamma_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Taking derivatives w.r.t. $\boldsymbol{\mu}_k$ and $\boldsymbol{\Sigma}_k$, $\forall k = 1, \dots, K$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\mu}_k} = \frac{\partial}{\partial \boldsymbol{\mu}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\mu}_k \text{ can be ignored})$$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\Sigma}_k} = \frac{\partial}{\partial \boldsymbol{\Sigma}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\Sigma}_k \text{ can be ignored})$$

GMM Parameter Estimation

- Given $\mathbb{E}[z_{nk}] = \gamma_{nk} = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$, the expected complete data log-lik.

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \gamma_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Taking derivatives w.r.t. $\boldsymbol{\mu}_k$ and $\boldsymbol{\Sigma}_k$, $\forall k = 1, \dots, K$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\mu}_k} = \frac{\partial}{\partial \boldsymbol{\mu}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\mu}_k \text{ can be ignored})$$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\Sigma}_k} = \frac{\partial}{\partial \boldsymbol{\Sigma}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\Sigma}_k \text{ can be ignored})$$

- For each k , it's a “weighted” version of the MLE for the multivar. Gaussian $\mathcal{N}(\mathbf{x} | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$

GMM Parameter Estimation

- Given $\mathbb{E}[z_{nk}] = \gamma_{nk} = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$, the expected complete data log-lik.

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \gamma_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Taking derivatives w.r.t. $\boldsymbol{\mu}_k$ and $\boldsymbol{\Sigma}_k$, $\forall k = 1, \dots, K$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\mu}_k} = \frac{\partial}{\partial \boldsymbol{\mu}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\mu}_k \text{ can be ignored})$$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\Sigma}_k} = \frac{\partial}{\partial \boldsymbol{\Sigma}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\Sigma}_k \text{ can be ignored})$$

- For each k , it's a “weighted” version of the MLE for the multivar. Gaussian $\mathcal{N}(\mathbf{x} | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$
- Can also solve for π_k likewise (subject to constraint $\sum_{k=1}^K \pi_k = 1$)

GMM Parameter Estimation

- Given $\mathbb{E}[z_{nk}] = \gamma_{nk} = \frac{\pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_{\ell=1}^K \pi_\ell \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_\ell, \boldsymbol{\Sigma}_\ell)}$, the expected complete data log-lik.

$$\mathcal{L} = \sum_{n=1}^N \sum_{k=1}^K \gamma_{nk} [\log \pi_k + \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)]$$

- Taking derivatives w.r.t. $\boldsymbol{\mu}_k$ and $\boldsymbol{\Sigma}_k$, $\forall k = 1, \dots, K$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\mu}_k} = \frac{\partial}{\partial \boldsymbol{\mu}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\mu}_k \text{ can be ignored})$$

$$\frac{\partial \mathcal{L}}{\partial \boldsymbol{\Sigma}_k} = \frac{\partial}{\partial \boldsymbol{\Sigma}_k} \sum_{n=1}^N \gamma_{nk} \log \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) = 0 \quad (\text{note: constants w.r.t. } \boldsymbol{\Sigma}_k \text{ can be ignored})$$

- For each k , it's a "weighted" version of the MLE for the multivar. Gaussian $\mathcal{N}(\mathbf{x} | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$
- Can also solve for π_k likewise (subject to constraint $\sum_{k=1}^K \pi_k = 1$)
- Derivations are a bit tedious (but straightforward). I will provide a note.

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k
 - Note that each x_n gets “soft” assignment $\gamma_{nk} \in (0, 1)$ to the k -th Gaussian

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k
 - Note that each x_n gets “soft” assignment $\gamma_{nk} \in (0, 1)$ to the k -th Gaussian (also $\sum_{k=1}^K \gamma_{nk} = 1$)

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k
 - Note that each x_n gets “soft” assignment $\gamma_{nk} \in (0, 1)$ to the k -th Gaussian (also $\sum_{k=1}^K \gamma_{nk} = 1$)
- The final expressions for updates of $\{\pi_k, \mu_k, \Sigma_k\}_{k=1}^K$

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k
 - Note that each \mathbf{x}_n gets “soft” assignment $\gamma_{nk} \in (0, 1)$ to the k -th Gaussian (also $\sum_{k=1}^K \gamma_{nk} = 1$)
- The final expressions for updates of $\{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$

$$\boldsymbol{\mu}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} \mathbf{x}_n$$

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k
 - Note that each \mathbf{x}_n gets “soft” assignment $\gamma_{nk} \in (0, 1)$ to the k -th Gaussian (also $\sum_{k=1}^K \gamma_{nk} = 1$)
- The final expressions for updates of $\{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$

$$\boldsymbol{\mu}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} \mathbf{x}_n$$

$$\boldsymbol{\Sigma}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} (\mathbf{x}_n - \boldsymbol{\mu}_k)(\mathbf{x}_n - \boldsymbol{\mu}_k)^\top$$

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k
 - Note that each \mathbf{x}_n gets “soft” assignment $\gamma_{nk} \in (0, 1)$ to the k -th Gaussian (also $\sum_{k=1}^K \gamma_{nk} = 1$)
- The final expressions for updates of $\{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$

$$\boldsymbol{\mu}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} \mathbf{x}_n$$

$$\boldsymbol{\Sigma}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} (\mathbf{x}_n - \boldsymbol{\mu}_k)(\mathbf{x}_n - \boldsymbol{\mu}_k)^\top$$

$$\pi_k = \frac{N_k}{N}$$

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k
 - Note that each \mathbf{x}_n gets “soft” assignment $\gamma_{nk} \in (0, 1)$ to the k -th Gaussian (also $\sum_{k=1}^K \gamma_{nk} = 1$)
- The final expressions for updates of $\{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$

$$\boldsymbol{\mu}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} \mathbf{x}_n$$

$$\boldsymbol{\Sigma}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} (\mathbf{x}_n - \boldsymbol{\mu}_k)(\mathbf{x}_n - \boldsymbol{\mu}_k)^\top$$

$$\pi_k = \frac{N_k}{N}$$

- Note that μ_k, Σ_k updates are similar to multivar. Gaussian MLE equations

GMM Parameter Estimation

- Suppose $N_k = \sum_{n=1}^N \gamma_{nk}$ is the effective number of observations assigned to Gaussian k
 - Note that each \mathbf{x}_n gets “soft” assignment $\gamma_{nk} \in (0, 1)$ to the k -th Gaussian (also $\sum_{k=1}^K \gamma_{nk} = 1$)
- The final expressions for updates of $\{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$

$$\boldsymbol{\mu}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} \mathbf{x}_n$$

$$\boldsymbol{\Sigma}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} (\mathbf{x}_n - \boldsymbol{\mu}_k)(\mathbf{x}_n - \boldsymbol{\mu}_k)^\top$$

$$\pi_k = \frac{N_k}{N}$$

- Note that $\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k$ updates are similar to multivar. Gaussian MLE equations
- Each $\mathbf{x}_n, n = 1, \dots, N$ contributes to each $\{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ update but **fractionally** (based on γ_{nk})

Summary: EM for Learning GMM

- Initialize the parameters $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ randomly, or using K -means

Summary: EM for Learning GMM

- Initialize the parameters $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ randomly, or using K -means
- Iterate until convergence (e.g., when $\log p(\mathbf{x}|\Theta)$ ceases to increase or Θ doesn't change by much)

Summary: EM for Learning GMM

- Initialize the parameters $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ randomly, or using K -means
- Iterate until convergence (e.g., when $\log p(\mathbf{x}|\Theta)$ ceases to increase or Θ doesn't change by much)
 - Given Θ , compute each expectation z_{nk} (posterior probability of $z_{nk} = 1$), $\forall n, k$

Summary: EM for Learning GMM

- Initialize the parameters $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ randomly, or using K -means
- Iterate until convergence (e.g., when $\log p(\mathbf{x}|\Theta)$ ceases to increase or Θ doesn't change by much)
 - Given Θ , compute each expectation z_{nk} (posterior probability of $z_{nk} = 1$), $\forall n, k$

$$\gamma_{nk} = \mathbb{E}[z_{nk}] \propto \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \quad (\text{and re-normalize s.t. } \sum_{k=1}^K \gamma_{nk} = 1)$$

Summary: EM for Learning GMM

- Initialize the parameters $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ randomly, or using K -means
- Iterate until convergence (e.g., when $\log p(\mathbf{x}|\Theta)$ ceases to increase or Θ doesn't change by much)
 - Given Θ , compute each expectation z_{nk} (posterior probability of $z_{nk} = 1$), $\forall n, k$

$$\gamma_{nk} = \mathbb{E}[z_{nk}] \propto \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \quad (\text{and re-normalize s.t. } \sum_{k=1}^K \gamma_{nk} = 1)$$

- Given “responsibilities” $\gamma_{nk} = \mathbb{E}[z_{nk}]$, and $N_k = \sum_{n=1}^N \gamma_{nk}$, update $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ as

Summary: EM for Learning GMM

- Initialize the parameters $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ randomly, or using K -means
- Iterate until convergence (e.g., when $\log p(\mathbf{x}|\Theta)$ ceases to increase or Θ doesn't change by much)
 - Given Θ , compute each expectation z_{nk} (posterior probability of $z_{nk} = 1$), $\forall n, k$

$$\gamma_{nk} = \mathbb{E}[z_{nk}] \propto \pi_k \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \quad (\text{and re-normalize s.t. } \sum_{k=1}^K \gamma_{nk} = 1)$$

- Given “responsibilities” $\gamma_{nk} = \mathbb{E}[z_{nk}]$, and $N_k = \sum_{n=1}^N \gamma_{nk}$, update $\Theta = \{\pi_k, \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k\}_{k=1}^K$ as


$$\boldsymbol{\mu}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} \mathbf{x}_n$$

$$\boldsymbol{\Sigma}_k = \frac{1}{N_k} \sum_{n=1}^N \gamma_{nk} (\mathbf{x}_n - \boldsymbol{\mu}_k)(\mathbf{x}_n - \boldsymbol{\mu}_k)^\top$$

$$\pi_k = \frac{N_k}{N}$$


GMM: Some Important Aspects

- GMM learns a probabilistic (“soft”) clustering as opposed to **hard clustering** (e.g., K -means)




GMM: Some Important Aspects

- GMM learns a probabilistic (“soft”) clustering as opposed to **hard clustering** (e.g., K -means)




- GMM doesn't assume the clusters to be spherical and equi-sized as opposed to K -means (recall that each Gaussian has a specific covariance which can control the shape of that cluster)




GMM: Some Important Aspects

- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



GMM: Some Important Aspects


- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit.

GMM: Some Important Aspects


- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit. Some remedies

GMM: Some Important Aspects


- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit. Some remedies
 - Assume the covariance matrix to be **diagonal** or **spherical** (but results might be poorer)

GMM: Some Important Aspects


- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit. Some remedies
 - Assume the covariance matrix to be **diagonal** or **spherical** (but results might be poorer)
 - Use **priors** on the parameters and do **MAP estimation** for parameters in the M step of EM

GMM: Some Important Aspects


- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit. Some remedies
 - Assume the covariance matrix to be **diagonal** or **spherical** (but results might be poorer)
 - Use **priors** on the parameters and do **MAP estimation** for parameters in the M step of EM
 - Can do **fully Bayesian inference** for GMM

GMM: Some Important Aspects


- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit. Some remedies
 - Assume the covariance matrix to be **diagonal** or **spherical** (but results might be poorer)
 - Use **priors** on the parameters and do **MAP estimation** for parameters in the M step of EM
 - Can do **fully Bayesian inference** for GMM
 - Helps learn K using marginal likelihood or **nonparametric Bayesian methods**

GMM: Some Important Aspects


- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit. Some remedies
 - Assume the covariance matrix to be **diagonal** or **spherical** (but results might be poorer)
 - Use **priors** on the parameters and do **MAP estimation** for parameters in the M step of EM
 - Can do **fully Bayesian inference** for GMM
 - Helps learn K using marginal likelihood or **nonparametric Bayesian methods**
 - Helps learn the hyperparameters

GMM: Some Important Aspects


- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit. Some remedies
 - Assume the covariance matrix to be **diagonal** or **spherical** (but results might be poorer)
 - Use **priors** on the parameters and do **MAP estimation** for parameters in the M step of EM
 - Can do **fully Bayesian inference** for GMM
 - Helps learn K using marginal likelihood or **nonparametric Bayesian methods**
 - Helps learn the hyperparameters
 - More on these aspects later..

GMM: Some Important Aspects

- GMM, just like K -means, can be sensitive to initialization (EM only converges to a local optima)



- Learning full covariances can be tricky: $\mathcal{O}(D^2)$ params to learn and can overfit. Some remedies
 - Assume the covariance matrix to be **diagonal** or **spherical** (but results might be poorer)
 - Use **priors** on the parameters and do **MAP estimation** for parameters in the M step of EM
 - Can do **fully Bayesian inference** for GMM
 - Helps learn K using marginal likelihood or **nonparametric Bayesian methods**
 - Helps learn the hyperparameters
 - More on these aspects later..
 - Use **low-rank Gaussians** for each mixture component (**mixture of factor analyzers**)

Mixture Models: Applications beyond Clustering

- Mixture models is a general framework for modeling grouped data

Mixture Models: Applications beyond Clustering


- Mixture models is a general framework for modeling grouped data
- Not limited only to clustering or density estimation

Mixture Models: Applications beyond Clustering

- Mixture models is a general framework for modeling grouped data
- Not limited only to clustering or density estimation
- **Mixture of Experts:** Each “expert” is a probabilistic supervised learning model $p(y|\mathbf{x}, \theta_k)$
 - Overall model is a convex combination of the experts

$$p(y|\mathbf{x}) = \sum_{k=1}^K \pi_k(\mathbf{x}) p(y|\mathbf{x}, \theta_k)$$

- Enables learning rich models (e.g., nonlinear reg.) from simpler models (e.g., linear reg.)



Next Class: The General EM Algorithm