

Analysis 1 Homework 9

A. Calculate each of the following limits, or explain why it does not converge. You may use any theorems that were stated in lecture.

i) $\lim_{x \rightarrow 1^+} |x-2|/(x-2)$

ii) $\lim_{x \rightarrow 1^+} |x-1|$

iii) $\lim_{x \rightarrow 1^+} |x-1|/(x-1)^2$

i) $\lim_{x \rightarrow 1^+} |x-2|/(x-2) = \left[\begin{array}{l} x-2 < 0 \\ x \rightarrow 1^+ \\ |x-2| = -x+2 \end{array} \right] \lim_{x \rightarrow 1^+} \frac{-x+2}{x-2} = \lim_{x \rightarrow 1^+} \frac{-(x-2)}{x-2} = -1$

ii) $\lim_{x \rightarrow 1^+} |x-1| = \left[\begin{array}{l} x \rightarrow 1^+ \\ x-1 > 0 \\ |x-1| = x-1 \end{array} \right] = \lim_{x \rightarrow 1^+} x-1 = 0$

iii) $\lim_{x \rightarrow 1^+} |x-1|/(x-1)^2 = \lim_{x \rightarrow 1^+} \frac{x-1}{(x-1)^2} = \lim_{x \rightarrow 1^+} \frac{1}{x-1} = \frac{1}{0^+} = +\infty$

B. i) Prove directly from the definitions that if $g: \mathbb{R} \rightarrow [1, \infty)$ is a function so that $\lim_{x \rightarrow c} g(x) = L$, then $\lim_{x \rightarrow c} 1/g(x) = 1/L$

ii) Prove the same result of the previous part, using Relating Sequences to Functions.

i) $g: \mathbb{R} \rightarrow [1, \infty)$

$\lim_{x \rightarrow c} g(x) = L$

$\lim_{x \rightarrow c} \frac{1}{g(x)} = \frac{1}{L}$

$\forall \epsilon > 0 \exists \delta > 0 \forall x \neq c \quad |x-c| < \delta \Rightarrow |g(x) - L| < \epsilon$

Let $\epsilon > 0 \exists \delta = \delta_1$ and let $x \neq c, |x-c| < \delta$

$\Rightarrow \left| \frac{1}{g(x)} - \frac{1}{L} \right| = \frac{|L - g(x)|}{g(x) \cdot L} < \frac{\epsilon}{1 \cdot 1} = \epsilon$

ii) $\lim_{x \rightarrow c} g(x) = L$

$\Leftrightarrow \lim_{n \rightarrow \infty} g(x_n) = L \quad \lim_{n \rightarrow \infty} x_n = c$

$\lim_{x \rightarrow c} \frac{1}{g(x)} = \frac{1}{L} \quad \forall x_n \rightarrow c \Rightarrow \lim_{n \rightarrow \infty} \frac{1}{g(x_n)} = \frac{1}{L}$

$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{g(x_n)} = \lim_{n \rightarrow \infty} \frac{1}{L} = \frac{1}{L}$

C. Prove that $\lim_{x \rightarrow c} f(x) = \infty$ if and only if for every sequence a_n that is in the domain of f and converges to c (but never equal to c), we have $\lim_{n \rightarrow \infty} f(a_n) = \infty$. (That is, prove a version of relating sequences to functions for infinite limits. Make sure you handle both directions of the if and only if!)

(\Rightarrow) Let $\lim_{x \rightarrow c} f(x) = \infty \Leftrightarrow \forall N > 0 \exists \delta > 0 \forall x \neq c \quad |x - c| < \delta \Rightarrow f(x) > N$ and let (a_n) , $a_n \in D_f$ and $\lim_{n \rightarrow \infty} a_n = c \Leftrightarrow \forall \epsilon > 0 \exists n_0 \forall n > n_0 \quad |a_n - c| < \epsilon$

$$\lim_{n \rightarrow \infty} f(a_n) = \infty \quad \forall N > 0 \exists n_0 \forall n > n_0$$

$$\text{Let } N > 0 \quad \exists n_0 \text{ and let } n > n_0 \quad (|a_n - c| < \epsilon) \quad x = a_n \quad |x - c| < \delta \Rightarrow f(a_n) = f(x) > N$$

(\Leftarrow) If $\forall n \quad \lim_{n \rightarrow \infty} a_n = c \quad \lim_{n \rightarrow \infty} f(a_n) = \infty \Rightarrow \lim_{x \rightarrow c} f(x) = \infty$

Let $\forall n \quad \lim_{n \rightarrow \infty} a_n = c$, it holds $\lim_{n \rightarrow \infty} f(a_n) = \infty$, and if we assume that $\lim_{x \rightarrow c} f(x) \neq \infty$

$$\neg (\forall N > 0 \exists \delta > 0 \forall x \neq c \quad |x - c| < \delta \Rightarrow f(x) > N)$$

$$\exists N > 0 \forall \delta > 0 \exists x \neq c \quad |x - c| < \delta \wedge f(x) \leq N$$

$$\exists N > 0 \forall \delta = \frac{1}{N} \exists x_n = a_n \neq c \quad |a_n - c| < \delta \wedge f(a_n) \leq N$$

This is in contradiction with $\lim_{n \rightarrow \infty} f(a_n) = \infty$

D. Which of the following families of real functions are algebras of functions?
(For each, either show it is an algebra of functions, or else identify a necessary property that is not satisfied)

i) Polynomials of degree at most 2. (That is polynomials of the form ax^2+bx+c , where any or all of a, b, c may be 0.

ii) Functions f such that $f(x) \leq 6$

iii) Functions f that are everywhere defined, and that are continuous at the point $c=2$.

i) f_1 and $f_2 \in \mathcal{P}_2 = \{f \mid f \text{ is polynomial with degree at most } 2\}$

$$\mathcal{P}_2 = \{f \mid f = ax^2 + bx + c, a, b, c \in \mathbb{R}\}$$

$$f_1(x), f_2(x) \in \mathcal{P}_2$$

$$\Rightarrow f_1(x) = a_1x^2 + b_1x + c_1 \quad f_2(x) = \text{~~polynomial~~} \quad a_2x^2 + b_2x + c_2$$

$$\begin{aligned} \Rightarrow (f_1 + f_2)(x) &= f_1(x) + f_2(x) = a_1x^2 + b_1x + c_1 + a_2x^2 + b_2x + c_2 = \\ &= \underbrace{(a_1 + a_2)}_ax^2 + \underbrace{(b_1 + b_2)}_bx + \underbrace{(c_1 + c_2)}_c = \\ &= ax^2 + bx + c \in \mathcal{P}_2 \end{aligned}$$

$$\text{ii) } \mathcal{P}_6 = \{f \mid f(x) \leq 6\}$$

$$\text{If } f_1(x), f_2(x) \in \mathcal{P}_6$$

$$f_1(x) + f_2(x) \in \mathcal{P}_6$$

Doesn't hold, because:

$$\text{for example } f_1(x) = 5 \leq 6 \Rightarrow f_1 \in \mathcal{P}_6$$

$$f_2(x) = 5 \leq 6 \Rightarrow f_2 \in \mathcal{P}_6$$

$$\text{but } (f_1 + f_2)(x) = f_1(x) + f_2(x) = 5 + 5 = 10 > 6$$

$$\Rightarrow f_1 + f_2 \notin \mathcal{P}_6$$

$$\text{iii) } \mathcal{N}_c = \{f \mid f: \mathbb{R} \rightarrow \mathbb{R} \text{ and } f \text{ is continuous at } c\}$$

$$f_1, f_2 \in \mathcal{N}_c$$

$$\Rightarrow (f_1 + f_2)(x) = f_1(x) + f_2(x)$$

$$f_1 + f_2: \mathbb{R} \rightarrow \mathbb{R}$$

E. Prove that $f(x) = x \cdot |x|$ is continuous at all points c in \mathbb{R} .

$$\lim_{x \rightarrow x_0} f(x) = f(x_0)$$

$$\lim_{x \rightarrow c} f(x) = f(c)$$

$$\lim_{x \rightarrow c} x \cdot |x| = \lim_{x \rightarrow c} x \cdot x = \lim_{x \rightarrow c} x^2 = c^2 = c \cdot c = c \cdot |c| = f(c)$$

$$\text{def: } \forall \epsilon > 0 \quad \exists \delta > 0 \quad \forall x \quad |x - x_0| < \delta \Rightarrow |f(x) - f(x_0)| < \epsilon$$

$$\epsilon > 0 \quad \exists \delta > 0 \quad \forall x \quad |x - x_0| < \delta$$

$$\lim_{x \rightarrow c} x|x| = \lim_{x \rightarrow c} x(-x) = -\lim_{x \rightarrow c} x^2 = -c^2 = c(-c) = c \cdot |c| = f(c)$$

$$\Rightarrow |f(x) - f(x_0)| = |x - x_0| < \delta = \epsilon$$

$$\Rightarrow f(x) = x$$

$$\lim_{x \rightarrow 0^-} x|x| = \lim_{x \rightarrow 0^-} x(-x) = -\lim_{x \rightarrow 0^-} x^2 = -0^2 = 0 = c \cdot |c| = f(c) \quad \text{for } c=0$$

$$\lim_{x \rightarrow 0^+} x|x| = \lim_{x \rightarrow 0^+} x \cdot x = \lim_{x \rightarrow 0^+} x^2 = 0^2 = 0 = c \cdot |c| = f(c) = f(0)$$