Spatial Generalized Linear Mixed Models with Application to Prevalence Mapping

空间广义线性混合模型及其在预测流行病中的应用 2015 级硕士学位论文答辩

学生: 黄湘云

导师: 李再兴 专业: 统计学

方向:数据分析与统计计算

理学院 计算数学与统计系

2017年11月19日



- ① 引言 研究意义 文献综述 主要内容
- ② 模型 (SGLMM) 模型结构 计算方法 数据分析
- 3 结论与展望

例例例例例

- 1 radionuclide concentrations on Rongelap Island
- 2 childhood malaria in the gambia
- 3 Loa loa prevalence in Cameroon and surrounding areas

Introduction

000


Diggle et al. (2002)


- First item in the list
- · Second item
- and so on
 - · First item in the list
 - Second item
 - and so on
- the effects of child level covariates (age and bed net use)
- village level covariates (the primary health care and greenness of surrounding vegetation)
- separate components for residual spatial
- non-spatial extrabinomial variation


 \mathbb{R}^n


$$\log\{p_{ij}/(1-p_{ij})\} = \alpha + \beta' z_{ij} + U_i + S(x_i)$$











References I

Diggle, Peter, Moyeed, Rana, Rowlingson, Barry, & Thomson, Madeleine. 2002. Childhood malaria in the Gambia: a case-study in model-based geostatistics. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **51**(4), 493–506.

Softwares and Tools



图: GNU R INLA Stan PyMC3

11 / 12

Github



