
Data analysis for digestate experiments


Sasha D. Hafner

14 October, 2022

Plots

```
ggplot(dat, aes(exper, e.rel, colour = treat)) +
  geom_jitter(height = 0)
  25 -
  20 -
                                                                              treat
                                                                                CC-CM
                                                                                   D-CM
                                                                                   U-CM
  15 -
  10-
                   Experiment 1
                                                  Experiment 2
                                     exper
ggplot(dat, aes(as.integer(factor(exper)), e.rel, colour = treat, group = interaction(treat, rep))) +
  geom_point() +
  geom_line()
```





Stats

Set reference to untreated cattle manure.


```
dat$treat <- factor(dat$treat, levels = c('U-CM', 'CC-CM', 'D-CM'))</pre>
m1 <- aov(e.rel ~ treat + exper, data = dat)</pre>
summary(m1)
##
               Df Sum Sq Mean Sq F value
## treat
                2 83.07
                           41.53
                                 4.266 0.035753 *
## exper
                1 208.56 208.56 21.421 0.000391 ***
               14 136.31
## Residuals
                            9.74
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
summary.lm(m1)
##
## Call:
## aov(formula = e.rel ~ treat + exper, data = dat)
## Residuals:
                1Q Median
##
                                3Q
## -4.6939 -2.1599 -0.2792 1.6681 5.6906
##
## Coefficients:
                     Estimate Std. Error t value Pr(>|t|)
##
```


```
## (Intercept)
                                           9.393 2.02e-07 ***
                       13.816
                                   1.471
## treatCC-CM
                       -2.295
                                          -1.274 0.223428
                                   1.801
## treatD-CM
                        2.953
                                   1.801
                                           1.639 0.123404
## experExperiment 2
                        6.808
                                   1.471
                                           4.628 0.000391 ***
                  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## Signif. codes:
## Residual standard error: 3.12 on 14 degrees of freedom
## Multiple R-squared: 0.6815, Adjusted R-squared: 0.6132
## F-statistic: 9.984 on 3 and 14 DF, p-value: 0.0008875
```

CC-CM and D-CM are clearly different. But neither is clearly different from the reference. Makes interpretation just a bit tricky. Some evidence of a difference but presumably digestion pH effect is moderated by low DM, and variability was high, so power is low.


Check residuals.

plot(m1, ask = FALSE)





Constant Leverage: Residuals vs Factor Levels



Factor Level Combinations

Looks fine, no need to transform.