

Comparison of abmneo and ABM - ABM part


Sasha D. Hafner

30 March, 2026 09:20


```
devtools::load_all('../..//ABM')
```

```
## i Loading ABM
```

```
mass_series <- read.csv('level_pig_od_ref.csv')  
plot(slurry_mass ~ time, data = mass_series, type = 'o')
```



```
temp_dat <- data.frame(time = seq(0, 365, length.out = 12))  
temp_dat$temp_C <- 10 + 10 * sin((temp_dat$time - 130) / 365 * 2 * pi)  
plot(temp_C ~ time, data = temp_dat, type = 'o')
```



```

arrh_pars <- list(lnA = 29.96298,
  E_CH4 = 81000,
  A = c(xa_dead= 3.61383 * 10^12, starch = 5 * 3.61383 * 10^12, Cfat = 0.1 * 3.61383 * 10^12),
  E = c(xa_dead= 81557, starch = 81557, Cfat = 81557, CPs = 81557, CPf = 81557, RFd = 81557),
  R = 8.314,
  VS_CH4 = 6.67,
  scale_alpha_opt = list(VSd = 0.46056, notVSd = 0.556886, CP = 2.065),
  kl = c(NH3 = 336, NH3_floor = 0.36*337, H2S = 0.02),
  scale_EF_NH3 = 0.0000048)

grp_pars <- list(grps = c('m0', 'm1', 'm2','m3','m4','m5','sr1'),
  yield = c(default = 0.05, sr1 = 0.065),
  xa_fresh = c(m0 = 0.06, m1 = 0.06, m2 = 0.06, m3 = 0.06, m4 = 0.06, m5 = 0.06, sr1 = 0),
  xa_init = c(all = 0.06),
  decay_rate = c(all = 0.02),
  ks_coefficient = c(default = 1.153337, sr1 = 0.461335),
  qhat_opt = c(m0 = 0.46289, m1 = 0.74568, m2 = 0.9006605, m3 = 2.79672, m4 = 4.66012, m5 = 55, sr1 = 43.75),
  T_opt = c(m0 = 18, m1 = 18, m2 = 28, m3 = 36, m4 = 43.75, m5 = 55, sr1 = 43.75),
  T_min = c(m0 = 0, m1 = 9.67, m2 = 9.67, m3 = 15, m4 = 26.25, m5 = 30, sr1 = 0),
  T_max = c(m0 = 25, m1 = 25, m2 = 38, m3 = 45, m4 = 51.25, m5 = 60, sr1 = 51.25),
  ki_NH3_min = c(all = 0.015),
  ki_NH3_max = c(all = 5.37),
  ki_NH4_min = c(all = 2.7),
  ki_NH4_max = c(all = 6.8492),
  ki_H2S_slope = c(default = -0.10623, sr1 = -0.1495),
  ki_H2S_int = c(default = 1.02, sr1 = 1.42),

```

```

ki_H2S_min = c(default = 0.08),
IC50_low = c(default = 0.20854, sr1 = 0.2772),
ki_HAC = c(default = 0.31),
pH_UL = c(default = 8.0),
pH_LL = c(default = 6.5, sr1 = 5.5))

```

Write mng_pars in order to insert slurry_mass data

```

mng_pars <- list(slurry_prod_rate = 5700,
  slurry_mass = mass_series,
  storage_depth = 2,
  resid_depth = 0,
  floor_area = 0,
  area = 0.7,
  empty_int = 0,
  temp_C = temp_dat,
  wash_water = 0,
  wash_int = NA,
  rest_d = 0,
  cover = 'none',
  resid_enrich = 0.9,
  graze = c(start = 'May', duration = 0, hours_day = 0),
  scale = c(ks_coefficient = 1, qhat_opt = 1, xa_fresh = 1, yield = 1, alpha_opt = 1)
)

```

Take man_pars from man_pars_pig2.0

```

man_pars_pig2.0 <- list(conc_fresh = list(sulfide = 0.01, urea = 3.17, sulfate = 0.01, TAN = 0.0, starch = 0.0,
  VFA = 1.7, xa_dead = 0, Cfat = 27.6, CPs = 21.1 * 0.55, CPf = 21.1,
  ash = 15), pH = 7, dens = 1000)

```

```

devtools::load_all('../..//ABM')

```

i Loading ABM

```

system.time(
  abm_out <- abm(
    man_pars = man_pars_pig2.0,
    mng_pars = mng_pars,
    arrh_pars = arrh_pars,
    grp_pars = grp_pars,
    evap_pars = list(evap = 0, rain = 0),
    startup = 2
  )
)

```

##

Startup run 1x ->

```

## Warning in abm_variable(days = days, delta_t = delta_t, times = times, y = y, : Negative slurry mass
## Check parameters and try again.

```

```

## Warning in emptyStore(y, resid_mass = slurry_mass[j], resid_enrich =
## pars$resid_enrich): Emptying skipped because of low slurry level.
## Warning in emptyStore(y, resid_mass = slurry_mass[j], resid_enrich =
## pars$resid_enrich): Emptying skipped because of low slurry level.
## Warning in emptyStore(y, resid_mass = slurry_mass[j], resid_enrich =


```




```
## arguments overwritten by slurry_mass: slurry_prod_rate, empty_int, resid_depth, wash_water, wash_int
## rain = 1.9 kg/m2/day
## evaporation = 0 kg/m2/day

## user system elapsed
## 0.538 0.001 0.538


plot(slurry_mass ~ time, data = abm_out, type = 'l')
lines(slurry_mass ~ time, data = mass_series, type = 'l', col = 'red')
```




```
plot(temp_C ~ time, data = abm_out, type = 'l')
```



```
plot(CH4_emis_rate ~ time, data = abm_out, type = 'l')
```



```
matplot(abm_out$time, abm_out[, nn <- c('m0', 'm1', 'm2', 'm3', 'm4', 'm5', 'sr1')], col = cc <- c('blue'
legend('topright', legend = nn, col = cc, lty = 1:length(nn))
```




```
matplot(abm_out$time, abm_out[, nn <- paste0(c('m0','m1','m2', 'm3', 'm4', 'm5', 'sr1'), '_conc')], col =
legend('topright', legend = nn, col = cc, lty = 1:length(nn))
```



```

      .jt[, nn <- paste0(c("m0", "m1", "m2", "m3", "m4", "m5", "sr1"),


```




```

matplot(abm_out$time, abm_out[, nn <- c('starch', 'Cfat', 'CPs', 'CPf', 'RFd')], col = cc <- c('blue',


```




```
matplot(abm_out$time, abm_out[, nn <- paste0(c('starch', 'Cfat', 'CPs', 'CPf', 'RFd'), '_conc')], col =
legend('topright', legend = nn, col = cc, lty = 1:length(nn))
```



```
plot(VFA ~ time, data = abm_out, type = 'l')
```



```
plot(starch_conc ~ time, data = abm_out, type = 'l')
```



```
plot(VFA_conc ~ time, data = abm_out, type = 'l')
```

