MC 501 Inferencing Data in R rubric

100 points. Due Monday, December 7, 11:59 PM.

How this is scored. Add the points from each row; the total is out of 100. A first attempt is graded forgivingly: a part that is there but incomplete earns partial credit, and 0 is kept for a part that is missing. Attempts are unlimited, so you can revise an early submission right up to the deadline.

- Run all four tests: a chi-square test of independence with `v2v::run_chi_square()`; an independent t-test with `v2v::run_t_test()` (Welch's version); a one-way ANOVA with `aov()` and `summary()`, then `TukeyHSD()`; and a linear regression with `lm()` and `summary()`, with `lm.beta()` for the standardized coefficient.
- The effectsize package returns each interval: `cramers_v()` for the chi-square, `cohens_d()` for the t-test, `omega_squared()` and `eta_squared()` on the ANOVA, and `eta_squared()` on the regression model, which for one predictor is R-squared. Install it with install.packages("effectsize") if the setup guide's optional step did not.
- Every script opens with a settings block: load the data, rename the variables you use to outcome, `group2`, `group3`, predictor, `cat_a`, and `cat_b`, and make the categorical ones factors. The code below the block uses only those names. This is what lets the same script run on your own data for the White Paper.
- How this loses points: going wide instead of deep. Four tests each reported with their assumptions, effect size, and interval score above four tests reported as four bare sentences.
- What it feeds: your White Paper, which runs this same script on your own variables after you change its settings block. The Twitch results stay here; the paper reports only what your own data shows, and only the tests your questions call for.

Inferencing Data in R, 100 points

Criterion	Exemplary	Proficient	Developing	Beginning
Hypotheses 8 points	**8 points.** Four hypotheses in plain English, written before any code, one for each test (a chi-square test of independence, an independent t-test, a one-way ANOVA across three or more groups, and a linear regression with a continuous outcome), each naming its variables and the relationship you expect.	**6 to 7 points.** Four stated before the code, but one variable is vague, or one hypothesis is a topic instead of a relationship.	**4 to 5 points.** Fewer than four, or stated after the results.	**0 to 3 points.** Missing.
Test choice against a rejected alternative 12 points	**12 points.** Each of the four tests justified against one alternative you rejected and why.	**9 to 11 points.** The right four tests, but a rejected alternative or its reason is missing on one.	**6 to 8 points.** A test that does not fit its hypothesis's variable types.	**0 to 5 points.** No tests chosen.

Criterion	Exemplary	Proficient	Developing	Beginning
Assumptions and diagnostics 20 points	**20 points.** Each test's assumptions checked with the diagnostic shown rather than asserted (expected cell counts for the chi-square, the two groups' spread for the t-test, equal variances across groups for the ANOVA, with Welch's ANOVA from `oneway.test()` where they differ, and the residual plots from plot(model) for the regression), and what you did about any that fail.	**15 to 19 points.** Checked and shown, but nothing said about one that fails.	**10 to 14 points.** Assumptions listed or asserted ("the data are normal") without being shown, or checked for some tests only.	**0 to 9 points.** Missing.
APA reporting 12 points	**12 points.** All four tests run in your document and are reported in APA format with the statistic, degrees of freedom, p, and n; for the ANOVA, Tukey's HSD names which groups differ.	**9 to 11 points.** One of the four pieces is missing on a test, or the Tukey step is missing.	**6 to 8 points.** The statistic and p only, or numbers pasted in instead of produced by the code.	**0 to 5 points.** Not reported.

Criterion	Exemplary	Proficient	Developing	Beginning
Effect sizes with their intervals 20 points	**20 points.** Each effect size with its 95% confidence interval, in the units of the data as well as by its conventional label: Cramér's V; Cohen's d; omega-squared with eta-squared beside it; and for the regression R-squared, the slope in the outcome's units, and the standardized coefficient.	**15 to 19 points.** All four with intervals, but one carries the label only, or eta-squared or the standardized coefficient is missing.	**10 to 14 points.** Computed without intervals, or the wrong measure for a test.	**0 to 9 points.** Missing.
Power for one focal test 10 points	**10 points.** For one test of your choice: what effect this sample could detect, and what that means for a null result.	**8 to 9 points.** The detectable effect, without what it means for a null result.	**5 to 7 points.** A power figure with no link to your own design (a generic eighty percent).	**0 to 4 points.** Missing.
Significance against size 10 points	**10 points.** A paragraph that would survive a reviewer, separating statistical significance from practical size across all four results.	**8 to 9 points.** Separates the two, but for some of the results only.	**5 to 7 points.** Treats a small p as a large effect.	**0 to 4 points.** Missing.
What the results do not license 8 points	**8 points.** One paragraph, including the gap between the number of messages and the number of channels behind any channel-level variable.	**6 to 7 points.** A limit, but the gap between messages and channels is not named.	**4 to 5 points.** A generic limit ("more research is needed").	**0 to 3 points.** Missing.

Before you submit

Inferential Check recomputes a chi-square or a t test from your own table or group figures and returns the statistic, the degrees of freedom, p, the effect size, the assumptions your data may be breaking, and the APA sentence to compare against what you wrote. It is a second opinion on your R output, not a replacement for it. It does not grade, and nothing you enter leaves your machine. It does not cover ANOVA or regression, so check those two against the worked examples in Chapter 13.