



MAARULA CLASSES

FOR DETAILED VISIT



www.questions.maarula.in



0512-3163515



https://youtube.com/@maarulaclasses?si=XkKXyJ_2a16NiK-K



9554548576

MAARULAMATHEM E-LEARNING APP

SOLVED WITH TRICK

MAH MCA CET
2026



SLOT-02



By Amit Katiyar
(MCA-JNU)

1. Topic Wise Weightage & Detailed Trend Analysis.
2. All 17 Years Paper With Smart Solution.
3. Syllabus & Important Documents.
4. Topic Wise All PYQ's NIMCET-Free Test.



MAARULA CLASSES

TARGET- NIMCET / CUET.PG By: Amit Katiyar (MCA-JNU)

MAH MCA CET-2026

Slot-02



Scan the QR & Download Our App Now.

1. If $f(x)$, $g(x)$ be twice differentiable functions on $[0,2]$ satisfying $f''(x) = g''(x)$, $f'(1) = 2$, $g'(1) = 4$ and $f(2) = 3$, $g(2) = 9$, then $f(x) - g(x)$ at $x = 4$ equals
(a) 0 (b) -10
(c) 8 (d) 2
2. The order of differential equation of the family of circles with one diameter along the line $y = x$ is
(a) 1 (b) 2
(c) 3 (d) 0
3. If $x = \frac{e^{2x} + 1}{e^{2x} - 1}$ then $\frac{dy}{dx}$ is
(a) $\frac{1}{1+x^2}$ (b) $\frac{1}{x^2-1}$
(c) $\frac{1}{1-x^2}$ (d) $\frac{1}{x^2-1}$
4. Two measurements of a cylinder are varying in such a way that the volume is kept constant. if the rate of change of radius (r) and height (h) are equal in magnitude but opposite in sign then
(a) $r = 2h$ (b) $h = 2r$
(c) $h = r$ (d) $h = 4r$
5. Ratio of two numbers X and Y is 3:2. If decrease 60 from X and add 60 to Y , then the ratio of X and Y is 18:17. The original value of X is
(a) 440 (b) 460
(c) 450 (d) 420
6. The coefficients of variation of two distributions are 75 and 80, and their standard deviations are 15 and 16 respectively. Their arithmetic mean respectively, are
(a) 5 and 5. (b) 5 and 6.
(c) 20 and 20. (d) 500 and 500.
7. Let $A = \{-3, -2, -1, 0, 1, 2, 3\}$. Let be a relation on A defined by $x R y$ if and only if $0 \leq x^2 + 2y \leq 4$. Let l be the number of elements in R and m be the minimum number of elements required to be added in R to make it a reflexive relation. Then $l + m$ is equal to
(a) 17 (b) 18
(c) 19 (d) 20
8. Let $S = N \cup \{0\}$. Define a relation R from S to Q .
 $Q = \{(x, y): \log_e y = x \log_e \left(\frac{2}{5}\right), x \in S, y \in Q\}$. Then the sum of elements in the range R is equal to
(a) $\frac{3}{2}$ (b) $\frac{9}{2}$
(c) $\frac{10}{9}$ (d) $\frac{5}{3}$
9. Two finite sets have m and n elements respectively. The total number of subsets of first set is 120 more than the total number of subsets of the second set. The values of m and n respectively are:
(a) 4, 3 (b) 7, 3
(c) 6, 3 (d) 3, 7
10. Side of square is three less than the base of the triangle. Height of triangle is 24 m. Area of triangle is 180 m^2 . The area of the square in m^2 is
(a) 144 (b) 256
(c) 196 (d) 324
11. The radius of a cylinder, a cone and a sphere are in the ratio 1:2:3 and the ratio of their heights is 5:6:7. If the volume of the sphere is 4851 cm^3 , the the sum of volumes of cylinder and cone (in cm^3) is
(a) 672.5 (b) 1501.5
(c) 771.5 (d) 671.5
12. One of the roots of the equation $8x^3 - 6x + 1$ is
(a) $\cos 10^\circ$ (b) $\cos 30^\circ$
(c) $\sin 30^\circ$ (d) $\cos 80^\circ$
13. The equation $a \sin x + \cos 2x = 2a - 7$ possesses a solution if
(a) $a > 6$ (b) $2 \leq a \leq 6$
(c) $a < 2$ (d) $a = 7$
14. The sum of all natural numbers between 250 and 1000, which are exactly divisible by 3 is
(a) 156375 (b) 155375
(c) 156575 (d) 156535
15. If A and B are arbitrary events, then
(a) $P(A \cap B) \geq P(A) + P(B) - 1$
(b) $P(A \cap B) \leq P(A) + P(B) - 1$
(c) $P(A \cap B) = P(A) + P(B) - 1$
(d) Insufficient information

ADDRESS: 117/466, O Block, Geeta Nagar, Sharda Nagar, NEAR: ANURAG HOSPITAL Kanpur, Uttar Pradesh 208025

website: www.question.maarula.in

0512-3163515

9935985550

9554548576



16. If the equation $x^4 - 4x^3 + ax^2 + bx + 1 = 0$ has four positive roots then (a, b) is
 (a) (-4, 6) (b) (6, -4)
 (c) (-4, -6) (d) (4, -6)
17. Let $a_1, a_2, a_3, \dots$ be in arithmetic progression and $q_1, q_2, q_3, \dots$ be in geometric progression, such that $a_1 = q_1 = 2$ and $a_{10} = q_{10} = 3$ then:
 (a) $a_7 a_{19}$ is not an integer
 (b) $a_{19} q_7$ is an integer
 (c) $a_7 a_{19} = a_{19} q_{10}$
 (d) Cannot be worked out from the given information
18. If the roots of $px^2 + qx + 2 = 0$ are reciprocal of each other then
 (a) $p = 0$ (b) $p = -2$
 (c) $p = \pm 2$ (d) $p = 2$
19. The solution of the equation is $2^{\frac{3}{\log_{3x}}} = \frac{1}{64}$ is
 (a) 3 (b) $\frac{1}{3}$
 (c) $\frac{1}{\sqrt{3}}$ (d) $\frac{1}{3^2}$
20. In the binomial expansion of $(a - b)^n, n \leq 5$, the sum of 5th and 6th term is zero. Then $\frac{a}{b}$ equals
 (a) $\frac{n-5}{6}$ (b) $\frac{n-4}{5}$
 (c) $\frac{5}{n-4}$ (d) $\frac{6}{n-5}$
21. The value of $\sum_{r=1}^n \frac{n P_r}{r!}$ is
 (a) 2^n (b) $2^n - 1$
 (c) 2^{n-1} (d) $2^n + 1$
22. The locus of the middle points of focal chords of the parabola $y^2 = 4ax$ is
 (a) $y^2 = 2a(x - a)$
 (b) $y^2 = 2a(x + a)$
 (c) $x^2 = 2a(y - a)$
 (d) $x^2 = 2a(y + a)$
23. The triangle formed by the equations $x + y = 0, 3x + y - 4 = 0, x + 3y - 4 = 0$
 (a) equilateral
 (b) isosceles
 (c) scalene
 (d) cannot be worked out
24. The equations to straight lines passing through the points (4, 5) and equally inclined to the lines $3x = 4y + 7$ and $5y = 12x + 6$ are
 (a) $9x - 7y = 1$ and $7x + 9y = 73$
 (b) $7x - 9y = 1$ and $7x + 9y = 73$
 (c) $9x - 7y = 1$ and $9x + 7y = 73$
 (d) $7x - 7y = 1$ and $9x + 9y = 73$
25. The triangle PQR is inscribed in the circle $x^2 + y^2 = 25$. If Q and R have coordinates (3, 4) and (-4, 3) respectively, then $\angle QPR$ is equal to
 (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{3}$
 (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{6}$
26. A certain distance is covered by a car at a certain speed. If scooter covers half the distance in double the time, the ratio of the speed of the scooter to that of the car is:
 (a) 1:2 (b) 2:1
 (c) 4:1 (d) 1:4
27. In how many ways can a committee of 3 members be formed from 7 persons?
 (a) 21 (b) 35
 (c) 70 (d) 210
28. If $\log_2 X = 5$ then the value of X is:
 (a) 10 (b) 16
 (c) 32 (d) 64
29. Two coins are tossed simultaneously. What is the probability of getting at least one head?
 (a) 1/4 (b) 1/2
 (c) 3/4 (d) 1/3
30. Two series A and B have the same mean. The standard deviation of A is 5 and that of B is 10. Which statement is correct?
 (a) Series A is more variable
 (b) Series B is more consistent
 (c) Series A is more consistent
 (d) Both Series have equal variability
31. Excluding stoppages, the speed of a bus is 54 kmph and including stoppages, it is 45 kmph. For how many minutes does the bus stop per hour?
 (a) 9 (b) 10
 (c) 15 (d) 12





MAARULA CLASSES

TARGET- NIMCET / CUET.PG By: Amit Katiyar (MCA-JNU)

MAH MCA CET-2026


Slot-02



Scan the QR & Download Our App Now.

ANSWER KEY

1.	2.	3.	4.	5.
b	b	c	a	d
6.	7.	8.	9.	10.
c	b	d	b	a
11.	12.	13.	14.	15.
b	d	b	a	a
16.	17.	18.	19.	20.
b	c	d	c	b
21.	22.	23.	24.	25.
b	a	b	a	c
26.	27.	28.	29.	30.
d	b	c	c	c
31.				
b				



ADDRESS: 117/466, O Block, Geeta Nagar, Sharda Nagar, NEAR: ANURAG HOSPITAL Kanpur, Uttar Pradesh 208025

website: www.question.maarula.in



0512-3163515



9935985550



9554548576