

Auteur: Noran Ben Said
Studierichting: Informatica
Oefeningenreeks 1C nr. 48.8

$$z^6 = -\sqrt{3} - i$$

$$r = \sqrt{(-\sqrt{3})^2 + (-1)^2} = 2$$

$$\theta = \frac{7\pi}{6}$$

$$-\sqrt{3} - i = 2 \operatorname{cis} \frac{7\pi}{6}$$

$$z_k = \sqrt[6]{2} \operatorname{cis} \left(\frac{\frac{7\pi}{6} + k2\pi}{6} \right)$$

$$= \sqrt[6]{2} \operatorname{cis} \left(\frac{7\pi}{36} + k\frac{\pi}{3} \right)$$

$$z_0 = \sqrt[6]{2} \operatorname{cis} \frac{7\pi}{36}$$

$$z_1 = \sqrt[6]{2} \operatorname{cis} \frac{19\pi}{36}$$

$$z_2 = \sqrt[6]{2} \operatorname{cis} \frac{31\pi}{36}$$

$$z_3 = \sqrt[6]{2} \operatorname{cis} \frac{43\pi}{36}$$

$$z_4 = \sqrt[6]{2} \operatorname{cis} \frac{55\pi}{36}$$

$$z_5 = \sqrt[6]{2} \operatorname{cis} \frac{67\pi}{36}$$

References