

Auteur: Abdul Ahmed
Studierichting: Informatica
Oefeningenreeks 2E nr. 26.12

$$\begin{aligned}\log_x \left(\frac{\log_2 6}{\log_{12} 6} + \log_2 \frac{1}{3} \right) &= \log_2 \left((2x)^{\frac{1}{6}} \right) \\ \Leftrightarrow \log_x \left(\frac{(\log_2 6) \cdot (\log_2 12)}{\log_2 6} + \log_2 \frac{1}{3} \right) &= \log_2 \left((2x)^{\frac{1}{6}} \right) \\ \Leftrightarrow \log_x \left(\log_2 12 + \log_2 \frac{1}{3} \right) &= \log_2 \left((2x)^{\frac{1}{6}} \right) \\ \Leftrightarrow \log_x (\log_2 4) &= \log_2 \left((2x)^{\frac{1}{6}} \right) \\ \Leftrightarrow \log_x 2 &= \frac{1}{6} \log_2 (2x) \\ \Leftrightarrow 6 \log_x 2 &= \log_2 2x \\ \Leftrightarrow 6 \frac{\log_2 2}{\log_2 x} &= 1 + \log_2 x \\ \Leftrightarrow \frac{6}{\log_2 x} &= 1 + \log_2 x\end{aligned}$$

Stel $t = \log_2 x$

$$\begin{aligned}\Rightarrow \frac{6}{t} &= 1 + t \\ \Leftrightarrow t^2 + t - 6 &= 0\end{aligned}$$

De oplossingen voor t zijn dan:

$$\begin{aligned}t_1 &= \frac{-1 + \sqrt{1 + 24}}{2} = -3 \\ t_2 &= \frac{-1 - \sqrt{25}}{2} = 2\end{aligned}$$

De oplossingen voor x zijn dan:

$$\begin{aligned}\log_2 x_1 &= -3 \text{ of } \log_2 x_2 = 2 \\ \Leftrightarrow x_1 &= \frac{1}{8} \text{ of } x_2 = 4\end{aligned}$$

References