

Auteur: Sabawoon Enayat
Studierichting: Informatica
Oefeningenreeks 6A nr. 7.5

$$\begin{aligned} &\text{Opgave: } x_1 \cdot x_3 = 36 \text{ en } x^2 + x^4 = 60 \\ \Rightarrow &\begin{cases} x_1 \cdot (x_1 \cdot q^2) &= 36 \\ x_1 \cdot q + x_1 \cdot q^3 &= 36 \end{cases} \Rightarrow \begin{cases} (x_1 \cdot q)^2 &= 36 \\ x_1(q + q^3) &= 60 \end{cases} \\ \Rightarrow &\begin{cases} x_1 \cdot q &= 6 \\ x_1(q + q^3) &= 60 \end{cases} \Rightarrow \begin{cases} x_1 &= \frac{6}{q} \\ x_1 \cdot (q + q^3) &= 60 \end{cases} \\ \Rightarrow &\frac{6}{q}(q + q^3) = 60 \Rightarrow 6q^2 + 6 = 60 \Rightarrow q^2 + 1 = 10 \Rightarrow q^2 = 9 \\ \Rightarrow &q = \pm 3 \Rightarrow x_1 = \frac{6}{q} = 2 \text{ of } x_1 = -2 \\ \Rightarrow &x_2 = x_1 \cdot q = 6 \\ \Rightarrow &x_3 = x_2 \cdot q = 18 \text{ of } x_3 = x_2 \cdot q = -18 \\ \Rightarrow &(2, 6, 18, \dots) \text{ of } (-2, 6, -18, \dots) \end{aligned}$$

References