

Auteur: Niels Hertoghs
 Studierichting: Informatica
 Oefeningenreeks 7D nr. 1.1

$$f(x, y) = x^3 + x^2y^3 - 3y^4 \text{ in } (1, -1), \text{ graad } 2$$

$$f(1, -1) = -3$$

$$\frac{\partial f}{\partial x}(x, y) = 3x^2 + 2xy^3 \Rightarrow \frac{\partial f}{\partial x}(1, -1) = 1$$

$$\frac{\partial f}{\partial y}(x, y) = 3x^2y^2 - 12y^3 \Rightarrow \frac{\partial f}{\partial y}(1, -1) = 15$$

$$\frac{\partial^2 f}{\partial x^2}(x, y) = 6x + 3y^3 \Rightarrow \frac{\partial^2 f}{\partial x^2}(1, -1) = 4$$

$$\frac{\partial^2 f}{\partial y^2}(x, y) = 6x^2y - 36y^2 \Rightarrow \frac{\partial^2 f}{\partial y^2}(1, -1) = -42$$

$$\frac{\partial^2 f}{\partial x \partial y}(x, y) = \frac{\partial^2 f}{\partial y \partial x}(x, y) = 6xy^2$$

$$\Rightarrow \frac{\partial^2 f}{\partial x \partial y}(1, -1) = \frac{\partial^2 f}{\partial y \partial x}(1, -1) = 6$$

$$T_2(x, y) = f(a, b) + \frac{\partial f}{\partial y}(a, b)(y - b) + \frac{\partial f}{\partial x}(a, b)(x - a)$$

$$+ \frac{1}{2} \left[\frac{\partial^2 f}{\partial x^2}(a, b)(x - a)^2 + \frac{\partial^2 f}{\partial x \partial y}(a, b)(x - a)(y - b) + \frac{\partial^2 f}{\partial y^2}(a, b)(y - b)^2 \right]$$

$$= -14 + 3x - 33y + 2x^2 - 21y^2 + 6xy$$

References