

THE DIFFERENTIAL GEOMETRY DIFFERENTIAL TOPOLOGY DUMP		
ERNEST YEUNG ERNESTYALUMNI@GMAIL.COM		
CONTENTS		
Part 1. Combinatorics, Probability Theory	2	32
Part 2. Linear Algebra Review	2	33
Part 3. Manifolds	3	
1. Inverse Function Theorem	3	
2. Immersions	5	
3. Submersions; Rank Theorem	6	
4. Submanifolds; immersed submanifold, embedded submanifolds, regular submanifolds	8	
5. Curves, Integral Curves, and Flows	8	
6. Tensors	9	
Part 4. Lie Groups, Lie Algebra	11	
7. Lie Groups	11	
Part 5. Cohomology; Stoke’s Theorem	23	
8. Stoke’s Theorem	23	
Part 6. Prástaro	24	
Part 7. Connections	25	
9. Connections of Vector Bundles	25	
Part 8. Holonomy	25	
10. Path Groupoid of a smooth manifold; generalization of paths	28	
Part 9. Complex Manifolds	29	
Part 10. Jets, Jet bundles, h-principle, h-Prinzipien	29	
Part 11. Morse Theory	30	
11. Morse Theory introduction from a physicist	30	
12. Lagrange multipliers	31	
Part 12. Classical Mechanics	31	
13. Classical Mechanics	31	
	14. Fluid Mechanics, Fluid Flow	32
	15. Thermodynamics	33
	Part 13. Classical Mechanics applications	33
	16. Orbital Mechanics	34
	Part 14. General Relativity	37
	Part 15. WE Heraeus International Winter School on Gravity and Light	37
	Introduction (from EY)	37
	17. Lecture 1: Topology	37
	Topology Tutorial Sheet	38
	18. Lecture 2: Topological Manifolds	38
	Tutorial Topological manifolds	39
	19. Lecture 3: Multilinear Algebra	40
	20. Lecture 4: Differentiable Manifolds	42
	Tutorial 4 Differentiable Manifolds	43
	21. Lecture 5: Tangent Spaces	44
	Lecture 6: Fields	46
	Lecture 7: Connections	48
	Lecture 8: Parallel Transport & Curvature (International Winter School on Gravity and Light 2015)	50
	Tutorial 8 Parallel transport & Curvature	52
	Lecture 9: Newtonian spacetime is curved!	53
	22. Lecture 10: Metric Manifolds	55
	23. Symmetry	57
	24. Integration	59
	25. Lecture 13: Relativistic spacetime	61
	26. Lecture 14: <u>Matter</u>	62
	27. Lecture 15: Einstein gravity	64
	Tutorial 13 Schwarzschild Spacetime	65
	28.	66
	29.	66
	30. Canonical Formulation of GR I	66
	31.	66
	32.	66
	33.	66
	34. Lecture 22: <u>Black Holes</u>	66

Date: 28 juillet 2016.
Key words and phrases. Differential Geometry, Differential Topology.

EY : Exercise B.22(d) is useful for showing the chart and atlas of a Grassmannian manifold, found in the More examples, for smooth manifolds.

Proof. (a) Recall the **rank-nullity theorem**:

Theorem 2 (Rank-Nullity Theorem).

$$(5) \qquad \qquad \qquad \dim(im(S)) + \dim(ker(S)) = \dim V$$

Now

$$\begin{aligned} \text{rank}(S) + \dim(\ker(S)) &\equiv \dim(\text{im}(S)) + \dim(\ker(S)) = \dim V \\ \implies \boxed{\text{rank}(S) &\leq \dim V} \end{aligned}$$

If $\text{rank}(S) = \dim V$,

then by rank-nullity theorem, $\dim(\ker(S)) = 0$, implying that $\ker S = \{0\}$.

Suppose $v_1, v_2 \in V$ and that $S(v_1) = S(v_2)$. By linearity of S , $S(v_1) - S(v_2) = S(v_1 - v_2) = 0$, which implies, since $\ker S = \{0\}$, that $v_1 - v_2 = 0$.

$\implies v_1 = v_2$. Then by definition of injectivity, S injective.

If S injective, then $S(v) = 0$ implies $v = 0$. Then $\ker S = \{0\}$. Then by rank-nullity theorem, $\text{rank}(S) = \dim V$.

(b) $\forall w \in \text{im}(S)$, $w \in W$. Clearly $\text{rank } S \leq \dim W$.

If S surjective, $\text{im}(S) = W$. Then $\dim(\text{im}(S)) = \text{rank } S = \dim W$.

If $\text{rank } S = \dim W = m$, then $\text{im}(S)$ has basis $\{y_i\}_{i=1}^m$, $y_i \in \text{im}(S)$, so $\exists x_i \in V$, $i = 1 \dots m$ s.t. $S(x_i) = y_i$, with $\{S(x_i)\}_{i=1}^m$ linearly independent.

Since $\{S(x_i)\}_{i=1}^m$ linearly independent and $\dim W = m$, $\{S(x_i)\}_{i=1}^m$ basis for W .

$\forall w \in W$, $w = \sum_{i=1}^m w^i S(x_i) = S(\sum_{i=1}^m w^i x_i)$. $\sum_{i=1}^m w^i x_i \in V$. S surjective.

(c)

(d) Now

$$\dim V = \text{rank } TS + \text{nullity } TS$$

$$\dim V = \text{rank } S + \text{nullity } S$$

$\ker S \subseteq \ker TS$, clearly, so $\text{nullity } S \leq \text{nullity } TS$

$$\implies \boxed{\text{rank } TS \leq \text{rank } S}$$

If $\text{rank } TS = \text{rank } S$,

then $\text{nullity } S = \text{nullity } TS$

Suppose $w \in \text{Im } S \bigcap \ker T$, $w \neq 0$

Then $\exists v \in S$, s.t. $w = S(v)$ and $T(w) = 0$

Then $T(w) = TS(v) = 0$. So $v \in \ker TS$

$v \notin \ker S$ since $w = S(v) \neq 0$

This implies $\text{nullity } TS > \text{nullity } S$. Contradiction.

$\implies \text{Im } S \bigcap \ker T = 0$

If $\text{Im } S \bigcap \ker T = 0$,

Consider $v \in \ker TS$. Then $TS(v) = 0$.

. Then $S(v) \in \ker T$

$S(v) = 0$; otherwise, $S(v) \in \text{Im } S$, contradicting given $\text{Im } S \bigcap \ker T = 0$

$v \in \ker S$

$\ker TS \subseteq \ker S$

$\implies \ker TS = \ker S$

So $\text{nullity } TS = \text{nullity } S$

$\implies \text{rank } TS = \text{rank } S$

(e)

(f)

(g)

□

Part 3. Manifolds

1. INVERSE FUNCTION THEOREM

Shastri (2011) had a thorough and lucid and explicit explanation of the Inverse Function Theorem [5]. I will recap it here. The following is also a blend of Wienhard's Handout 4 <https://web.math.princeton.edu/~wienhard/teaching/M327/handout4.pdf>

Definition 1. Let (X, a) metric space.

contraction $\phi : X \rightarrow X$ if \exists constant $0 < c < 1$ s.t. $\forall x, y \in X$

$$d(\phi(x), \phi(y)) \leq cd(x, y)$$

Theorem 3 (Contraction Mapping Principle). Let (X, d) complete metric space.

Then \forall contraction $\phi : X \rightarrow X$, $\exists ! y \in X$ s.t. $\phi(y) = y$, y fixed pt.

Proof. Recall def. of complete metric space X , X metric space s.t. \forall Cauchy sequence in X is convergent in X (i.e. has limit in X).

$$x_1 = \phi(x_0)$$

$$x_2 = \phi(x_1)$$

$$\vdots$$

$\forall x_0 \in X$, Define \vdots

$$x_j = \phi(x_{j-1})$$

$$\vdots$$

$$x_n = \phi(x_{n-1})$$

$$d(x_{n+1}, x_n) = d(\phi(x_n), \phi(x_{n-1})) \leq cd(x_n, x_{n-1}) \leq \dots \leq c^n d(x_1, x_0)$$

for some $0 < c < 1$.

$$d(x_m, x_n) \leq d(x_n, x_{n-1}) + d(x_{n-1}, x_m) \leq d(x_n, x_{n-1}) + d(x_{n-1}, x_{n-2}) + \dots + d(x_{m+1}, x_m) \leq \sum_{k=n-1}^m c^k d(x_1, x_0)$$

Thus, $\forall \epsilon > 0$, $\exists n_0 > 0$, (n_0 large enough) s.t. $\forall m, n \in \mathbb{N}$ s.t. $n_0 < n < m$,

$$d(x_m, x_n) \leq \sum_{k=n-1}^m c^k d(x_1, x_0) < \epsilon d(x_1, x_0)$$

Thus, $\{x_n\}$ Cauchy sequence. Since X complete, \exists limit pt. $y \in X$ of $\{x_n\}$.

$$\phi(y) = \phi(\lim_n x_n) = \lim_n \phi(x_n) = \lim_n x_{n+1} = y$$

Since by def. of y limit pt. of $\{x_n\}$, $\forall \epsilon > 0$, then $\{n || x_n - y| \leq \epsilon, n \in \mathbb{N}\}$ is infinite.

Consider $\delta > \mathbb{N}$. Consider $\{n || x_n - y| \leq \delta, n \in \mathbb{N}\}$

$\exists N_\delta \in \mathbb{N}$ s.t. $\forall n > N_\delta$, $|x_n - y| < \delta$; otherwise, $\forall N_\delta$, $\exists n > N_\delta$ s.t. $|x_n - y| \geq \delta$. Then $\{n || x_n - y| \leq \delta, n \in \mathbb{N}\}$ finite.

Contradiction.

ϕ cont. so by def. $\forall \epsilon > 0$, $\exists \delta > 0$ s.t. if $|x_n - y| < \delta$, then $|\phi(x_n) - \phi(y)| < \epsilon$.

Pick N_δ s.t. $\forall n > N_\delta$, $|x_n - y| < \delta$, and so $|\phi(x_n) - \phi(y)| < \epsilon$. There are infinitely many $\phi(x_n)$'s that satisfy this, and so $\phi(y)$ is a limit pt.

If $\exists y_1, y_2 \in X$ s.t. $\phi(y_1) = y_1$, then
 $\phi(y_2) = y_2$

$$d(y_1, y_2) = d(\phi(y_1), \phi(y_2)) \leq cd(y_1, y_2) \text{ with } c < 1$$

so $c = 1$

Theorem 4 (Inverse Function Theorem). *Suppose open $U \subset \mathbb{R}^n$, let $C^1 f : U \rightarrow \mathbb{R}^n$, $x_0 \in U$ s.t. $Df(x_0)$ invertible. Then \exists open neighborhoods $V \ni x_0$, $W \ni f(x_0)$ s.t. $V \subseteq U$ and $W \subseteq \mathbb{R}^n$, respectively, and s.t.*

- (i) $f : V \rightarrow W$ bijection
- (ii) $g = f^{-1} : V \rightarrow U$ differentiable, i.e. $g = f^{-1} : W \rightarrow V$ is C^1
- (iii) $D(f^{-1})$ cont. on W .
- (iv) $Dg(y) = (Df(g(y)))^{-1} \quad \forall y \in W$

Also, notice that $f(g(y)) = y \forall y \in W$.

Proof. Consider $\tilde{f}(x) = (Df(x_0))^{-1}(f(x + x_0) - f(x_0))$. Then
 $\tilde{f}(0) = 0$ and

$$D\tilde{f} = (Df(x_0))^{-1}(Df(x + x_0) - 0)$$
$$D\tilde{f}(0) = (Df(x_0))^{-1}Df(x_0) = 1$$

So let $\tilde{f} \rightarrow f$ (notation) and so assume, without loss of generality, that $U \ni 0$, $f(0) = 0$, $Df(0) = 1$
Choose $0 < \epsilon \leq \frac{1}{2}$. Let $0 < \delta < 1$ s.t. open ball $V = B_\delta(0) \subseteq U$, and $\|Df(x) - 1\| < \epsilon$. $\forall x \in U$, since Df cont. at 0.
Let $W = f(V)$.

$\forall y \in W$, define $\phi_y : V \rightarrow \mathbb{R}^n$
 $\phi_y(x) = x + (y - f(x))$

$$D(\phi_y)(x) = 1 + -Df(x) \quad \forall x \in V$$
$$\|D(\phi_y)(x)\| = \|1 - Df(x)\| \leq \epsilon < 1$$

$\forall x_1, x_2 \in V$, by mean value Thm. (not the equality that is only valid in 1-dim., but the inequality, that's valid for \mathbb{R}^d ,
 $\|\phi_y(x_1) - \phi_y(x_2)\| \leq \|D(\phi_y)(x')\| \|x_1 - x_2\|$

for some $x' = cx_2 + (1 - c)x_1$, $c \in [0, 1]$. V only needed to be convex set.
 $\implies \|\phi_y(x_1) - \phi_y(x_2)\| \leq \epsilon \|x_1 - x_2\|$

Then ϕ_y contraction mapping.
Suppose $f(x_1) = f(x_2) = y$, $x_1, x_2 \in V$.

$$\phi_y(x_1) = x_1$$
$$\phi_y(x_2) = x_2$$
$$\|\phi_y(x_1) - \phi_y(x_2)\| = \|x_1 - x_2\| \leq \epsilon \|x_1 - x_2\| \quad \forall \epsilon > 0 \implies x_1 = x_2$$

$\implies f|_U$ injective.
 $W = f(V)$, so $f : V \rightarrow W$ surjective. f bijective.
Fix $y_0 \in W$, $y_0 = f(x_0)$, $x_0 \in V$.
Let $r > 0$ s.t. $B_r(x_0) \subset V$.
Consider $B_{r\epsilon}(y_0)$. If $y \in B_{r\epsilon}(y_0)$.

$$r\epsilon > \|y - y_0\| = \|y - f(x_0)\| = \|\phi_y(x_0) - x_0\| \text{ with}$$
$$\phi_y(x) = x + (y - f(x))$$

If $x \in B_r(x_0)$,

$$\|\phi_y(x) - x_0\| \leq \|\phi_y(x) - \phi_y(x_0)\| + \|\phi_y(x_0) - x_0\| \leq \epsilon \|x - x_0\| + r\epsilon < 2r\epsilon = r$$

Thus $\phi(B_r(x_0)) = B_r(x_0)$.
By contraction mapping principle, $\exists a \in B_r(x_0)$, s.t. $\phi_y(a) = a$. Then $\phi_y(a) = a + (y - f(a)) = a \implies f(a) = y$.
 $y \in f(V) = W$.
So $B_{r\epsilon}(y_0) \subset W$. W open.

Let $\text{Mat}(n, n) \equiv$ space of all $n \times n$ matrices; $\text{Mat}(n, n) = \mathbb{R}^{n^2}$.

□

There is a proof of the implicit function theorem and its various forms in Shastri (2011) [5], but I found Wienhard's Handout 4 for Math 327 to be clearer.¹

Theorem 5 (Implicit Function Theorem). *Let open $U \subset \mathbb{R}^{m+n} \equiv \mathbb{R}^m \times \mathbb{R}^n$*

$C^1 f : U \rightarrow \mathbb{R}^n$
 $(a, b) \in U$ s.t. $f(a, b) = 0$ and $D_y f|_{(a, b)}$ invertible.
Then \exists open $V \ni (a, b)$, $V \subset U$
 \exists open neighborhood $W \ni a$, $W \subseteq \mathbb{R}^m$
 $\exists!$ $C^1 g : W \rightarrow \mathbb{R}^n$ s.t.

$$\{(x, y) \in V | f(x, y) = 0\} = \{(x, g(x)) | x \in W\}$$

Moreover,

$$dg_x = - (d_y f)^{-1} \Big|_{(x, g(x))} d_x f|_{(x, g(x))}$$

and g smooth if f .

Proof. Define $F : U \rightarrow \mathbb{R}^{m+n}$

$$F(x, y) = (x, f(x, y))$$

Then $F(a, b) = (a, 0)$ (given), and

$$DF = \begin{bmatrix} 1 & \\ \frac{\partial f^i(x, y)}{\partial x^j} & \frac{\partial f^i(x, y)}{\partial y^j} \end{bmatrix} \equiv \begin{bmatrix} 1 & \\ D_x f & D_y f \end{bmatrix}$$

$DF(a, b)$ invertible.
By inverse function theorem, since $DF(a, b)$ invertible at pt. (a, b) ,

\exists open neighborhoods $V \ni (a, b) \subseteq \mathbb{R}^m \times \mathbb{R}^n$ s.t. F diffeomorphism with $F^{-1} : \widetilde{W} \rightarrow V$.
 $\widetilde{W} \ni (a, 0) \subseteq \mathbb{R}^m \times \mathbb{R}^n$

Set $W = \{x \in \mathbb{R}^m | (x, 0) \in \widetilde{W}\}$. Then $\pi_1(\widetilde{W}) = W$ open in \mathbb{R}^m .
Define $g : W \rightarrow \mathbb{R}^n$,

$$g(x) = \pi_2 \circ F^{-1}(x, 0) \text{ or}$$
$$F^{-1}(x, 0) = (h(x), g(x))$$

Now $FF^{-1}(x, 0) = (x, 0) = (h(x), f(h(x), g(x)))$ so $h(x) = x \forall x \in W$, $0 = f(x, g(x))$.
Then

$$\{(x, y) \in V | f(x, y) = 0\} = \{(x, y) \in V | F(x, y) = (x, 0)\} = \{(x, g(x)) | x \in W, 0 = f(x, g(x))\}$$

Since π smooth and F^{-1} is C^1 , g is C^1 .
To reiterate, $f(x, g(x)) = 0$ on W .

¹<https://web.math.princeton.edu/~wienhard/teaching/M327/handout4.pdf>

Using chain rule while differentiating $f(x, g(x)) = 0$,

$$\partial_{x^j} f(x, g(x)) = \frac{\partial f(x, g(x))}{\partial x^k} \frac{\partial x^k}{\partial x^j} + \frac{\partial f(x, g(x))}{\partial y^k} \frac{\partial g^k(x)}{\partial x^j} = D_x f|_{(x, g(x))} + (D_y f)|_{(x, g(x))} \cdot (Dg)_x = 0 \text{ or}$$

$$(Dg)_x = - (D_y f)|_{x, g(x)} D_x f|_{(x, g(x))}$$

2. IMMERSIONS

Definition 2 (Immersion). *smooth $f : M \rightarrow N$, s.t. $Df(p) : T_p M \rightarrow T_{f(p)} N$ injective. Then f **immersion** at p .*

Absil, Mahony, and Sepulchre [7] pointed out that another definition for a *immersion* can utilize the theorem that rank of $Df \equiv DF = \dim T_p M$. Indeed, recall these facts from linear algebra:
for $T : V \rightarrow W$,

It's always true that $\text{rank} T \leq V$, and
 $\text{rank} T \leq W$

$\text{rank} T = \dim V$ iff T injective.
 $\text{rank} T = \dim W$ iff T surjective.

$$\begin{array}{ccc} T_x M & \xrightarrow{DF(x)} & T_{F(x)} N = T_y N \\ \uparrow & & \uparrow \\ x \in M & \xrightarrow{F} & y = F(x) \in N \end{array}$$

$$M \xrightarrow{F} N$$

Now

$$\begin{aligned} \dim T_x M &= \dim M \\ \dim T_{F(x)} N &= \dim N \end{aligned}$$

And

$$\text{rank}(DF(x)) \equiv \text{rank of } F$$

I know that the notation above is confusing, but this is what all Differential Geometry books apparently mean when they say "rank of F ".

Now

$$\text{rank}(DF(x)) = \dim(\text{im}(DF(x))) = \dim T_x M \text{ iff } DF(x) \text{ injective}$$

If $\forall x \in M$, this is the case, then F an **immersion**.

Apply the rank-nullity theorem in this case:

$$\begin{aligned} \text{rank}(DF(x)) + \dim \ker(DF(x)) &= \dim T_x M = \dim M \\ \implies \text{rank}(DF(x)) &= \dim M \leq \dim T_{F(x)} N = \dim N \text{ or } \dim M \leq \dim N \end{aligned}$$

Now

$$\text{rank}(DF(x)) = \dim T_{F(x)} N \text{ iff } DF(x) \text{ surjective}$$

If $\forall x \in M$, this is the case, then F an **submersion**.

$$\text{rank}(DF(x)) = \dim T_{F(x)} N = \dim N \leq \dim M$$

Shastri (2011) has this as the "Injective Form of Implicit Function Theorem", Thm. 1.4.5, pp. 23 and Guillemin and Pollack (2010) has this as the "Local Immersion Theorem" on pp. 15, Section 3 "The Inverse Function Theorem and Immersions" [4].

□ **Theorem 6** (Local immersion Theorem i.e. Injective Form of Implicit Function Theorem). *Suppose $f : M \rightarrow N$ immersion at p , $q = f(p)$.*

Then \exists local coordinates around p, q , x, y , respectively s.t. $f(x_1 \dots x_m) = (x_1 \dots x_m, 0 \dots 0)$.

Proof. Choose local parametrizations

$$\begin{array}{ccc} U \subseteq M & \xrightarrow{f} & N \supseteq V \\ \downarrow \phi & & \downarrow \psi \\ \phi(U) & \xrightarrow{f} & \psi(V) \end{array} \quad \begin{aligned} \phi(p) &= x \\ \psi(q) &= y \end{aligned}$$

$D(\psi f \phi^{-1}) \equiv Df$. $Df(p)$ injective (given f immersion). $Df(p) \in \text{Mat}(n, m)$

By change of basis in \mathbb{R}^n , assume $Df(p) = \begin{pmatrix} I_m \\ 0 \end{pmatrix}$.

Now define $G : \phi(U) \times \mathbb{R}^{n-m} \rightarrow \mathbb{R}^n$

$$G(x, z) = f(x) + (0, z)$$

Thus, $DG(x, z) = 1$ and for open $\phi(U) \times U_2$, $G(\phi(U) \times U_2)$ open.

By inverse function theorem, G local diffeomorphism of \mathbb{R}^n , at 0.

Now $f = G \circ i$, where i is canonical immersion.

$$\begin{aligned} G(x, 0) &= f(x) \\ \implies G^{-1}G(x, 0) &= (x, 0) = G^{-1}f(x) \end{aligned}$$

Use $\psi \circ G$ as the local parametrization of N around pt. q . Shrink U, V so that

$$\begin{array}{ccc} U \subseteq M & \xrightarrow{f} & N \supseteq V \\ \downarrow \phi & & \downarrow \psi \circ G \\ \phi(U) & \xrightarrow{i} & \psi \circ G(V) \end{array}$$

□

Theorem 7 (Implicit Function Thm.). *Let open subset $U \subseteq \mathbb{R}^n \times \mathbb{R}^d$, $(x, y) = (x^1 \dots x^n, y^1 \dots y^k)$ on U . Suppose smooth $\Phi : U \rightarrow \mathbb{R}^k$, $(a, b) \in U$, $c = \Phi(a, b)$*

If $k \times k$ matrix $\frac{\partial \Phi^i}{\partial y^j}(a, b)$ nonsingular, then \exists neighborhoods $V_0 \subseteq \mathbb{R}^n$ of a and smooth $F : V_0 \rightarrow W_0$ s.t.
 $W_0 \subseteq \mathbb{R}^k$ of b

$\Phi^{-1}(c) \cap (V_0 \times W_0)$ is graph of F , i.e.
 $\Phi(x, y) = c$ for $(x, y) \in V_0 \times W_0$ iff $y = F(x)$.

3. SUBMERSIONS; RANK THEOREM

cf. pp. 20, Sec. 4 "Submersions", Ch. 1 of Guillemin and Pollack (2010) [4].

Consider $X, Y \in \mathbf{Man}$, s.t. $\dim X \geq \dim Y$.

Definition 3 (submersion). *If $f : X \rightarrow Y$,
if $Df_x \equiv df_x$ is surjective, $f \equiv \textbf{submersion}$ at x .*

Recall that,

$$\begin{aligned} Df_x : T_x X &\rightarrow T_{f(x)} Y \\ \dim T_x X &\geq \dim T_{f(x)} Y \\ \text{rank } Df_x &\leq \dim T_{f(x)} Y, \text{ in general, while} \\ \text{rank } Df_x &= \dim T_{f(x)} Y \text{ iff } Df_x \text{ surjective} \end{aligned}$$

Canonical submersion is standard projection:

If $\dim X = k$, $k \geq l$,

$\dim Y = l$

$$(a_1 \dots a_k) \mapsto (a_1 \dots a_l)$$

Theorem 8 (Local Submersion Theorem). *Suppose $f : X \rightarrow Y$ submersion at x , and $y = f(x)$, Then \exists local coordinates around x, y s.t.*

$$f(x_1 \dots x_k) = (x_1 \dots x_l)$$

i.e. f locally equivalent to canonical submersion near x

Proof. I'll have a side-by-side comparison of my notation and the 1 used in Guillemin and Pollack (2010) [4] where I can.

For charts $(U, \phi), (V, \psi)$ for X, Y , respectively, $y = f(x)$ for $x \in X$,

$$\begin{array}{ccc} U \subseteq X & \xrightarrow{f} & Y \supseteq V \\ \downarrow \phi & & \downarrow \psi \circ G \\ \mathbb{R}^k & \xrightarrow{\mathbf{i}} & \mathbb{R}^l \end{array} \quad \begin{array}{ccc} x & \xrightarrow{f} & f(x) = y \\ \downarrow \phi & & \downarrow \psi \\ \phi(x) = (a^1 \dots a^k) & \xrightarrow{g} & g(\phi(x)) = g(a^1 \dots a^k) = \psi(y) \end{array}$$

Dg_x surjective, so assume it's a $l \times k$ matrix $\begin{bmatrix} \mathbf{1}_l & 0 \end{bmatrix}$.

Define

$$\begin{aligned} (6) \quad G : U \subset \mathbb{R}^k &\rightarrow \mathbb{R}^k \\ G(a) \equiv G(a^1 \dots a^k) &:= (g(a), a_{l+1}, \dots, a_k) \end{aligned}$$

Now

$$(7) \quad DG(a) = \begin{bmatrix} \mathbf{1}_l & 0 \\ & \mathbf{1}_{k-l} \end{bmatrix} = \mathbf{1}_k$$

so G local diffeomorphism (at 0).

So $\exists G^{-1}$ as local diffeomorphism of some U' of a into $U \subset \mathbb{R}^k$.

By construction,

$$(8) \quad g = \mathbb{P}_l \circ G$$

where \mathbb{P}_l is the *canonical submersion*, the projection operator onto \mathbb{R}^l .

$$g \circ G^{-1} = \mathbb{P}_l$$

(since G diffeomorphism)

$$\begin{array}{ccc} U \subseteq X & \xrightarrow{f} & V \subseteq Y \\ \uparrow \phi^{-1} \circ G^{-1} & & \uparrow \psi^{-1} \\ \mathbb{R}^k & \xrightarrow{\mathbb{P}_l} & \mathbb{R}^l \end{array} \quad \text{for} \quad \begin{array}{ccc} \phi^{-1} \circ G^{-1}(a) \equiv \phi^{-1} \circ G^{-1}(a^1 \dots a^k) = x & \xrightarrow{f} & f(x) = y = \psi^{-1}(a^1 \dots a^l) \\ \uparrow \phi^{-1} \circ G^{-1} & & \uparrow \psi^{-1} \\ (a^1 \dots a^k) & \xrightarrow{\mathbb{P}_l} & (a^1 \dots a^l) \end{array}$$

\implies

□

"An obvious corollary worth noting is that if f is a submersion at x , then it is actually a submersion in a whole neighborhood of x ." Guillemin and Pollack (2010) [4]

Suppose f submersion at $x \in f^{-1}(y)$.

By local submersion theorem

$$f(x_1 \dots x_k) = (x_1 \dots x_l)$$

Choose $y = (0, \dots, 0)$.

Then, near x , $f^{-1}(y) = \{(0, \dots, 0, x_{l+1} \dots x_k)\}$ i.e. let $V \ni x$ neighborhood of x , define $(x_1 \dots x_k)$ on V .

Then $f^{-1}(y) \cap V = \{(0 \dots 0, x_{l+1}, \dots x_k) | x_1 = 0, \dots x_l = 0\}$.

Thus $x_{l+1}, \dots x_k$ form a coordinate system on open set $f^{-1}(y) \cap V \subseteq f^{-1}(y)$.

Indeed,

$$\begin{array}{ccc} U \subseteq X & \xrightarrow{f} & V \subseteq Y \\ \downarrow \phi & & \downarrow \psi \\ \mathbb{R}^k & \xrightarrow{\mathbb{P}_l} & \mathbb{R}^l \end{array} \quad \begin{array}{ccc} x & \xrightarrow{f} & f(x) = y \\ \downarrow \phi & & \downarrow \psi \\ \phi(x) = (x^1 \dots x^k) & \xrightarrow{\mathbb{P}_l} & (x^1 \dots x^l) \end{array}$$

and now

$$\begin{array}{ccc} f^{-1}(y) & \xleftarrow{f^{-1}} & y \\ \uparrow \phi^{-1} & & \downarrow \psi \\ \{(0, \dots, 0, x^1 \dots x^k)\} & \xleftarrow{\mathbb{P}_l^{-1}} & (0 \dots 0) \end{array}$$

3.1. Rank Theorem. Lee (2012) [3] in pp. 85, Ch. 4 Submersions, Immersions, and Embeddings, combines Theorems 6, 8 (local immersion and local submersion theorems, respectively) into the "Rank Theorem" (cf. Thm 4.12 "Rank Theorem" of Lee (2012)):

Theorem 9 (Rank Theorem). *Suppose smooth manifolds M, N , $\dim M = m$, $\dim N = n$, smooth map $F : M \rightarrow N$, F has constant rank r .*

$\forall p \in M$, \exists smooth charts (U, φ) for M , centered at p , (V, ψ) for N , centered at $F(p)$, s.t.

$$F(U) \subseteq V$$

in which F has coordinate representation of form

$$(9) \quad \widehat{F}(x^1 \dots x^r, x^{r+1} \dots x^m) = (x^1 \dots x^r, 0 \dots 0)$$

Particularly, if F smooth submersion,

$$\widehat{F}(x^1 \dots x^n, x^{n+1} \dots x^m) = (x^1 \dots x^n$$

and if F smooth immersion

$$\widehat{F}(x^1 \dots x^m) = (x^1 \dots x^m, 0 \dots 0)$$

$$\begin{array}{ccc} \mathbb{R}^m & \xrightarrow{\psi F \varphi^{-1}} & \mathbb{R}^n \\ \uparrow \varphi & & \uparrow \psi \\ U \subset M & \xrightarrow{F} & V \subset N \end{array}$$

Also remember that $DF(p) : T_p M \rightarrow T_{F(p)} N$

Proof. $DF(p)$ has rank r (given). Then $DF(p)$ is some $r \times r$ submatrix of a $n \times m$ matrix s.t. $\det DF(p)$ nonzero.

By change of basis in \mathbb{R}^n , or reordering coordinates, assume $DF(p)$ is upper left submatrix $\left(\frac{\partial F^i}{\partial x^j} \right) \quad \forall i, j = 1, \dots, r$.

Relabel standard coordinate as

$$(x, y) = (x^1 \dots x^r, y^1 \dots y^{m-r}) \in \mathbb{R}^m$$

$$(v, w) = (v^1 \dots v^r, w^1 \dots w^{n-r}) \in \mathbb{R}^n$$

By initial translations of coordinates, assume without loss of generality $p = (0, 0)$, $F(p) = (0, 0)$

Suppose

$$F(x, y) = (Q(x, y), R(x, y))$$

for some smooth maps $Q : U \rightarrow \mathbb{R}^r$, $R : U \rightarrow \mathbb{R}^{n-r}$

Define

$$\varphi : U \rightarrow \mathbb{R}^m$$

$$\varphi(x, y) = (Q(x, y), y)$$

so

$$D\varphi(0, 0) = \begin{pmatrix} \frac{\partial Q^i}{\partial x^j}(0, 0) & \frac{\partial Q^i}{\partial y^j}(0, 0) \\ 0 & \delta_j^i \end{pmatrix}$$

$D\varphi(0, 0)$ nonsingular, since $\det \frac{\partial Q^i}{\partial x^j} \neq 0$ (by hypothesis).

By inverse function thm., \exists connected neighborhoods U_0 of $(0, 0)$, \widetilde{U}_0 of $\varphi(0, 0) = (0, 0)$ s.t.

$$\varphi : U_0 \rightarrow \widetilde{U}_0$$

is a diffeomorphism.

By shrinking U_0, \widetilde{U}_0 , assume \widetilde{U}_0 open cube.

Write $\varphi^{-1}(x, y) = (A(x, y), B(x, y))$, for some smooth functions $A : \widetilde{U}_0 \rightarrow \mathbb{R}^r$,

$$B : \widetilde{U}_0 \rightarrow \mathbb{R}^{m-r}$$

$$(x, y) = \varphi(A(x, y), B(x, y)) = (Q(A(x, y), B(x, y)), B(x, y))$$

$$\begin{aligned} B(x, y) &= y \\ \implies \varphi^{-1}(x, y) &= (A(x, y), y) \end{aligned}$$

$$\varphi \varphi^{-1} = 1 \implies x = Q(A(x, y), y)$$

Recall that we had hypotehsized that

$$F(x, y) = (Q(x, y), R(x, y))$$

Then

$$F \circ \varphi^{-1}(x, y) = F(A(x, y), y) = (Q(A(x, y), y), R(A(x, y), y)) = (x, R(A(x, y), y))$$

and so

$$F \circ \varphi^{-1}(x, y) = (x, \widetilde{R}(x, y))$$

where $\widetilde{R} : \widetilde{U}_0 \rightarrow \mathbb{R}^{n-r}$

$$\widetilde{R}(x, y) = R(A(x, y), y)$$

Compute

$$D(F \circ \varphi^{-1})(x, y) = \begin{pmatrix} \delta_j^i & 0 \\ \frac{\partial \widetilde{R}^i}{\partial x^j}(x, y) & \frac{\partial \widetilde{R}^i}{\partial y^j}(x, y) \end{pmatrix}$$

Since composing with a diffeomorphism doesn't change rank of map, $D(F \circ \varphi^{-1})$ has rank r everywhere in \widetilde{U}_0 .

$\left(\frac{\partial \widetilde{R}^i}{\partial x^j}(x, y) \right) \quad j = 1 \dots r$ are linearly independent, so $\frac{\partial \widetilde{R}^i}{\partial y^j}(x, y) = 0$ on \widetilde{U}_0 , so \widetilde{R}^i independent of y^j .

Let $S(x) = \widetilde{R}(x, 0)$, then

$$F \circ \varphi^{-1}(x, y) = (x, S(x))$$

Let open $V_0 \subseteq V$, $(0, 0) \in V$ be an open subset $V_0 = \{(v, w) \in V : (v, 0) \in \widetilde{U}_0\}$.

Then V_0 is a neighborhood of $(0, 0)$.

Because \widetilde{U}_0 is a cube, $F \circ \varphi^{-1}(x, y) = (x, S(x))$,

$$F \circ \varphi^{-1}(\widetilde{U}_0) \subseteq V_0$$

so $F(U_0) \subseteq V_0$.

Define $\psi : V_0 \rightarrow \mathbb{R}^n$

$$\psi(v, w) = (v, w - S(v))$$

Because $\psi^{-1}(s, t) = (s, t + S(s))$,

it is a diffeomorphism.

Thus (V_0, ψ) is a smooth chart.

$$\psi \circ F(\varphi^{-1}(x, y)) = \psi(x, S(x)) = (x, S(x) - S(x)) = (x, 0)$$

□

Definition 4 (regular value). *For smooth $f : X \rightarrow Y$, $X, Y \in \mathbf{Man}$,*

*$y \in Y$ is a **regular value** for f if $Df_x : T_x X \rightarrow T_y Y$ surjective $\forall x$ s.t. $f(x) = y$.*

*$y \in Y$ **critical value** if y not a regular value of f .*

Absil, Mahony, and Sepulchre [7] pointed out that another definition for a *regular value* can utilize the theorem that rank of $Df \equiv DF = \dim T_p N = \dim N$, iff $DF(p)$ surjective, for $p \in M$, $F : M \rightarrow N$. Then **regular value** $y \in N$, of F , if rank of $F \equiv \text{rank}(DF(x)) = \dim N$, $\forall x \in F^{-1}(y)$, for $F : M \rightarrow N$.

Theorem 10 (Preimage theorem). *If y regular value of $f : X \rightarrow Y$, $f^{-1}(y)$ is a submanifold of X , with $\dim f^{-1}(y) = \dim X - \dim Y$*

Proof. Given y is a regular value of $f : X \rightarrow Y$, $\forall x \in f^{-1}(y)$, $Df_x : T_x X \rightarrow T_y Y$ is surjective. By local submersion theorem,

$$f(x^1 \dots x^k) = (x^1 \dots x^l) = y$$

Since $x \in f^{-1}(y)$, $(x^1 \dots x^k) = (y^1 \dots y^l, x^{l+1} \dots x^k)$.

For this chart for (U, φ) , $U \ni x$, consider $(U \cap f^{-1}(y), \psi)$ with $\psi(x) = (x^{l+1} \dots x^k) \quad \forall x \in U \cap f^{-1}(y)$. $\forall f^{-1}(y)$ submanifold with $\dim f^{-1}(y) = k - l = \dim X - \dim Y$.

Examples for emphasis

If $\dim X > \dim Y$,

if $y \in Y$, regular value of $f : X \rightarrow Y$,
 f submersion, $\forall x \in f^{-1}(y)$

If $\dim X = \dim Y$,

f local diffeomorphism $\forall x \in f^{-1}(y)$

If $\dim X < \dim Y$, $\forall y \in f(X)$ is a critical value.

Example: $O(n)$ as a submanifold of $\text{Mat}(n, n)$

Given $\text{Mat}(n, n) \equiv M(n) = \{n \times n \text{ matrices}\}$ is a manifold; in fact $\text{Mat}(n, n) \cong \mathbb{R}^{n^2}$, Consider $O(n) = \{A \in \text{Mat}(n, n) | AA^T = 1\}$.

$$(10) \quad AA^T \in \text{Sym}(n) \equiv S(n) = \{S \in \text{Mat}(n, n) | S^T = S\} = \{\text{symmetric } n \times n \text{ matrices}\}$$

$\text{Sym}(n)$ submanifold of $\text{Mat}(n, n)$, $\text{Sym}(n)$ diffeomorphic to \mathbb{R}^k (i.e. $\text{Sym}(n) \cong \mathbb{R}^k$), $k := \frac{n(n+1)}{2}$.

$$f : \text{Mat}(n, n) \rightarrow \text{Sym}(n)$$

$$f(A) = AA^T$$

Notice f is smooth,

$$f^{-1}(1) = O(n)$$

$$Df_A(B) = \lim_{s \rightarrow 0} \frac{f(A + sB) - f(A)}{s} = \lim_{s \rightarrow 0} \frac{(A + sB)(A^T + sB^T) - AA^T}{s} = AB^T + BA^T$$

If $Df_A : T_A \text{Mat}(n, n) \rightarrow T_{f(A)} \text{Sym}(n)$ surjective when $A \in f^{-1}(1) = O(n)$ (???).

Proposition 1. *If smooth $g_1 \dots g_l \in C^\infty(X)$ on X are independent $\forall x \in X$, s.t. $g_i(x) = 0$, $\forall i = 1 \dots l$, then $Z = \{x \in X | g_1(x) = \dots = g_l(x) = 0\} = \text{set of "common zeros"}$ is a submanifold of X s.t. $\dim Z = \dim X - l$.*

Take note that $g_1 \dots g_l$ are independent at x means, really, that $D(g_1)_x \dots D(g_l)_x$ are linearly independent on $T_x X$.

Proof. Suppose smooth $g_1 \dots g_l \in C^\infty(X)$ on manifold X s.t. $\dim X = k \geq l$.

Consider $g = (g_1 \dots g_l) : X \rightarrow \mathbb{R}^l$, $Z \equiv g^{-1}(0)$.

Since $\forall g_i$ smooth, $D(g_i)_x : T_x X \rightarrow \mathbb{R}$ linear.

Now for

$$Dg_x = (D(g_1)_x \dots D(g_l)_x) : T_x X \rightarrow \mathbb{R}^l$$

By rank-nullity theorem (linear algebra), Dg_x surjective iff $\text{rank } Dg_x = l$ i.e. l functionals $D(g_1)_x \dots D(g_l)_x$ are linearly independent on $T_x X$.

"We express this condition by saying the l functions $g_1 \dots g_l$ are independent at x ." (Guillemin and Pollack (2010) [4])

4. SUBMANIFOLDS; IMMERSED SUBMANIFOLD, EMBEDDED SUBMANIFOLDS, REGULAR SUBMANIFOLDS

Definition 5 (Embedded Submanifold).

Recall immersion:

$F : M \rightarrow N$ immersion iff DF injective, i.e. iff $\text{rank } DF = \dim M$.

Consider manifolds $M \subseteq N$.

Consider inclusion map $i : M \rightarrow N$.

$$i : x \mapsto x$$

If i immersion, $Di(x) = \frac{\partial y^i}{\partial x^j} = \delta_j^i$ if $y^i = x^i$, $\forall i = 1, \dots, \dim M$.

□ **Definition 6** (immersed submanifold). ***immersed submanifold** $M \subseteq N$ if inclusion $i : M \rightarrow N$ is an immersion.*

cf. 3.3 Embedded Submanifolds of Absil, Mahony, and Sepulchre [7], also Ch. 5 Submanifolds, pp. 108, **Immersed Submanifolds** of John Lee (2012) [3].

Immersed submanifolds often arise as images of immersions.

Proposition 2 (Images of Immersions as submanifolds). *Suppose smooth manifold M , smooth manifold with or without boundaries N ,*

injective, smooth immersion $F : M \rightarrow N$ (F injective itself, not just immersion)

Let $S = F(M)$.

Then S has unique topology and smooth structure of smooth submanifolds of N s.t. $F : M \rightarrow S$ diffeomorphism.

cf. Prop. 5.18 of John Lee (2012) [3].

Proof. Define topology of S : set $U \subseteq S$ open iff $F^{-1}(U) \subseteq M$ open ($F^{-1}(U \cap V) = F^{-1}(U) \cap F^{-1}(V)$, $F^{-1}(U \cup V) = F^{-1}(U) \cup F^{-1}(V)$).

Define smooth structure of S : $\{F(U), \varphi \circ F^{-1} | (U, \varphi) \in \text{atlas for } M, \text{ i.e. } (U, \varphi) \text{ any smooth chart of } M\}$.

"smooth compatibility condition":

$$(\varphi_2 \circ F^{-1})(\varphi_1 F^{-1})^{-1} = \varphi_2 \circ F^{-1} F \varphi_1^{-1} = \varphi_2 \varphi_1^{-1}$$

since $\varphi_2 \varphi_1^{-1}$ diffeomorphism ($\varphi_2 \varphi_1^{-1}$ bijection and it and inverse is differentiable)

F diffeomorphism onto $F(M)$.

and these are the only topology and smooth structure on S with this property:

$$S \xrightarrow{F^{-1}} M \xrightarrow{F} N \quad = \quad S \hookrightarrow M$$

and F^{-1} diffeomorphism, F smooth immersion, so $i : S \rightarrow M$ smooth immersion. □

5. CURVES, INTEGRAL CURVES, AND FLOWS

cf. John Lee (2012) [3], Ch. 9, deals with time-dependent vector fields and I don't see other texts or references handling such an important, but overlooked, case.

5.1. **Curves in Euclidean space.** cf. 4.1 Jeffrey Lee (2009) [2]

If C is 1-dim. submanifold of \mathbb{R}^n , $p \in C$, \exists chart (V, y) of C , $p \in V$ s.t. $y(V)$ is a connected open interval $I \subset \mathbb{R}$, inverse map $y^{-1} : I \rightarrow V \subset M$ is a local parametrization.

idea is to extract information that's appropriately independent of parametrization.

If $\gamma : I \rightarrow \mathbb{R}^n$, $c : J \rightarrow \mathbb{R}^n$ curves with same image, c is a **positive reparametrization** of γ if \exists smooth $h : J \rightarrow I$ with $h' > 0$ s.t. $c = \gamma \circ h$, in this case, γ, c have same **sense** and same **orientation**.

Assume $\gamma : I \rightarrow \mathbb{R}^n$ has $\|\gamma'\| > 0$, Such a curve is **regular**, i.e. curve is an immersion (Recall, an immersion would be smooth $\gamma : I \rightarrow \mathbb{R}^n$, s.t. $D\gamma(p) : T_p I \rightarrow T_{\gamma(p)} \mathbb{R}^n$ injective).

Definition 7 (unit tangent field in Euclidean space, 4.3 Lee (2009) [2]). *If $\gamma : I \rightarrow \mathbb{R}^n$ regular curve, then $\mathbf{T}(t) := \frac{\gamma'(t)}{\|\gamma'(t)\|}$ defines **unit tangent field** along γ ($\|T\| = 1$)*

length of a curve defined on closed interval $\gamma : [t_1, t_2] \rightarrow \mathbb{R}^n$,

$$L = \int_{t_1}^{t_2} \|\gamma'(t)\| dt$$

Define **arc length function** for curve $\gamma : I \rightarrow \mathbb{R}^n$ by choosing $t_0 \in I$,

$$s = h(t) := \int_{t_0}^t \|\gamma'(\tau)\| d\tau$$

If curve is smooth and regular, then $h' = \|\gamma'(\tau)\| > 0$, so by inverse function theorem, \exists smooth h^{-1} (since h' invertible (with $1/h'$)).

If $c(s) = \gamma h^{-1}(s)$, then $\|c'\|(s) := \|c'(s)\| = 1, \quad \forall s$

$$\begin{aligned} c'(s) &= (\gamma(h^{-1}(s)))' = \gamma'(h^{-1}(s)) \cdot \frac{dh^{-1}}{ds}(s) \\ \|c'(s)\| &= \|\gamma'(t)\| \left\| \frac{dh^{-1}}{ds}(s) \right\| = h' \cdot \frac{1}{h'} = 1 \end{aligned}$$

Curves parametrized by arc length are **unit speed curves**.

For a unit speed curve, $\frac{dc}{ds}(s) = \mathbf{T}(s)$.

Definition 8 (curvature vector, curvature function, principal normal in Euclidean space, 4.4 Lee (2009) [2]). *Let $c : I \rightarrow \mathbb{R}^n$ be a unit speed curve. vector valued function*

$$\kappa(s) := \frac{d\mathbf{T}}{ds}(s)$$

is called the **curvature vector**.

$$\kappa(s) := \|\kappa(s)\| = \left\| \frac{d\mathbf{T}}{ds}(s) \right\|$$

is called the **curvature function**.

If $\kappa(s) > 0$, then define principal normal

$$\mathbf{N}(s) = \left\| \frac{d\mathbf{T}}{ds}(s) \right\|^{-1} \frac{d\mathbf{T}}{ds}(s)$$

s.t. $\frac{d\mathbf{T}}{ds} = \kappa \mathbf{N}$

5.2. **Integral Curves.** cf. Ch. 9 of Lee (2012) [3]

integral curves - smooth curves whose velocity at each point is equal to the value of the smooth vector field there.

Let smooth manifold M with or without boundary.

If smooth curve $\gamma : J \rightarrow M$, then $\forall t \in J$, velocity vector $\gamma'(t)$ is a vector in $T_{\gamma(t)} M$.

Definition 9 (Integral Curve Γ). *Integral curve: \mathbf{v}, γ .*

If vector field \mathbf{v} on M ,

integral curve of \mathbf{v} is differentiable curve $\gamma : J \rightarrow M$, whose velocity $\forall p \in M$, $V_{\gamma(t)}$

$$\gamma'(t) = V_{\gamma(t)}, \forall t \in J$$

Finding integral curves boils downs to solving a system of ordinary differential equations in a smooth chart.

Suppose smooth vector field V on M , smooth curve $\gamma : J \rightarrow M$

On smooth coordinate domain $U \subseteq M$,

Write γ in local coordinates as

$$\begin{aligned} \gamma(t) &= (\gamma^1(t), \dots, \gamma^n(t)) \\ \gamma^i(t) &= x^i \circ \gamma(t) \text{ where } x^i : U \subseteq M \rightarrow \mathbb{R} \end{aligned}$$

Then condition for γ to be an integral curve.

$$\gamma'(t) \equiv \dot{\gamma}(t) = V_{\gamma(t)}$$

can be written as

$$\dot{\gamma}^i(t) \frac{\partial}{\partial x^i} \Big|_{\gamma(t)} = V^i(\gamma(t)) \frac{\partial}{\partial x^i} \Big|_{\gamma(t)}$$

which reduces to an autonomous system of ordinary differential equations (ODEs)

$$\begin{aligned} (11) \quad \dot{\gamma}^1(t) &= V^1(\gamma^1(t), \dots, \gamma^n(t)) \\ &\vdots \\ \dot{\gamma}^n(t) &= V^n(\gamma^1(t), \dots, \gamma^n(t)) \end{aligned}$$

20200306 EY: I use this notation which is more useful for me, e.g. taking the double derivative of the curve γ :

$$\begin{aligned} \dot{\gamma}(t) &= v(\gamma(t)) \in T_{\gamma(t)} U \\ \dot{\gamma}(t) &= v^i(\gamma^1(t), \dots, \gamma^n(t)) \\ \frac{d}{dt} \dot{\gamma}^i(t) &= \frac{d}{dt} v^i(\gamma^1(t), \dots, \gamma^n(t)) = \frac{d}{dt} v^i(x^1, \dots, x^n) \\ &= \frac{\partial v^i}{\partial x^j}(x) \dot{x}^j = \dot{\gamma}^j \frac{\partial \dot{\gamma}^i}{\partial x^j} \end{aligned}$$

6. TENSORS

I'll go through Ch.7 *Tensors* of Jeffrey Lee (2009) [2].

Definition 10 (7.1[2]). *Let V, W be modules over commutative ring R , with unity.*

Then, algebraic W -valued tensor on V is multilinear map.

$$(12) \quad \tau : V_1 \times V_2 \times \dots \times V_m \rightarrow W$$

where $V_i = \{V, V^*\} \quad \forall i = 1, 2, \dots, m$.

If for r, s s.t. $r + s = m$, there are $r \quad V_i = V^*, s \quad V_i = V$, tensor is r -contravariant, s -covariant; also say tensor of total type $\binom{r}{s}$.

EY : 20170404 Note that

$$(\tau_{\beta}^{i\alpha} \frac{\partial}{\partial x^i} \text{ or } \tau_{\beta}^{i\alpha} e_i)(\omega_j dx^j \text{ or } \omega_j e^j \in V^*)$$

$$(\tau_{i\alpha}^{\beta} dx^i \text{ or } \tau_{i\alpha}^{\beta} e^i)(X^j \frac{\partial}{\partial x^j} \text{ or } X^j e_j \in V)$$

\exists natural map $V \rightarrow V^{**}$, $\tilde{v} : \alpha \mapsto \alpha(v)$. If this map is an isomorphism, V is **reflexive** module, and identify V with V^{**} .
 $v \mapsto \tilde{v}$

Exercise 7.5. Given vector bundle $\pi : E \rightarrow M$, open $U \subset M$, consider sections of π on U , i.e. cont. $s : U \rightarrow E$, where $(\pi \circ s)(u) = u$, $\forall u \in U$.

Consider $E^* \ni \omega = \omega_i e^i$.

$\forall s \in \Gamma(E)$, $\omega(s) = \omega_i(s(x))^i$, $\forall x \in U \subset M$. So define $\tilde{s} : \omega, x \mapsto \omega(s(x))$, $\forall x \in U$.

If $\tilde{s} = 0$, $\tilde{s}(\omega, x) = \omega(s(x)) = 0 \quad \forall \omega \in E^*, \forall x \in U$, and so $s = 0$. (Let $\omega_i = \delta_{iJ}$ for some J , and so $s^J(x) = 0 \quad \forall J$).

$s = 0$. So $\ker(s \mapsto \tilde{s}) = \{0\}$ (so condition for injectivity is fulfilled).

Since $\tilde{s} : \omega, x \mapsto \omega(s(x))$, $\forall \omega \in E^*, \forall x \in U$, $s \mapsto \tilde{s}$ is surjective.

$s \mapsto \tilde{s}$ is an isomorphism so $\Gamma(E)$ is a *reflexive* module.

Proposition 3. For R a ring (special case), \exists module homomorphism:

tensor product space \rightarrow tensor, as a multilinear map, i.e. \exists

$$(13) \quad (\otimes_{i=1}^r V) \otimes (\otimes_{j=1}^s V^*) \rightarrow T_s^r(V; R) \\ u_1 \otimes \cdots \otimes u_r \otimes \beta^1 \otimes \cdots \otimes \beta^s \in (\otimes^r V) \otimes (\otimes^s V^*) \mapsto (\alpha^1 \dots \alpha^r, v_1 \dots v_s) \mapsto \alpha^1(u_1) \dots \alpha^r(u_r) \beta^1(v_1) \dots \beta^s(v_s)$$

Indeed, consider

$$(\alpha^1 \dots \alpha^r, v_1 \dots v_s) \in \underbrace{V^* \times \cdots \times V^*}_r \times \underbrace{V \times \cdots \times V}_s \mapsto \alpha^1(u_1) \dots \alpha^r(u_r) \beta^1(v_1) \dots \beta^s(v_s)$$

and so for

$$\alpha^i = \alpha_{\mu}^i e^{\mu}, \quad i = 1, 2, \dots, r, \mu = 1, 2, \dots, \dim V^* \quad \alpha^i(u_i) = \alpha_{\mu}^i u_i^{\mu} \\ v_i = v_i^{\mu} e_{\mu}, \quad i = 1, 2, \dots, s, \mu = 1, 2, \dots, \dim V \quad \beta^i(v_i) = \beta_{\mu}^i v_i^{\mu}$$

So that

$$\alpha^1(u_1) \dots \alpha^r(u_r) \beta^1(v_1) \dots \beta^s(v_s) = \alpha_{\alpha_1}^1 u_1^{\alpha_1} \dots \alpha_{\alpha_r}^r u_r^{\alpha_r} \beta_{\mu_1}^1 v_1^{\mu_1} \dots \beta_{\mu_s}^s v_s^{\mu_s} = \\ = (u_1^{\alpha_1} \dots u_r^{\alpha_r} \beta_{\mu_1}^1 \dots \beta_{\mu_s}^s)(\alpha_{\alpha_1}^1 \dots \alpha_{\alpha_r}^r v_1^{\mu_1} \dots v_s^{\mu_s})$$

Identify $u_1 \otimes \cdots \otimes u_r \otimes \beta^1 \otimes \cdots \otimes \beta^s$ with this multiplinear map.

Proposition 4. If V is finite-dim. vector space, or if $V = \Gamma(E)$, for vector bundle $E \rightarrow M$, map

$$(14) \quad (\otimes_{i=1}^r V) \otimes (\otimes_{j=1}^s V^*) \rightarrow T_s^r(V; R)$$

is an isomorphism.

Definition 11. tensor that can be written as

$$(15) \quad u_1 \otimes \cdots \otimes u_r \otimes \beta^1 \otimes \cdots \otimes \beta^s \equiv u_1 \otimes \cdots \otimes \beta^s$$

is *simple* or *decomposable*.

Now well that not *all* tensors are simple.

Definition 12 (7.7[2], tensor product). $\forall S \in T_{s_1}^{r_1}(V)$, $\forall T \in T_{s_2}^{r_2}(V)$, define tensor product

$$(16) \quad S \otimes T \in T_{s_1+s_2}^{r_1+r_2}(V) \\ S \otimes T(\theta^1 \dots \theta^{r_1+r_2}, v_1 \dots v_{s_1+s_2}) := S(\theta^1 \dots \theta^{r_1}, v_1 \dots v_{s_1})T(\theta^{r_1+1} \dots \theta^{r_1+r_2}, v_{s_1+1} \dots v_{s_1+s_2})$$

Proposition 5 (7.8[2]).

$$\tau^{i_1 \dots i_r}_{j_1 \dots j_s} e_{i_1} \otimes \cdots \otimes e_{i_r} \otimes e^{j_1} \otimes \cdots \otimes e^{j_s} = \tau(e^{i_1} \dots e^{i_r}, e_{j_1} \dots e_{j_s}) e_{i_1} \otimes \cdots \otimes e_{i_r} \otimes e^{j_1} \otimes \cdots \otimes e^{j_s} = \tau$$

So $\{e_{i_1} \otimes \cdots \otimes e_{i_r} \otimes e^{j_1} \otimes \cdots \otimes e^{j_s} | i_1 \dots i_r, j_1 \dots j_s \in 1 \dots n\}$ spans $T_s^r(V; R)$

Exercise 7.11. Let basis for V $e_1 \dots e_n$, corresponding dual basis for V^* $e^1 \dots e^n$

Let basis for V $\bar{e}_1 \dots \bar{e}_n$, corresponding dual basis for V^* $\bar{e}^1 \dots \bar{e}^n$

s.t.

$$\bar{e}_i = C_{\bar{i}}^k e_k$$

$$\bar{e}^i = (C^{-1})^i_{\bar{k}} e^{\bar{k}}$$

EY:20170404, keep in mind that

$$Ax = e_i A^i_{\bar{k}} e^{\bar{k}}(x^j e_j) = e_i A^i_{\bar{j}} x^j = A^i_{\bar{j}} x^j e_i$$

$$Ae_j = e_k A^k_{\bar{i}} e^{\bar{i}}(e_j) = A^k_{\bar{j}} e_k = \bar{e}_j$$

$$\bar{\tau}^i_{jk} \bar{e}_i \otimes \bar{e}^j \otimes \bar{e}^k = \bar{\tau}^i_{jk} C^l_{\bar{i}} e_l (C^{-1})^j_m e^m (C^{-1})^k_n e^n = \bar{\tau}^i_{jk} C^l_{\bar{i}} (C^{-1})^j_m (C^{-1})^k_n = \tau^l_{mn}$$

$$\bar{\tau}^i_{jk} = C^c_k C^b_j (C^{-1})^i_a \tau^a_{bc}$$

On Remark 7.13 of Jeffrey Lee (2009) [2]: first, egregious typo for $L(V, V)$; it should be $L(V, W)$. Onward, for $L(V, W)$,

consider $W \otimes V^* \ni w \otimes \alpha$ s.t.

$$(w \otimes \alpha)(v) = \alpha(v)w \in W, \forall v \in V, \text{ so } w \otimes \alpha \in L(V, W)$$

Now consider (category of) left R -module,

$$(17) \quad {}_R \mathbf{Mod} \ni {}_{\text{Mat}_{\mathbb{K}}(N, M)} \mathbb{K}^N$$

where

$$V = \mathbb{K}^N$$

$$W = \mathbb{K}^M$$

For $A \in \text{Mat}_{\mathbb{K}}(N, M)$, $x \in \mathbb{K}^N$,

$$e_i A^i_{\bar{\mu}} e^{\bar{\mu}}(x^{\nu} e_{\nu}) = Ax = e_i A^i_{\bar{\mu}} x^{\mu}, \quad i = 1, 2, \dots, M, \mu = 1, 2, \dots, N$$

$$A \in \text{Mat}_{\mathbb{K}}(N, M) \cong W \otimes V^* \cong L(V, W)$$

Consider

$$\alpha \in (\mathbb{K}^N)^* = V^* \quad \alpha = \alpha_{\mu} e^{\mu}$$

$$w \in \mathbb{K}^M = W \quad w = w^i e_i$$

$$\alpha \otimes w = w \otimes \alpha = w^i \alpha_{\mu} e_i \otimes e^{\mu}$$

(remember, isomorphism between $\text{Mat}_{\mathbb{K}}(N, M)$ and $W \otimes V^*$ guaranteed, if V, W are free R -modules, $R = \mathbb{K}$).

Let V, W be left R -modules, i.e. $V, W \in {}_R \mathbf{Mod}$.

$$V^* \in \mathbf{Mod}_R$$

For $V^* \otimes W \in \mathbf{Mod}_R \otimes {}_R \mathbf{Mod}$

$$\alpha \in V^*, w \in W$$

$$(\alpha \otimes w)(v) = \alpha(v)w, \text{ for } v \in V \in {}_R \mathbf{Mod}$$

But $(w \otimes \alpha)(v) = w\alpha(v)$.

Note $\alpha(v) \in R$.

Let V, W be right R -modules, i.e. $V, W \in \mathbf{Mod}_R$.

$$V^* \in {}_R \mathbf{Mod}$$

For $W \otimes V^* \in \mathbf{Mod}_R \otimes {}_R\mathbf{Mod}$.

$$\alpha \in V^*, w \in W$$

$$(v)(w \otimes \alpha) = w\alpha(v), \text{ with } \alpha(v) \in R, v \in V$$

So $W \otimes V^* \cong L(V, W)$, for $V, W \in \mathbf{Mod}_R$

Definition 13 (7.20[2], **contraction**). *Let (e_1, \dots, e_n) basis for V , $(e^1 \dots e^n)$ dual basis. If $\tau \in T_s^r(V)$, then for $k \leq r, l \leq s$, define*

$$(18) \quad \begin{aligned} & C_l^k \tau \in T_{s-1}^{r-1}(V) \\ & C_l^k \tau (\theta^1 \dots \theta^{r-1}, w_1 \dots w_{s-1}) := \\ & \sum_{a=1}^n \tau (\theta^1 \dots \underbrace{e^a}_{kth \text{ position}} \dots \theta^{r-1}, w_1 \dots \underbrace{e_a}_{ith \text{ position}} \dots w_{s-1}) \end{aligned}$$

C_l^k is called **contraction**, for some single $1 \leq k \leq r$, some single $1 \leq l \leq s$,

$$C_l^k : T_s^r(V) \rightarrow T_{s-1}^{r-1}(V)$$

s.t.

$$(C_l^k \tau)^{i_1 \dots \widehat{i_k} \dots i_r}_{j_1 \dots \widehat{j_l} \dots j_s} := \tau^{i_1 \dots a \dots i_r}_{j_1 \dots a \dots j_s}$$

Universal mapping properties can be invoked to give a basis free definition of contraction (EY : 20170405???).
IN general,

$$\forall v_1 \dots v_s \in V, \forall \alpha^1 \dots \alpha^r \in V^*$$

so that

$$\begin{aligned} v_j &= v_j^\mu e_\mu \quad j = 1 \dots s, \quad \mu = 1, \dots \dim V \\ \alpha^i &= \alpha_\mu^i e^\mu \quad i = 1 \dots r, \quad \mu = 1 \dots \dim V^* \end{aligned}$$

then $\forall \tau \in T_s^r(V)$,

$$\begin{aligned} \tau(\alpha^1 \dots \alpha^r, v_1 \dots v_s) &= \tau(\alpha_{\mu_1}^1 e^{\mu_1} \dots \alpha_{\mu_r}^r e^{\mu_r}, v_1^{\nu_1} e_{\nu_1} \dots v_s^{\nu_s} e_{\nu_s}) = \\ &= \alpha_{\mu_1}^1 \dots \alpha_{\mu_r}^r v_1^{\nu_1} \dots v_s^{\nu_s} \tau(e^{\mu_1} \dots e^{\mu_r}, e_{\nu_1} \dots e_{\nu_s}) = \alpha_{\mu_1}^1 \dots \alpha_{\mu_r}^r v_1^{\nu_1} \dots v_s^{\nu_s} \tau^{\mu_1 \dots \mu_r}_{\nu_1 \dots \nu_s} \end{aligned}$$

which is equivalent to

$$\begin{array}{ccc} \tau \in T_s^r(V) & \xrightarrow{\alpha^1 \otimes \dots \otimes \alpha^r \otimes v_1 \otimes \dots \otimes v_s \otimes} & \alpha^1 \otimes \dots \otimes \alpha^r \otimes v_1 \otimes \dots \otimes v_s \otimes \tau \\ & & \downarrow \\ & & C_{s+1}^1 C_{s+2}^2 \dots C_{r+s}^r C_1^r C_2^{r+1} \dots C_s^{r+s} \\ & & \downarrow \\ & & \tau(\alpha^1 \dots \alpha^r, v_1 \dots v_s) \in R \end{array}$$

where I've tried to express the right- R -module, "right action" on $\alpha^1 \otimes \dots \otimes \alpha^r \otimes v_1 \otimes \dots \otimes v_s \in V^* \otimes \dots \otimes V$.
Conlon (2008) [16]

Part 4. Lie Groups, Lie Algebra

7. LIE GROUPS

- : Lie Groups
- : Groups
- : Ring
- : group algebra
- : Group Ring
- : Representation Theory
- : Modules
- : kG -modules

From Sec. 8.1 "Noncommutative Rings" of Rotman (2010) [9]:

Definition 14. *ring R - additive abelian group equipped with multiplication $R \times R \rightarrow R$ s.t. $\forall a, b \in R$
 $(a, b) \mapsto ab$*

- (i) $a(bc) = (ab)c$
- (ii) $a(b+c) = ab+ac, (b+c)a = ba+ca$
- (iii) $\exists 1 \in R$ s.t. $\forall a \in R, 1a = a = a1$

Example 8.1[9]

- (ii) group algebra kG , k commutative ring, G group, "its additive abelian group is free k -module having basis labeled by elements of G ,
i.e. $\forall a \in kG, a = \sum_{g \in G} a_g g, a_g \in k, \forall g \in G, a_g \neq 0$ for only finitely many $g \in G$.

$$\begin{aligned} & \text{define (ring) multiplication } kG \times kG \rightarrow kG \quad \forall a, b \in kG, \quad a = \sum_{g \in G} a_g g \quad \text{to be} \\ & ab = ab \quad b = \sum_{h \in G} b_h h \end{aligned}$$

$$\left(\sum_{g \in G} a_g g \right) \left(\sum_{h \in G} b_h h \right) = \sum_{z \in G} \left(\sum_{gh=z} a_g b_h \right) z$$

Definition 15. *Given R ring, left R -module is (additive) abelian group M equipped with*

*scalar multiplication $R \times M \rightarrow M$ s.t. $\forall m, m' \in M, \forall r, r', 1 \in R$
 $(r, m) \mapsto rm$*

- (i) $r(m+m') = rm+rm'$
- (ii) $(r+r')m = rm+r'm$
- (iii) $(rr')m = r(r'm)$
- (iv) $1m = m$

EY : 20150922 Example : for kG -module V^σ , for $r \in kG$, so $r = \sum_{g \in G} a_g g$
 $R \times M \rightarrow M \quad kG \times V \rightarrow V$
 $(r, m) \mapsto rm \implies (r, v) \mapsto tv$

For some representation $\sigma : G \rightarrow GL(V)$,

$$rv = \sum_{g \in G} a_g g \cdot v = \sum_{g \in G} a_g \sigma_g(v)$$

So a kG -module needs to be associated with some chosen representation.

Note for V as an additive abelian group, $\forall u, v, w \in V$,

$$v + w = w + v, (u + v) + w = u + (v + w)$$

$$v + 0 = v \quad \forall v \in V \text{ for } 0 \in V$$

$$v + (-v) = 0 \quad \forall v \in V$$

So a vector space can be an additive abelian group.

Note that

$$r(v + w) = \left(\sum_{g \in G} a_g g \right) (v + w) = \left(\sum_{g \in G} a_g \sigma_g \right) (v + w) = \sum_{g \in G} a_g \sigma_g(v) + \sum_{g \in G} a_g \sigma_g(w) = rv + rw$$

$$(r + r')v = \left(\sum_{g \in G} a_g g + b_g g \right) v = \sum_{g \in G} (a_g \sigma_g + b_g \sigma_g) v = \sum_{g \in G} a_g \sigma_g(v) + \sum_{g \in G} b_g \sigma_g(v) = rv + r'v$$

$$(rr')v = \left(\sum_{g \in G} a_g g \sum_{h \in G} b_h h \right) v = \left(\sum_{z \in G} \sum_{gh=z} a_g b_h z \right) v = \sum_{z \in G} \sum_{gh \in z} a_g b_h \sigma_z(v) = \sum_{g \in G} \sum_{h \in G} a_g b_h \sigma_g \sigma_h(v)$$

since $\sigma(gh) = \sigma(g)\sigma(h) = \sigma_g \sigma_h = \sigma_{gh}$ (σ homomorphism)

$$1v = \sigma(1)v = 1v = v$$

From Sec. 8.3 “Semisimple Ring” of Rotman (2010) [9]:

Definition 16. k -representation of group G is homomorphism

$$\sigma : G \rightarrow GL(V)$$

where V is vector field over field k

Proposition 6 (8.37 Rotman (2010)[9]). $\forall k$ -representation $\sigma : G \rightarrow GL(V)$ equips V with structure of left kG -module, denote module by V^σ .

Conversely, \forall left kG -module V determines k -representation $\sigma : G \rightarrow GL(V)$

Proof. Given $\sigma : G \rightarrow GL(V)$,

$$\sigma_g =: \sigma(g) : V \rightarrow V$$

define

$$kG \times V \rightarrow V$$

$$\left(\sum_{g \in G} a_g g \right) v = \sum_{g \in G} a_g \sigma_g(v)$$

$$v, w \in V$$

Let $r, r', 1 \in kG$

$$r = \sum_{g \in G} a_g g$$

$$r(v + w) = \left(\sum_{g \in G} a_g g \right) (v + w) = \left(\sum_{g \in G} a_g \sigma_g \right) (v + w) = \sum_{g \in G} a_g \sigma_g(v) + \sum_{g \in G} a_g \sigma_g(w) = rv + rw$$

$$(r + r')v = \left(\sum_{g \in G} a_g g + b_g g \right) v = \sum_{g \in G} (a_g \sigma_g + b_g \sigma_g) v = \sum_{g \in G} a_g \sigma_g(v) + \sum_{g \in G} b_g \sigma_g(v) = rv + r'v$$

$$(rr')v = \left(\sum_{g \in G} a_g g \sum_{h \in G} b_h h \right) v = \left(\sum_{z \in G} \sum_{gh=z} a_g b_h z \right) v = \sum_{z \in G} \sum_{gh \in z} a_g b_h \sigma_z(v) = \sum_{g \in G} \sum_{h \in G} a_g b_h \sigma_g \sigma_h(v)$$

since $\sigma(gh) = \sigma(g)\sigma(h) = \sigma_g \sigma_h = \sigma_{gh}$ (σ homomorphism)

$$1v = \sigma(1)v = 1v = v$$

Conversely, assume V left kG -module.

If $g \in G$, then $v \mapsto gv$ defines $T_g : V \rightarrow V$. T_g nonsingular since $\exists T_g^{-1} = T_{g^{-1}}$

Define $\sigma : G \rightarrow GL(V)$

$$\sigma : g \mapsto T_g$$

σ k -representation

$$\sigma(gh) = T_{gh} = T_g T_h = \sigma(g)\sigma(h)$$

$$\sigma(gh)(v) = T_{gh}v = ghv = T_g T_h v = \sigma(g)\sigma(h)v \quad \forall v \in V$$

□

Proposition 7. Let group G , let $\sigma, \tau : G \rightarrow GL(V)$ be k -representations, field k .

If V^σ, V^τ corresponding kG -modules in Prop. 6 (Prop. 8.37 in Rotman (2010) [9]), then

$V^\sigma \simeq V^\tau$ as kG -modules iff \exists nonsingular $\varphi : V \rightarrow V$ s.t.

$$\varphi\tau(g) = \sigma(g)\varphi \quad \forall g \in G$$

Proof. If $\varphi : V^\tau \rightarrow V^\sigma$ kG -isomorphism, then $\varphi : V \rightarrow V$ isomorphism s.t.

$$\varphi(\sum a_g gv) = (\sum a_g g)\varphi(v) \quad \forall v \in V, \forall g \in G$$

in V^τ , $kG \times V \rightarrow V$

$$gv = \tau(g)(v)$$

in V^σ , $kG \times V \rightarrow V$ scalar multiplication

$$gv = \sigma(g)(v)$$

$$\implies \forall g \in G, v \in V, \quad \varphi(\tau(g)(v)) = \sigma(g)(\varphi(v))$$

I think

$$\varphi(gv) = \varphi(\tau(g)(v)) = g\varphi(v) = \sigma(g)\varphi(v)$$

$$\implies \varphi\tau(g) = \sigma(g)\varphi \quad \forall g \in G$$

Conversely, if \exists nonsingular $\varphi : V \rightarrow V$ s.t. $\varphi\tau(g) = \sigma(g)\varphi \quad \forall g \in G$

$$\varphi\tau(g)v = \varphi(\tau(g)v) = \sigma(g)\varphi(v) \quad \forall g \in G, \forall v \in V$$

Consider scalar multiplication

$$kG \times V \rightarrow V$$

$$\sum_{g \in G} a_g g(v) = \sum_{g \in G} a_g \tau_g(v)$$

$$\varphi \left(\sum_{g \in G} a_g \tau_g(v) \right) = \varphi \left(\sum_{g \in G} a_g \tau(g)v \right) = \sum_{g \in G} a_g \sigma(g) \sigma(g) \varphi(v) = \left(\sum_{g \in G} a_g g \right) \varphi(v)$$

□

Admittedly, after this exposition from Rotman (2010) [9], I still didn't understand how kG -modules relate to representation theory and group rings. I turned to Baker (2011) [10], which we'll do right now. Note that I found a lot of links to online resources on representation theory from Khovanov's webpage <http://www.math.columbia.edu/~khovanov/resource/>.

Note,

Definition 17. *vector subspace $W \subseteq V$ is called a*

G -submodule, G -subspace, EY : 20150922 “invariant” subspace?

if $\forall g \in G$, for representation $\rho : G \rightarrow GL_k(V)$, $\rho_g(w) \in W$, $\forall w \in W$, $\forall g \in G$ i.e. closed under “action of elements of G ” with $\rho_g =: \rho(g) : V \rightarrow V$

Given basis $\mathbf{v} = \{v_1 \dots v_n\}$ for V , $\dim_k V = n$, $\forall g \in G$,

$$\rho_g v_j = \rho(g) v_j = r_{kj}(g) v_k$$

for, indeed,

$$\rho_g x^j v_j = \rho(g) x^j v_j = x^j \rho(g) v_j = x^j r_{kj}(g) v_k = r_{kj} x^j v_k$$

so that

$$\rho : G \rightarrow GL_k(V)$$

$$\rho(g) = [r_{ij}(g)]$$

Example 2.1 (Baker (2011) [10]): Let $\rho : G \rightarrow GL_k(V)$ where $\dim_k V = 1$

$$\forall v \in V, v \neq 0, \forall g \in G, \lambda_g \in k \text{ s.t. } g \cdot v = \rho_g(v) = \lambda_g v$$

$$\rho(hg)v = \rho_h \rho_g v = \lambda_{hg} v = \lambda_h \lambda_g v \implies \lambda_{hg} = \lambda_h \lambda_g$$

$\implies \exists$ homomorphism $\Lambda : G \rightarrow k^\times$

$$\Lambda(g) = \lambda_g$$

From Sec. 2.2 “ G -homomorphisms and irreducible representations” of Baker (2011) [10], suppose $\rho : G \rightarrow GL_k(V)$ are 2 representations
 $\sigma : G \rightarrow GL_k(W)$

Many names for the same thing: G -equivalent, G -linear, G -homomorphism, EY : 20150922 kG -isomorphic?

If $\forall g \in G$,

$$\begin{array}{ccc} V & \xrightarrow{\varphi} & V \\ \tau_g \downarrow & & \downarrow \sigma_g \\ V & \xrightarrow{\varphi} & V \end{array} \iff V^\tau \xrightarrow{\varphi} V^\sigma$$

Indeed, define

$$\varphi : V^\tau \rightarrow V^\sigma$$

$$\varphi(v + w) = \varphi(v) + \varphi(w)$$

$$\varphi(rv) = \varphi\left(\sum_{g \in G} a_g g \cdot v\right) = \varphi\left(\sum_{g \in G} a_g \tau_g(v)\right) = \sum_{g \in G} a_g \varphi(\tau_g(v)) = \sum_{g \in G} a_g \sigma_g \cdot \varphi(v) = r \varphi(v)$$

EY : 20150922 So φ is a kG -isomorphism between left kG modules V^τ and V^σ if it's bijective and is “linear” in “scalars” $r \in kG$, i.e. $\varphi(rv) = r\varphi(v)$.

Define action of G on $\text{Hom}_k(V, W)$ ($\text{Hom}_k(V, W)$ is the vector space of k -linear transformations $V \rightarrow W$)

$$\begin{array}{ccc} V & \xrightarrow{f} & W \\ v & \xmapsto{f} & f(v) \end{array}$$

$$\forall f \in \text{Hom}_k(V, W), f : V \rightarrow W$$

$$f(v) \in W$$

Consider

$$G \times \text{Hom}_k(V, W) \rightarrow \text{Hom}_k(V, W)$$

$$(g \cdot f) \mapsto (\sigma_g f) \circ \rho_{g^{-1}} \text{ i.e. } (g \cdot f)(v) = \sigma_g f(\rho_{g^{-1}} v) \quad (f \in \text{Hom}_k(V, W))$$

Let $g, h \in G$,

$$(gh \cdot f)(v) = g \cdot \sigma_h f(\rho_{h^{-1}} v) = \sigma_g \sigma_h f \rho_{h^{-1}} \rho_{g^{-1}}(v) = (\sigma_{gh} f \rho_{(gh)^{-1}})(v)$$

Thus, $G \times \text{Hom}_k(V, W) \rightarrow \text{Hom}_k(V, W)$ is thus another G -representation of G .

$$(g \cdot f) \mapsto (\sigma_g f) \circ \rho_{g^{-1}}$$

For k -representation ρ , if the only G -subspaces of V are $\{0\}$, V , ρ **irreducible** or **simple**.

$$\rho_g(\{0\}) = \{0\}$$

$$\rho_g(V) = V$$

given subrepresentation $W \subseteq V$, V/W admits linear action of G , $\bar{\rho}_W : G \rightarrow GL_k(V/W)$ quotient representation

$$\bar{\rho}_W(g)(v + W) = \rho(g)(v) + W$$

if $v' - v \in W$

$$\rho(g)(v') + W = \rho(g)(v + (v' - v)) + W = (\rho(g)(v) + \rho(g)(v' - v)) + W = \rho(g)(v) + W$$

Proposition 8 (2.7 Baker (2011)[10]). *if $f : V \rightarrow W$ G -homomorphism, then*

(a) *$\ker f$ is G -subspace of V*

(b) *$\text{im} f$ is G -subspace of W*

$$\begin{array}{ccc} V & \xrightarrow{f} & W \\ \rho_g \downarrow & & \downarrow \sigma_g \\ V & \xrightarrow{f} & W \end{array}$$

Proof. Recall

(a) Let $v \in \ker f$. Then $\forall g \in G$,

$$f(\rho_g v) = \sigma_g f(v) = 0$$

so $\rho_g v \in \ker f$, $\forall g \in G$. So $\ker f$ is G -subspace of V

(b) Let $w \in \text{im} f$. So $w = f(u)$ for some $u \in V$

$$\sigma_g w = \sigma_g f(u) = f(\rho_g u) \in \text{im} f$$

So $\text{im} f$ is G -subspace of W

□

Theorem 11 (Schur's Lemma). *Let $\rho : G \rightarrow GL_{\mathbb{C}}(V)$ be irreducible representations of G over field $k = \mathbb{C}$; let $f : V \rightarrow W$ be*

$$\sigma : G \rightarrow GL_{\mathbb{C}}(W)$$

G -linear map.

- (a) if $f \neq 0$, f isomorphism. True $\forall k$ field, not just \mathbb{C}
 (b) if $V = W$, $\rho = \sigma$, then for some $\lambda \in \mathbb{C}$, f given by $f(v) = \lambda v$ ($v \in V$) (true for algebraically closed fields)

Proof. (a) By Prop. 8, $\ker f \subseteq V$, $\text{im} f \subseteq W$ are G -subspaces.
 For ρ , only G -subspaces are 0 or V , so if $\ker f = V$, $f = 0$. If $\ker f = 0$, f injective.
 For σ , only G -subspaces are 0 or V , so $\text{im} f = 0$, $f = 0$. If $\text{im} f = V$, f surjective.
 $\implies f$ isomorphism.
 (b) Let $\lambda \in \mathbb{C}$ be an eigenvalue of f , $f(v_0) = \lambda v_0$ eigenvector, $v_0 \neq 0$.
 Let linear $f_\lambda : V \rightarrow V$ s.t.

$$f_\lambda(v) = f(v) - \lambda v \quad (v \in V)$$

$$\forall g \in G$$

$$\rho_g f_\lambda(v) = \rho_g f(v) - \rho_g \lambda v = f(\rho_g v) - \lambda \rho_g v = f_\lambda(\rho_g v)$$

So f_λ is G -linear, for

$$\begin{array}{ccc} V & \xrightarrow{f} & V \\ \rho_g \downarrow & & \downarrow \rho_g \\ V & \xrightarrow{f} & V \end{array}$$

Since $f_\lambda(v_0) = 0$, by Prop. 8, $\ker f_\lambda = V$, (for $\ker f_\lambda \neq 0$ and so $\ker f_\lambda = V$)
 By rank-nullity theorem, $\dim V = \dim \ker f_\lambda + \dim \text{im} f_\lambda$.
 So $\text{im} f_\lambda = 0$, and so $f_\lambda(v) = 0$ ($\forall v \in V$) $\implies f(v) = \lambda v$

□

Schur's lemma, at least the first part, implies that the left kG -modules associated with representations ρ, σ are kG -isomorphic, i.e.

$$\begin{array}{ccc} V & \xrightarrow{f} & W \\ \rho_g \downarrow & & \downarrow \sigma_g \\ V & \xrightarrow{f} & W \end{array} \iff V^\rho \stackrel{f}{\simeq} V^\sigma$$

with f being an isomorphism between V^ρ and V^σ s.t.

$$\begin{aligned} f(v+w) &= f(v) + f(w) \quad \forall v, w \in (V^\sigma, +) \\ f(rv) &= rf(v) \quad \forall r = \sum_{g \in G} a_g g \in kG \end{aligned}$$

Kosmann-Schwarzbach's **Groups and Symmetries**[11] is a very lucid text that's mathematically rigorous enough and practical for physicists. It's really good and very clear. Let's follow its development for $SU(2)$, $SO(3)$, $SL(2, \mathbb{C})$ and corresponding Lie algebras $\mathfrak{su}(2)$, $\mathfrak{so}(3)$, $\mathfrak{sl}(2, \mathbb{C})$.

From Chapter 2 "Representations of Finite Groups" of Kosmann-Schwarzbach (2010) [11]

Definition 18 (2.1 Kosmann-Schwarzbach (2010)[11]). On $L^2(G)$, scalar product defined by

$$\langle f_1 | f_2 \rangle = \frac{1}{|G|} \sum_{g \in G} \overline{f_1(g)} f_2(g)$$

$f_1, f_2 \in \mathcal{F}(G) \equiv \mathbb{C}[G]$ vector space of functions on G taking values on \mathbb{C}

Definition 19 (2.3 Kosmann-Schwarzbach (2010)[11]). Let (E, ρ) be representation of G

character of $\rho \equiv \chi_\rho : G \rightarrow \mathbb{C}$

$$\chi_\rho(g) = \text{tr}(\rho(g)) = \sum_{i=1}^n (\rho(g))_{ii}$$

Note: equivalent representations have same character
 each conjugacy class of G , function χ_p is constant

Looking at Def. 18

$$\langle \chi_{\rho_1} | \chi_{\rho_2} \rangle = \frac{1}{|G|} \sum_{g \in G} \chi_{\rho_1}(g^{-1}) \chi_{\rho_2}(g)$$

since $\overline{\chi_{\rho_1(g)}} = \chi_{\rho_1}(g^{-1})$ by unitarity of representation with respect to scalar product \langle , \rangle

Proposition 9 (2.7 Kosmann-Schwarzbach (2010)[11]). Let (E_1, ρ_1) be representations of G , let linear $u : E_1 \rightarrow E_2$.
 (E_2, ρ_2)

Then \exists linear T_u s.t.

$$(19) \quad \begin{aligned} T_u &: E_1 \rightarrow E_2 \\ T_u &= \frac{1}{|G|} \sum_{g \in G} \rho_2(g) u \rho_1(g)^{-1} \end{aligned}$$

□

so that $\rho_2(g) T_u = T_u \rho_1(g) \quad \forall g \in G$

Proof.

$$\rho_2(g) T_u = \frac{1}{|G|} \sum_{h \in G} \rho_2(gh) u \rho_1(h^{-1}) = \frac{1}{|G|} \sum_{k \in G} \rho_2(k) u \rho_1(k^{-1}g) = T_u \rho_1(g)$$

□

Thus, diagrammatically, we have that

$$\begin{array}{ccccc} & & E_1 & \xrightarrow{T_u} & E_2 \\ & & \downarrow \rho_1(g) & & \downarrow \rho_2(g) \\ E_1 & \xrightarrow{u} & E_2 & \implies & E_1 & \xrightarrow{T_u} & E_2 \end{array}$$

From Definition 1.12 of Kosmann-Schwarzbach [11], "representations ρ_1 and ρ_2 are called **equivalent** if there is a bijective intertwining operator for ρ_1 and ρ_2 ." So I will interpret this as if an intertwining operator is not bijective, then the representations ρ_1, ρ_2 are not equivalent.

Proposition 10 (2.8 Kosmann-Schwarzbach (2010)[11]). Let (E_1, ρ_1) be irreducible representations of G , let linear $u : E_1 \rightarrow E_2$,
 (E_2, ρ_2)

define T_u by $T_u = \frac{1}{|G|} \sum_{g \in G} \rho_2(g) u \rho_1(g)^{-1}$ by Eq. 19.

- (i) If ρ_1, ρ_2 inequivalent, then $T_u = 0$

(ii) If $E_1 = E_2 = E$ and $\rho_1 = \rho_2 = \rho$, then

$$T_u = \frac{\text{tr}(u)}{\dim E} 1_E$$

Proof. (i) if ρ_1, ρ_2 are inequivalent, by definition, T_u is not isomorphic. Then by Schur's lemma (first part), $T_u = 0$
(ii) By Schur's lemma, $T_u(v) = \lambda v \quad \forall v \in E = E_1 = E_2$. So $T_u = \lambda 1_E$. $\text{tr} T_u = \lambda \dim E$ or $\lambda = \frac{\text{tr} T_u}{\dim E}$. Thus, $T_u = \frac{\text{tr} T_u}{\dim E} 1_E$ \square

Let $(e_1 \dots e_n)$ basis of E

$(f_1 \dots f_p)$ basis of F

$$\forall u \in \mathcal{L}(E, F), \quad \begin{aligned} u : E &\rightarrow F \\ u(x) &= u(x^j e_j) = x^j u(e_j) = x^j u^i_j f_i \quad \text{for } x = x^j e_j \in E \\ u &= u^i_j e^j \otimes f_i \quad \quad \quad y = y^i f_i \in F \end{aligned}$$

For

$$\begin{aligned} T : E^* \otimes F &\rightarrow \mathcal{L}(E, F) \\ T(\xi \otimes y) &= u^i_j e^j \otimes f_i \text{ i.e. set } T(\xi \otimes y) \text{ to this } u \\ T(\xi \otimes y) &= T(\xi_l e^l \otimes y^k f_k) = \xi_l y^k T(e^l \otimes f_k) = (\xi_l y^k T_{kj}^{li}) e^j \otimes f_i \implies \xi_l y^k T_{kj}^{li} = u^i_j \end{aligned}$$

Exercises. Exercises of Ch. 2 Representations of Finite Groups [11]

Exercise 2.6. [11] *The dual representation.*

Let (E, π) representation of group G .

$\forall g \in G, \xi \in E^*, x \in E$, set $\langle \pi^*(g)(\xi), x \rangle = \langle \xi, \pi(g^{-1})(x) \rangle$

(a) *dual* (or *contragredient*) of π , $\pi^* : G \rightarrow \text{End}(E^*)$, π^* is a representation, since

$$\begin{aligned} \langle \pi^*(gh)(\xi), x \rangle &= \langle \xi, \pi((gh)^{-1})(x) \rangle = \langle \xi, \pi(h^{-1}g^{-1})(x) \rangle = \langle \xi, \pi(h^{-1})\pi(g^{-1})(x) \rangle = \langle \xi, \pi(h^{-1})(\pi(g^{-1})(x)) \rangle = \\ &= \langle \pi^*(h)(\xi), \pi(g^{-1})(x) \rangle = \langle \pi^*(g)\pi^*(h)(\xi), x \rangle \end{aligned}$$

since this is true, $\forall x \in E, \forall \xi \in E^*, \pi^*(gh) = \pi^*(g)\pi^*(h)$.

dual π^* of π is a representation.

(b) Consider $G \times \mathcal{L}(E, F) \rightarrow \mathcal{L}(E, F)$.

$$g \cdot u = \rho(g) \circ u \circ \pi(g^{-1})$$

Define

$$\begin{aligned} \sigma : G &\rightarrow \text{End}(\mathcal{L}(E, F)) \\ \sigma(g) : \mathcal{L}(E, F) &\rightarrow \mathcal{L}(E, F) \\ \sigma(g)(u) &= \rho(g) \circ u \circ \pi(g^{-1}) \end{aligned}$$

Let $(e_1 \dots e_n)$ be a basis of E . Let $\xi = \xi_i e^i \in E^*, x = x^j e_j \in E$.

Consider the isomorphism $T : E^* \otimes F \rightarrow \mathcal{L}(E, F)$ defined as²

$$\begin{aligned} T : E^* \otimes F &\rightarrow \mathcal{L}(E, F) = \text{Hom}(E, F) \\ \xi \otimes y &\mapsto (x \mapsto \xi(x)y) \end{aligned}$$

Choose bases $(e_1 \dots e_n)$ of E
 $(e^1 \dots e^n)$ of E^* . Then
 $(f_1 \dots f_p)$ of F

$$T(e^j \otimes f_i)(x) = T(e^j \otimes f_i)(x^k e_k) = \delta^j_k x^k f_i = x^j f_i$$

$$T(e^j \otimes f_i)(e_k) = \delta^j_k f_i$$

Consider

$$u \in \mathcal{L}(E, F)$$

$$u : E \rightarrow F$$

$$u(x) = u(x^j e_j) = x^j u(e_j) = x^j u^i_j f_i$$

$$u(e_j) = u^i_j f_i \text{ i.e. } u : e_j \rightarrow u^i_j f_i$$

Then $\forall u \in \mathcal{L}(E, F)$,

$$T(u^i_j e^j \otimes f_i)(e_k) = u^i_j \delta^j_k f_i = u^i_k f_i = u(e_k) \implies u = T(u^i_j e^j \otimes f_i)$$

so T is surjective.

With $T(\xi \otimes y) = T(\xi' \otimes y')$,

$$T(\xi \otimes y)(x) = T(\xi' \otimes y')(x)$$

$$\xi(x)y = \xi'(x)y' \implies \xi(x)y - \xi'(x)y' = 0$$

which implies that $\xi \otimes y = \xi' \otimes y'$. So T is injective. Or, one could consider that $T^{-1} : \mathcal{L}(E, F) \rightarrow E^* \otimes F$, $T^{-1} : u \mapsto u^i_j e^j \otimes f_i$, which is the inverse of T .

Remark 1.

$$E^* \otimes F \xrightarrow{T} \mathcal{L}(E, F) = \text{Hom}(E, F)$$

$$(\xi, y) \mapsto (x \mapsto \xi(x)y)$$

and so $(e^j \otimes f_i) \mapsto (x \mapsto e^j(x)f_i = x^j f_i)$

So $E^* \otimes F$ is isomorphic to $\mathcal{L}(E, F) = \text{Hom}(E, F)$

For representation π ,

$$\pi : G \rightarrow \text{End}(E)$$

$$\pi(g) : E \rightarrow E$$

$$\pi(g)(x) = \pi(g)(x^j e_j) = x^j \pi(g)(e_j) = x^j \pi(g)^i_j e_i = (\pi(g)^i_j x^j e_i$$

Consider this matrix formulation:

$$\begin{aligned} \pi^*(g)(\xi) &= \pi^*(g)(\xi_i e^i) = \xi_i \pi^*(g)(e^i) = \xi_i (\pi^*(g))^i_j e^j \\ &\implies \langle \pi^*(g)(\xi), x \rangle = \xi_i (\pi^*(g))^i_j x^j \end{aligned}$$

and

$$\langle \xi, \pi(g^{-1})(x) \rangle = \xi_i \pi(g^{-1})^i_j x^j$$

so that

$$\langle \pi^*(g)(\xi), x \rangle = \langle \xi, \pi(g^{-1})(x) \rangle \implies \pi(g^{-1})^i_j = (\pi^*(g))^i_j$$

Thus, given a choice of basis for E , the *dual* of π , $\pi^*(g)^i_j$, and $\pi(g^{-1})^i_j$ are formally equal.

²Mathematics stackexchange Isomorphism between Hom and tensor product [duplicate] <http://math.stackexchange.com/questions/428185/isomorphism-between-hom-and-tensor-product>
<http://math.stackexchange.com/questions/57189/understanding-isomorphic-equivalences-of-tensor-product>

So for a choice of basis of E and of F ,

$$(\pi^* \otimes \rho)(g)(\xi, y) = (\pi^*(g) \otimes \rho(g))(\xi, y) = \pi^*(g)\xi \otimes \rho(g)y = \xi_l \pi(g^{-1})^l_j e^j \otimes \rho(g)^i_k y^k f_i = \rho(g)^i_k y^k \xi_l \pi(g^{-1})^l_j e^j \otimes f_i$$

Applying T ,

$$\begin{array}{ccc} E^* \otimes F & \xrightarrow{T} & \mathcal{L}(E, F) \\ \downarrow (\pi^* \otimes \rho)(g) & & \downarrow \sigma(g) \\ E^* \otimes F & \xrightarrow{T} & \mathcal{L}(E, F) \end{array} \quad \begin{array}{ccc} (\xi, y) & \xrightarrow{T} & (x \mapsto \xi(x)y) = y^i \xi_j \\ \downarrow (\pi^* \otimes \rho)(g) & & \downarrow \sigma(g) \\ \pi^*(g)(\xi) \otimes \rho(g)y & \xrightarrow{T} & \rho(g)y^i \xi_j \pi(g^{-1}) = \rho(g)T(\xi, y)\pi(g^{-1}) \end{array}$$

Thus

Thus, representation $\sigma(g)$ is equivalent to representation $(\pi^* \otimes \rho)$, a tensor product of representations.

Exercise 2.15. Representation of $GL(2, \mathbb{C})$ on the polynomials of degree 2

Let group G , let representation ρ of G on $V = \mathbb{C}^n$, i.e. $\rho : G \rightarrow \text{End}(V)$

Let $P^{(k)}(V)$ vector space of complex polynomials on V that are homogeneous of degree k .

For $f \in P^{(k)}(V)$, the general form is

$$f = \sum_{\substack{i_1+i_2+\dots+i_n=k \\ 0 \leq i_j \leq k}} a_{i_1 i_2 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}$$

Given

$$\binom{n+k}{k} = \binom{k-1}{k-1} + \binom{k}{k-1} + \dots + \binom{n+k-1}{k-1} = \sum_{i=0}^n \binom{k-1+i}{k-1}$$

$\binom{k+n-1}{n-1}$ is number of monomials of degree k .

So $\dim P^{(k)}(V) = \binom{k+n-1}{n-1}$. This is a very lucid and elementary exposition on the basics of polynomials which I found was useful for the basic facts I forgot³.

So we have the graded algebra

$$P(V) = \bigoplus_{k=0}^{\infty} P^{(k)}(V)$$

$$\rho^{(k)} : G \rightarrow \text{End}(P^{(k)}(V))$$

$$\rho^{(k)}(g) : P^{(k)}(V) \rightarrow P^{(k)}(V)$$

$$\rho^{(k)}(g)(f) = f \circ \rho(g^{-1})$$

This is a representation of G since

(a)

$$\begin{aligned} \rho^{(k)}(gh)(f) &= f \circ \rho((gh)^{-1}) = f \circ \rho(h^{-1}g^{-1}) = f \circ \rho(h^{-1}\rho(g^{-1})) \\ \rho^{(k)}(g)\rho^{(k)}(h)(f) &= \rho^{(k)}(g)(f \circ \rho(h^{-1})) = f \circ \rho(h^{-1}) \circ \rho(g^{-1}) \implies \rho^{(k)}(gh) = \rho^{(k)}(g)\rho^{(k)}(h) \end{aligned}$$

(b) Choose basis $(e_1 \dots e_n)$ of V , $x = x^j e_j \in V$, $\rho : G \rightarrow \text{End}(V)$, and so $\rho(g)(x) = \rho(g)(x^j e_j) = x^j \rho(g)(e_j) = x^j (\rho(g))^i_j e_i$.

$$\text{With } \xi(e_i) = \xi_i \implies \langle \xi, \rho(g^{-1})x \rangle = \xi_i x^j (\rho(g^{-1}))^i_j$$

$$\forall \xi \in V^*, \xi = \xi_i e^i,$$

$$\begin{aligned} \rho^*(g)(\xi) &= \rho^*(g)(\xi_i e^i) = \xi_i \rho^*(g)^i_j e^j \\ \implies \langle \rho^*(g)(\xi), x \rangle &= \xi_i x^j (\rho^*(g))^i_j \implies (\rho^*(g))^i_j = (\rho(g^{-1}))^i_j \end{aligned}$$

$$\begin{aligned} \text{So } \forall f \in P^{(1)}(V), x \in V, \rho(g^{-1})x &= x^j (\rho(g^{-1}))^i_j e_i. \text{ So } f \circ \rho(g^{-1})(x) = \sum_{i=1}^n a_i (\rho(g^{-1}))^i_j x^j = \sum_{i=1}^n a_i (\rho^*(g))^i_j x^j \\ \implies \rho^{(1)}(g)(f) &= f \circ \rho^*(g) \end{aligned}$$

(c) Suppose $G = GL(2, \mathbb{C})$, $V = \mathbb{C}^2$, ρ fundamental representation $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $g^{-1} = \frac{1}{\det g} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ for $\det g = ad - bc$.

$$\text{Let } k = 2, \dim P^{(2)}(\mathbb{C}^2) = \binom{2+2-1}{2-1} = \binom{3}{1} = 3$$

$$\forall f \in P^{(2)}(\mathbb{C}^2), f(x, y) = Ax^2 + 2Bxy + Cy^2$$

Let

$$P^{(2)}(\mathbb{C}^2) \rightarrow \mathbb{C}^3$$

$$f(x, y) = Ax^2 + 2Bxy + Cy^2 \mapsto \begin{pmatrix} A \\ B \\ C \end{pmatrix} \in \mathbb{C}^3$$

Call this transformation T , $T : P^{(2)}(\mathbb{C}^2) \rightarrow \mathbb{C}^3$.

$$\forall \begin{pmatrix} A \\ B \\ C \end{pmatrix} \in \mathbb{C}^3, f(x, y) = Ax^2 + 2Bxy + Cy^2 \text{ and } Tf(x, y) = \begin{pmatrix} A \\ B \\ C \end{pmatrix}. \text{ } T \text{ surjective.}$$

Suppose $Tf(x, y) = Tf'(x, y)$,

$$\implies Ax^2 + 2Bxy + Cy^2 = A'x^2 + 2B'xy + C'y^2$$

$$\implies (A - A')x^2 + 2(B - B')xy + (C - C')y^2 = 0$$

Then since the monomials form a basis, and its basis elements are independent (by definition), then $A = A'$, $B = B'$, $C = C'$. T injective. So T is bijective, an isomorphism.

(This is all in **groups.sage**)

```
sage: P2CC.<x,y> = PolynomialRing(CC,2) # this declares a PolynomialRing of field of complex numbers,
# of order 2 (i.e. only 2 variables for a polynomial, such as x, y)
sage: A = var('A')
sage: assume(A, 'complex')
sage: B = var('B')
sage: assume(B, 'complex')
sage: C = var('C')
sage: assume(C, 'complex')
sage: f(x,y) = A*x**2 + 2*B*x*y + C*y**2

sage: a = var('a')
sage: assume(a, 'complex')
sage: b = var('b')
sage: assume(b, 'complex')
sage: c = var('c')
sage: assume(c, 'complex')
sage: d = var('d')
sage: assume(d, 'complex')
sage: g = Matrix([[a,b],[c,d]])
sage: X = Matrix([[x],[y]])
sage: f((g.inverse()*X)[0,0], (g.inverse()*X)[1,0]).expand()
sage: f((g.inverse()*X)[0,0], (g.inverse()*X)[1,0]).expand().coefficient(x^2).full_simplify()
(C*c^2 - 2*B*c*d + A*d^2)/(b^2*c^2 - 2*a*b*c*d + a^2*d^2)
sage: f((g.inverse()*X)[0,0], (g.inverse()*X)[1,0]).expand().coefficient(x*y).full_simplify()
-2*(C*a*c + A*b*d - (b*c + a*d)*B)/(b^2*c^2 - 2*a*b*c*d + a^2*d^2)
sage: f((g.inverse()*X)[0,0], (g.inverse()*X)[1,0]).expand().coefficient(y^2).full_simplify()
(C*a^2 - 2*B*a*b + A*b^2)/(b^2*c^2 - 2*a*b*c*d + a^2*d^2)
```

³Polynomials. Math 4800/6080 Project Course <http://www.math.utah.edu/~bertram/4800/PolyIntroduction.pdf>

So

$$\begin{aligned} \rho^{(2)}(g)(f)(x, y) &= f \circ \rho(g^{-1})(x, y) = \\ &= \frac{Cc^2 - 2Bcd + Ad^2}{(ad - bc)^2} x^2 + -2 \frac{(Cac + Abd - (bc + ad)B)}{(ad - bc)^2} xy + \frac{Ca^2 - 2Bab + Ab^2}{(ad - bc)^2} y^2 \end{aligned}$$

So define $\tilde{\rho}: G \rightarrow \text{End}(\mathbb{C}^3)$. $\tilde{\rho}$ is a representation, for

$$\forall v = \begin{pmatrix} A \\ B \\ C \end{pmatrix} \in \mathbb{C}^3, \quad \tilde{\rho}(gh)(v) = T \circ f \circ \rho((gh)^{-1}) = T \circ f \circ \rho(h^{-1}g^{-1}) = T \circ f \circ \rho(h^{-1})\rho(g^{-1})$$

$$\begin{aligned} \text{Now } \tilde{\rho}(h)(v) &= T \circ f \circ \rho(h^{-1}) \\ \implies \tilde{\rho}(g)\tilde{\rho}(h)(v) &= T \circ (f \circ \rho(h^{-1})) \circ \rho(g^{-1}) = T \circ f \circ \rho(h^{-1})\rho(g^{-1}) \text{ and so} \\ \tilde{\rho}(gh) &= \tilde{\rho}(g)\tilde{\rho}(h) \end{aligned}$$

And so

$$\tilde{\rho}^*(g)(v) = Tf\rho(g^{-1})$$

and consider this commutation diagram, that (helped me at least and) clarifies the relationships:

$$\begin{array}{ccc} P^{(2)}(\mathbb{C}^2) & \xrightarrow{T} & \mathbb{C}^3 \\ \rho^{(2)}(g) \downarrow & & \downarrow \tilde{\rho}(g) \\ P^{(2)}(\mathbb{C}^2) & \xrightarrow{T} & \mathbb{C}^3 \end{array} \quad \begin{array}{ccc} f & \xrightarrow{T} & \begin{pmatrix} A \\ B \\ C \end{pmatrix} \\ \rho^{(2)}(g) \downarrow & & \downarrow \tilde{\rho}(g) \\ f \circ \rho(g^{-1}) & \xrightarrow{T} & \begin{pmatrix} D \\ E \\ F \end{pmatrix} \end{array}$$

with

$$\begin{pmatrix} D \\ E \\ F \end{pmatrix} = \begin{pmatrix} \frac{Cc^2 - 2Bcd + Ad^2}{(ad - bc)^2} \\ -2 \frac{(Cac + Abd - (bc + ad)B)}{(ad - bc)^2} \\ \frac{Ca^2 - 2Bab + Ab^2}{(ad - bc)^2} \end{pmatrix}$$

Now define the dual $\tilde{\rho}^*$ as such:

$$\begin{aligned} \tilde{\rho}^*(g) : (\mathbb{C}^3)^* &\rightarrow (\mathbb{C}^3)^* \\ \tilde{\rho}^*(g) &= \tilde{\rho}(g^{-1}) \\ \forall \xi &\in (\mathbb{C}^3)^* \\ \tilde{\rho}^*(g)\xi &= \xi_i(\tilde{\rho}^*(g))^i_j e^j = \xi_i(\tilde{\rho}(g^{-1}))^i_j e^j \end{aligned}$$

$$\text{So for } v = \begin{pmatrix} A \\ B \\ C \end{pmatrix} \in \mathbb{C}^3, f = T^{-1}v = Ax^2 + 2Bxy + Cy^2 \in P^2(\mathbb{C}^2),$$

$$\tilde{\rho}(g^{-1})(v) = T \circ (f\rho(g)) = \begin{bmatrix} Aa^2 + 2Bac + Cc^2 \\ Aab + Bbc + Bad + Ccd \\ Ab^2 + 2Bbd + Cd^2 \end{bmatrix}$$

which was found using Sage Math:

```
sage: f((g*X)[0,0],(g*X)[1,0])
(a*x + b*y)^2*A + 2*(a*x + b*y)*(c*x + d*y)*B + (c*x + d*y)^2*C
sage: f((g*X)[0,0],(g*X)[1,0]).expand()
A*a^2*x^2 + 2*B*a*c*x^2 + C*c^2*x^2 + 2*A*a*b*x*y + 2*B*b*c*x*y + 2*B*a*d*x*y + 2*C*c*d*x*y + A*b^2*y^2 + 2*B*b*d*y^2 + C*d^2*y^2
sage: f((g*X)[0,0],(g*X)[1,0]).expand().coefficient(x^2)
A*a^2 + 2*B*a*c + C*c^2
sage: f((g*X)[0,0],(g*X)[1,0]).expand().coefficient(x*y)
2*A*a*b + 2*B*b*c + 2*B*a*d + 2*C*c*d
sage: f((g*X)[0,0],(g*X)[1,0]).expand().coefficient(y^2)
A*b^2 + 2*B*b*d + C*d^2
```

or

```
sage: T( f((g*X)[0,0],(g*X)[1,0]).expand() )
[A*a^2 + 2*B*a*c + C*c^2,
 2*A*a*b + 2*B*b*c + 2*B*a*d + 2*C*c*d,
 A*b^2 + 2*B*b*d + C*d^2]
```

So then

$$\tilde{\rho}(g^{-1}) = \begin{bmatrix} a^2 & 2ac & c^2 \\ 2ab & 2(ad + bc) & 2cd \\ b^2 & 2bd & d^2 \end{bmatrix}$$

So then

$$\tilde{\rho}^*(g) = \begin{bmatrix} a^2 & 2ac & c^2 \\ 2ab & 2(ad + bc) & 2cd \\ b^2 & 2bd & d^2 \end{bmatrix}$$

and operate on row vectors $\xi \in (\mathbb{C}^3)^*$ with $\tilde{\rho}^*(g)$ from the row vector's right.

More: Let $G = SU(2)$. Then $U = e^{i\phi} \begin{bmatrix} a & b \\ -\bar{b} & \bar{a} \end{bmatrix}$

$$\tilde{\rho}: SU(2) \rightarrow \text{End}(\mathbb{C}^3)$$

$$\tilde{\rho}(U): \mathbb{C}^3 \rightarrow \mathbb{C}^3$$

$$\tilde{\rho}(U)(v) = e^{-2i\varphi} \begin{bmatrix} A\bar{a}^2 + 2B\bar{a}\bar{b} + C\bar{b}^2 \\ -A\bar{a}\bar{b} + B + C\bar{a}\bar{b} \\ Ab^2 - 2Bab + Ca^2 \end{bmatrix}$$

$$\implies \tilde{\rho}(U) = e^{-2i\varphi} \begin{bmatrix} -\bar{a}^2 & 2\bar{a}\bar{b} & \bar{b}^2 \\ -\bar{a}\bar{b} & 1 & \bar{a}\bar{b} \\ b^2 & -2ab & a^2 \end{bmatrix}$$

cf. Ch. 5 Lie Groups of Jeffrey Lee (2009) [2]

Definition 20 (Lie Group). ***Lie Group** $G := \text{smooth manifold } G \text{ is a } \mathbf{Lie \ Group}$ if G is a group (abstract group), s.t.*

multiplication map $\mu: G \times G \rightarrow G$

$$\mu(g, h) = gh$$

inverse map $\text{inv}: G \rightarrow G$

$$\text{inv}(g) = g^{-1}$$

are C^∞ maps.

If group is abelian, use additive notation $g + h$ for group operation.

Definition 21 ($GL(n, \mathbb{R})$). *$GL(n, \mathbb{R}) := \text{group of all invertible real } n \times n \text{ matrices.}$*

global chart on $GL(n, \mathbb{R}) = \{x_j^i\}$, n^2 functions x_j^i , where if $A \in GL(n, \mathbb{R})$, then $x_j^i(A)$ is ij th entry of A .

Claim: $GL(n, \mathbb{R})$ is a Lie group.

Proof. multiplication is clearly smooth: $(AB)_{ij} = A_{ik}B_{kj}$,

$$\frac{\partial}{\partial x_m^i}(x_k^i(A)x_j^k(B)) = \delta_l^i \delta_k^m x_j^k(B) + x_k^i(A)\delta_l^k \delta_j^m$$

inversion map; appeal to formula for A^{-1} , $A^{-1} = \text{adj}(A)/\det(A)$, $\text{adj}(A) \equiv$ adjoint matrix (whose entries are cofactors).

$\implies A^{-1}$ depends smoothly on entries of A .

Similarly, $GL(n, \mathbb{C})$, group of invertible $n \times n$ complex matrices, is a Lie group.

Exercise 5.5. Let subgroup H of G , consider cosets gH , $g \in G$.

Recall G is disjoint union of cosets of H .

Claim: if H open, so are all its cosets. And H closed.

Proof. cf. [stackexchange: Open subgroups of a topological group are closed](#)

$gH = \{gh|h \in H\}$ is an open neighborhood of g (since $1 \in H$, and mapping $h \mapsto gh$ sends open sets to open sets, since its inverse, $gh \mapsto h$, is C^∞ (so continuous)).

$$\begin{array}{ll} gH \rightarrow H & H \rightarrow gH \\ gh \xrightarrow{g^{-1}} h = \mu(g^{-1}, gh) & h \xrightarrow{g} gh = \mu(g, h) \end{array}$$

Then \forall coset gH , gH is open.

Suppose $g' \in H^c \equiv G - H \equiv G \setminus H$.

Consider $h \in H$, if $g'h \in H$, then $g' = (g'h)h^{-1} \in H$ (recall $h^{-1} \in H$, and H is a subgroup).

Contradiction.

$\implies \forall g' \in H^c$, \exists open neighborhood $g'H \subset H^c$, so H^c open (by definition). Then H closed.

cf. Thm. 5.6 in Jeffrey Lee (2009) [2].

Theorem 12. *If G connected Lie group, U neighborhood of identity element e , then U generates the group, i.e. $\forall g \in G$, g is a product of elements of U .*

Proof. Note $V = \text{inv}(U) \cap U$ is an open neighborhood of e . Note $\text{inv}(V) = V$. $\text{inv}(V) \equiv V^{-1} = \{V^{-1}|v \in V\}$. We say that V is *symmetric*.

Claim: V generates G .

\forall open W_1 , open $W_2 \subset G$,

$W_1W_2 = \{w_1w_2|w_1 \in W_1, w_2 \in W_2\}$ is an open set being a union of open sets $\bigcup_{g \in W_1} gW_2$.

Thus, inductively defined sets

$$V^n = VV^{n-1}, \quad n = 1, 2, 3, \dots$$

are open.

$$e \in V \subset V^2 \subset \dots V^n \subset \dots$$

It's easy to check that each V^n is symmetric.

$$\text{inv}(V) = V$$

$$\text{inv}(V^2) = \text{inv}\left(\bigcup_{v \in V} vV\right) = V\text{inv}(V) = V = V^2$$

$$\text{inv}(V^{n+1}) = \text{inv}\left(\bigcup_{v \in V} vV^n\right) = V\text{inv}(V^n) = VV^n = V^{n+1}$$

so $V^\infty := \bigcup_{n=1}^\infty V^n$ is symmetric.

V^∞ closed under inversion, also multiplication. Thus V^∞ is an open subgroup.

From Exercise 5.5, Jeffrey Lee (2009) [2], i.e. Exercise **7**, V^∞ also closed, since G is connected, $V^\infty = G$. (a topological space X is **connected** iff the only open and closed (clopen) sets are \emptyset and X).

□

Definition 22. *Identity component of G, G_0 .*

$G_0 :=$ connected component of Lie group G that contains identity;

□ G_0 is a Lie group, and is generated by any open neighborhood of the identity.

Definition 23. *For Lie group G , fixed element $g \in G$,*

left translation (by g) $L_g : G \rightarrow G$, $L_gx = gx$, $\forall x \in G$

right translation (by g) $R_g : G \rightarrow G$, $R_gx = xg$, $\forall x \in G$

L_g, R_g are diffeomorphisms with $L_g^{-1} = L_{g^{-1}}, R_g^{-1} = R_{g^{-1}}$.

Definition 24 (Product Lie group). *If G, H are Lie groups, then product manifold $G \times H$ is a Lie group, where multiplication*

$$(g_1, h_1) \cdot (g_2, h_2) = (g_1g_2, h_1h_2)$$

*Lie group $G \times H$ is called **product Lie group***

e.g. product group $S^1 \times S^1 \equiv$ 2-torus group.

Generally, higher torus groups $T^n = S^1 \times \dots \times S^1$ (n factors).

Definition 25 (Lie subgroup of G, H). *Let H be an abstract subgroup of Lie group G .*

If H is a Lie group s.t. inclusion map $i : H \rightarrow G \equiv H \hookrightarrow G$ is an immersion, then H is a

□ ***Lie subgroup*** of G .

Recall $i : H \rightarrow G$ immersion iff Di injective, i.e. iff $\text{rank}Di = \dim H$

cf. Prop. 5.9 in Jeffrey Lee (2009) [2].

Proposition 11. *If H abstract subgroup of Lie group G , that's also a regular submanifold \equiv embedded submanifold, then H closed Lie subgroup.*

Recall that

$$\text{embedded submanifold} \equiv \text{regular submanifold}$$

Each name is used frequently and we shouldn't be biased against one or the other; we'll have to refer to both, to emphasize they're *exactly the same*.

embedded submanifold \equiv regular submanifold is an immersed submanifold s.t. inclusion map i is a topological embedding,

i.e. embedded submanifold \equiv regular submanifold $S \subset M$,

immersed submanifold S if $i : S \rightarrow M \equiv S \hookrightarrow M$ is an immersion, i.e. Di injective, i.e. $\text{rank}Di \equiv \dim S$.

topological embedding $:=$ homeomorphism onto its image, i.e.

injective cont. map $f : X \rightarrow Y$, X, Y topological spaces, is a **topological embedding**

if f is a homeomorphism between X and $f(X)$.

f homeomorphism is a bijection, continuous, and f^{-1} continuous.

e.g. \forall embedding $f : M \rightarrow N$, $f(M) \subset N$ naturally has the structure of an embedding submanifold \equiv regular submanifold.

Useful, intrinsic definition of **embedded submanifold** \equiv regular submanifold.

Let manifold M , $\dim M = n$, let $k \in \mathbb{Z}^+$, s.t. $0 \leq k \leq n$.

A k -dim. embedded submanifold \equiv regular submanifold S is subset $S \subset M$ s.t. $\forall p \in S$, \exists chart $(U \subset M, \varphi : U \rightarrow \mathbb{R}^n \ni 0)$, s.t. $\varphi(S \cap U)$ is the intersection of a k -dim. plane with $\varphi(U)$.

(pairs $(S \cap U, \varphi|_{S \cap U})$ form an atlas for differential structure on S).

Proof 1:

Proof. H subgroup of G , so
multiplication map $H \times H \rightarrow H$
inversion map $H \rightarrow H$
are restrictions of multiplication and inversion maps on G .

Since H regular submanifold, maps are smooth.
Recall H regular submanifold iff H immersive submanifold (i.e. $H \hookrightarrow G$ is an immersion) and H topological subspace of G , i.e. submanifold topology on H is same as subspace topology.

Claim: H closed.
Let $x_0 \in \overline{H}$
Let (U, x) be a chart adapted to H , whose domain contains e .
Let

$$\delta : G \times G \rightarrow G$$

$$\delta(g_1, g_2) = g_1^{-1}g_2$$

Choose open set V s.t. $e \in V \subset \overline{V} \subset U$.
By continuity map δ , find open neighborhood O of identity e s.t. $O \times O \subset \delta^{-1}(V)$
If $\{h_i\}$ sequence in H converging to $x_0 \in \overline{H}$, then $x_0^{-1}h_i \rightarrow e$ and $x_0^{-1}h_i \in O$ for all sufficiently large i .
Since $h_j^{-1}h_i = (x_0^{-1}h_j)^{-1}x_0^{-1}h_i$, $h_j^{-1}h_i \in V$ for sufficiently large i, j .

For any sufficiently large fixed j ,

$$\lim_{i \rightarrow 0} h_j^{-1}h_i = h_j^{-1}x_0 \in \overline{V} \subset U$$

Since U is domain of a single-slice chart, $U \cap H$ closed in U .

Thus, since $\forall h_j^{-1}h_i \in U \cap H$, $h_j^{-1}x_0 \in U \cap H \subset H$, \forall sufficiently large j .
 $\implies x_0 \in H$, and since x_0 arbitrary, done.

Proof 2:
cf. 9.2 The Closed Subgroup Theorem I of 427 Notes⁴

Proof. **Claim:** Since H is an embedded submanifold \equiv regular submanifold, \exists neighborhood U of 1 , $1 \in G$, s.t. $U \cap H$ closed in U .

Let $x_0 \in \overline{H}$, $\overline{H} \equiv$ closure of x_0 .
Then $x_0U^{-1} \subseteq G$ is a neighborhood of x_0 in G (since $1 \in U^{-1}$, $x_01 = x_0 \in x_0U^{-1}$)

$$\implies x_0U^{-1} \cap H \neq \emptyset$$

$\forall x \in x_0U^{-1} \cap H$, $x = x_0U^{-1}$ for some $u \in U$. Thus, $x^{-1}x_0 = u \in U$.

Now
 $L_{x^{-1}} : G \rightarrow G$ is a homeomorphism, so $L_{x^{-1}}(H) = H$. By continuity, $L_{x^{-1}}(\overline{H}) = \overline{H}$. Thus $x^{-1}x_0 \in \overline{H}$.

Claim: $x^{-1}x_0 \in H \cap U$.
Since $x^{-1}x_0 \in \overline{H} \cap U$, \exists sequence $\{h_i\} \subset H \cap U$ s.t. $h_0 \rightarrow x^{-1}x_0$.
But recall $H \cap U$ closed in U , so $x^{-1}x_0 \in H \cap U$.

$$\implies x_0 \in xH = H, \quad \overline{H} \subseteq H$$

Thus H closed.

⁴<https://faculty.math.illinois.edu/~lerman/519/s12/427notes.pdf>

Claim: If H abstract subgroup of Lie group G , that's also an embedded submanifold \equiv regular submanifold, then H is a Lie subgroup.

Recall that by definition, Lie group has group multiplication and inverse map to be C^∞ . Then, just show group multiplication is C^∞ , first.

Since G is a Lie group, then

$$\mu : G \times G \rightarrow G$$

$$\mu(x, y) = xy$$

is C^∞ (by definition).

Then $\mu : G \times G \rightarrow G$ cont.
Consider subgroup $H \subseteq G$ and $\mu : H \times H \rightarrow H$.
Since $H \times H \subseteq G \times G$, $\forall (x, y) \in H \times H$ (fix $(x, y) \in H \times H$), \forall neighborhood V of $\mu(x, y) = xy$, $V \subset G$, \exists neighborhood U of (x, y) s.t. $\mu(U) \subseteq V$ (by $\mu : G \times G \rightarrow G$ cont.).
Since H embedded submanifold \equiv regular submanifold of G ,
 \exists neighborhood $V' \subseteq V$ of $xy \in G$, coordinate map $\varphi : V' \rightarrow \mathbb{R}^n$ ($n = \dim G$) s.t.

$$\varphi(H \cap V') = \varphi(V') \cap (\mathbb{R}^k \times \{0\})$$

where $k = \dim H$
(since H is a k -dim. embedded submanifold \equiv regular submanifold, $H \subseteq G$, s.t. $\forall p \in H$, \exists chart $(V \subset G, \varphi : U \rightarrow \mathbb{R}^n \ni 0)$, s.t. $\varphi(U \cap V) = \varphi(V) \cap (\mathbb{R}^k \times \{0\})$).
Now

$$\varphi \circ \mu : \mu^{-1}(V') \cap U \rightarrow \mathbb{R}^n \text{ is } C^\infty, \text{ and } \varphi \circ \mu(\mu^{-1}(V') \cap U) \subseteq \mathbb{R}^k \times \{0\}$$

Let projection $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^k$ be the standard projection,

$$\pi \circ \varphi \circ \mu : \mu^{-1}(V') \cap U \rightarrow \mathbb{R}^k \text{ is } C^\infty$$

$\implies \mu$ is C^∞

From Chapter 4 “Lie Groups and Lie Algebras” of Kosmann-Schwarzbach (2010) [11]
While Proposition 2.6 of Kosmann-Schwarzbach (2010) [11] states that

$$\det(\exp(X)) = \exp(\operatorname{tr} X)$$

here are some other resources online that gave further discussion on the characteristic polynomial, $\det(A - \lambda 1)$ and the different terms of it, called Newton identities:

- http://scipp.ucsc.edu/~haber/ph116A/charpoly_11.pdf
- <http://math.stackexchange.com/questions/1126114/how-to-find-this-lie-algebra-proof-that-mathfraksl-is-tr>
- <http://mathoverflow.net/questions/131746/derivative-of-a-determinant-of-a-matrix-field>

Theorem 13 (5.1 [11]). *Consider $\mathfrak{g} = \{X = \gamma'(0) | \gamma : 1 \rightarrow G \text{ of class } C^1, \gamma(0) = 1\}$
Let Lie group G*

- \mathfrak{g} vector subspace of $\mathfrak{gl}(n, \mathbb{R})$
- $X \in \mathfrak{g}$ iff $\forall t \in \mathbb{R}, \exp(tX) \in G$
- if $X \in \mathfrak{g}$, if $g \in G$, then $gXg^{-1} \in \mathfrak{g}$
- \mathfrak{g} closed under matrix commutator, i.e. if $X, Y \in \mathfrak{g}$, $[X, Y] \in \mathfrak{g}$

Proof. (i)

- (ii) If $\exp(tX) \in G$, then $X \frac{d}{dt} \exp(tX) \Big|_{t=0} \in \mathfrak{g}$ (by def.)
 If $X \in \mathfrak{g}$, then by def., $X = \frac{d}{dt} \gamma(t) \Big|_{t=0}$ with $\gamma(t) \in G$.
 Now Taylor expand; $\forall k \in \mathbb{Z}^+$

$$\begin{aligned} \gamma\left(\frac{t}{k}\right) &= 1 + \frac{t}{k}X + O\left(\frac{1}{k^2}\right) = \exp\left(\frac{t}{k}X + O\left(\frac{1}{k^2}\right)\right) \\ &\implies \left(\gamma\left(\frac{t}{k}\right)\right)^k = \exp(tX) \\ \gamma\left(\frac{t}{k}\right) &\in G \quad \forall k \in \mathbb{Z}^+ \end{aligned}$$

G closed subgroup, so $\lim_{k \rightarrow \infty} (\gamma(\frac{t}{k}))^k = \exp(tX) \in G$

- (iii)
(iv)

Definition 26 (Lie algebra). *Lie algebra \mathfrak{g} , tangent space to G at 1, i.e. $\mathfrak{g} := T_1G$ is called Lie algebra of Lie group G .*

$$\mathfrak{g} := \{X = \gamma'(0) | \gamma : 1 \rightarrow G \text{ of class } C^1, \gamma(0) = 1\} = T_1G$$

This is based on Proposition 5.3 of Kosmann-Schwarzbach (2010) [11].

For Lie group

$$U(n) = \{U \in GL(n, \mathbb{C}) | UU^\dagger = 1\}$$

If $X \in \mathfrak{u}(n)$, then $\exp(tX) \in U(n)$. Then

$$\exp(tX) \exp(tX)^\dagger = (1 + tX + O(t^2))(1 + tX^\dagger + O(t^2)) = 1 + t(X + X^\dagger) + O(t^2) = 1 \forall t \in \mathbb{R} \implies X + X^\dagger = 0$$

i.e. $X \in \mathfrak{u}(n)$ is an anti-Hermitian complex $n \times n$ matrix.

$$\mathfrak{u}(n) = \{X \in \mathfrak{gl}(n, \mathbb{C}) | X + X^\dagger = 0\}$$

Physicists: $X = iA$ and so $A - A^\dagger$. $A \in \mathfrak{u}(n)$ is a Hermitian complex $n \times n$ matrix.

$$\mathfrak{u}(n) = \{A \in \mathfrak{gl}(n, \mathbb{C}) | A - A^\dagger = 0\}$$

Regardless, $\dim_{\mathbb{R}} \mathfrak{u}(n) = n^2 = 2n^2 - n^2$

For Lie group

$$SU(n) = \{U \in GL(n, \mathbb{C}) | UU^\dagger = 1, \det U = 1\}$$

Then

$$\mathfrak{su}(n) = \{X \in \mathfrak{gl}(n, \mathbb{C}) | X + X^\dagger = 1, \operatorname{tr} X = 0\}$$

is the Lie algebra of traceless anti-Hermitian complex $n \times n$ matrices, and that

$$\dim_{\mathbb{R}} \mathfrak{su}(n) = n^2 - 1$$

In summary,

$$\begin{array}{ccc} \mathfrak{u}(n) = \{X \in \mathfrak{gl}(n, \mathbb{C}) | X + X^\dagger = 0\} & & \mathfrak{su}(n) = \{X \in \mathfrak{gl}(n, \mathbb{C}) | X + X^\dagger = 0, \operatorname{tr} X = 0\} \\ \exp(tX) \downarrow & & \exp(tX) \downarrow \\ U(n) = \{U \in GL(n, \mathbb{C}) | UU^\dagger = 1\} & & SU(n) = \{U \in GL(n, \mathbb{C}) | UU^\dagger = 1, \det U = 1\} \end{array}$$

$$\dim_{\mathbb{R}} \mathfrak{u}(n) = n^2 \quad \dim_{\mathbb{R}} \mathfrak{su}(n) = n^2 - 1$$

From Chapter 5 “Lie Groups $SU(2)$ and $SO(3)$ ” of Kosmann-Schwarzbach (2010) [11],

7.0.1. *Bases of $\mathfrak{su}(2)$, Subsection 1.1 of Chapter 5 of Kosmann-Schwarzbach (2010) [11].* Recall that

$$\mathfrak{su}(n) = \{X \in \mathfrak{gl}(n, \mathbb{C}) | X + X^\dagger = 0, \operatorname{tr} X = 0\}$$

$$\exp(tX) \downarrow$$

$$SU(n) = \{U \in GL(n, \mathbb{C}) | UU^\dagger = 1, \det U = 1\}$$

$$\dim_{\mathbb{R}} \mathfrak{su}(n) = n^2 - 1$$

and so

$$\mathfrak{su}(2) = \{X \in \mathfrak{gl}(2, \mathbb{C}) | X + X^\dagger = 0, \operatorname{tr} X = 0\}$$

$$\exp(tX) \downarrow$$

$$SU(2) = \{U \in GL(n, \mathbb{C}) | UU^\dagger = 1, \det U = 1\}$$

$$\dim_{\mathbb{R}} \mathfrak{su}(2) = 3$$

Also, recall that $\mathfrak{g} \subseteq \mathfrak{gl}(n, \mathbb{C})$ is a vector subspace (13) and that $X \in \mathfrak{g}$ iff $\forall t \in \mathbb{R}, \exp(tX) \in G$.

if $X \in \mathfrak{g}$, if $g \in G$, then $gXg^{-1} \in \mathfrak{g}$

\mathfrak{g} closed under $\mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$

$$(X, Y) \mapsto [X, Y]$$

and so with \mathfrak{g} as a vector space, we can have a choice of bases.

$$\begin{aligned} \xi_1 &= \frac{i}{2} \begin{pmatrix} & 1 \\ 1 & \end{pmatrix} \\ \text{(a) } \xi_2 &= \frac{1}{2} \begin{pmatrix} & -1 \\ 1 & \end{pmatrix} \\ \xi_3 &= \frac{i}{2} \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} \\ &\text{satisfying} \end{aligned}$$

$$[\xi_k, \xi_l] = \epsilon_{klm} \xi_m$$

(b) *Physics*

$$\sigma_1 = -2i\xi_1 = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}$$
$$\sigma_2 = 2i\xi_2 = \begin{pmatrix} & -i \\ i & \end{pmatrix}$$
$$\sigma_3 = -2i\xi_3 = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$$

satisfying

$$[\sigma_k,\sigma_l] = 2i\epsilon_{klm}\sigma_m$$

EY : 20151001 Sage Math 6.8 doesn’t run on Mac OSX El Capitan: I suspect that it’s because in Mac OSX El Capitan, /usr cannot be modified anymore, even in an Administrator account. The TUG group for MacTeX had a clear, thorough, and useful (i.e. copy UNIX commands, paste, and run examples) explanation of what was going on:

<http://tug.org/mactex/elcapitan.html>

So keep in mind that my code for Sage Math is for Sage Math 6.8 that doesn’t run on Mac OSX El Capitan. I’ll also use sympy in Python as an alternative and in parallel.

One can check in sympy the traceless anti-Hermitian (or Hermitian) property of the bases and Pauli matrices, and the commutation relations (see groups.py):

```
import itertools
from itertools import product, permutations

import sympy
from sympy import I, LeviCivita
from sympy import Rational as Rat

from sympy.physics.matrices import msigma # <class 'sympy.matrices.dense.MutableDenseMatrix'>

def commute(A,B):
    """
    commute = commute(A,B)
    commute takes the commutator of A and B
    """
    return (A*B - B*A)

def xi(i):
    """
    xi = xi(i)
    xi is a function that returns the independent basis for
    Lie algebra su(2)\equiv su(2,\mathbb{C}) of Lie group SU(2) of
    traceless anti-Hermitian matrices, based on msigma of sympy
    cf. http://docs.sympy.org/dev/_modules/sympy/physics/matrices.html#msigma
    """
    if i not in [1,2,3]:
        raise IndexError("Invalid Pauli index")
    elif i==1:
        return I/Rat(2)*msigma(1)
    elif i==2:
        return -I/Rat(2)*msigma(2)
    elif i==3:
        return I/Rat(2)*msigma(3)

## check anti-Hermitian property and commutation relations with xi
# xi is indeed anti-Hermitian
xi(1) == -xi(1).adjoint() # True
xi(2) == -xi(2).adjoint() # True
xi(3) == -xi(3).adjoint() # True

# xi obeys the commutation relations
```

```
for i,j in product([1,2,3],repeat=2): print i,j

for i,j in product([1,2,3],repeat=2): print i,j, "\tCommutator:", commute(xi(i),xi(j))

## check traceless Hermitian property and commutation relations with Pauli matrices
# Pauli matrices i.e. msigam is indeed traceless Hermitian

msigma(1) == msigma(1).adjoint() # True
msigma(2) == msigma(2).adjoint() # True
msigma(3) == msigma(3).adjoint() # True

msigma(1).trace() == 0 # True
msigma(2).trace() == 0 # True
msigma(3).trace() == 0 # True

# Pauli matrices obey commutation relation
print "For Pauli matrices, the commutation relations are:\n"
for i,j in product([1,2,3],repeat=2): print i,j, "\tCommutator:", commute(msigma(i),msigma(j))

for i,j,k in permutations([1,2,3],3): print "Commute:", i,j,k, msigma(i), msigma(j), \
":_and_is_2*i_of_", msigma(k), commute(msigma(i),msigma(j)) == 2*I*msigma(k)*LeviCivita(i,j,k)
```

And finally the traceless property of the Pauli matrices:

```
>>> msigma(1).trace()
0
>>> msigma(2).trace()
0
>>> msigma(3).trace()
0
```

7.1. **Spin.** Let’s follow the development by Baez and Muniain (1994) on pp. 175 of the Section II.1 “Lie Groups”, the second (II) chapter on “Symmetry” [8].

Let $V = \mathbb{C}^2$, $G = SU(2)$. Then consider the graded algebra of polynomials on $V = \mathbb{C}^2 \ni (x,y)$

$$P(V) = \bigoplus_{k=0}^{\infty} P^{(k)}(V) = \bigoplus_{\substack{j=0 \\ 2j \in \mathbb{Z}}}^{\infty} P^{(2j)}(V) = \bigoplus_{\substack{j=0 \\ j \in \mathbb{Z}}}^{\infty} P^{(2j)}(V) \oplus \bigoplus_{\substack{j=1/2 \\ 2j \text{ odd}}}^{\infty} P^{(2j)}(V)$$
$$P^{(2j)}(V) \equiv \text{vector space of complex polynomials of degree } 2j$$

and recall this representation on $P^{(2j)}(V)$

$$\rho^{(2j)} : G \rightarrow \text{End}(P^{(2j)}(V))$$
$$\rho^{(2j)} : P^{(2j)}(V) \rightarrow P^{(2j)}(V)$$
$$\rho^{(2j)}(g)(f) = f \circ \rho(g^{-1}) \text{ where } \rho \text{ is the fundamental representation of } G = SU(2)$$
$$\rho^{(2j)}(g)(f)(v) = f \circ \rho(g^{-1})(v) \quad \forall f \in P^{(2j)}(V), \forall v \in V = \mathbb{C}^2$$

Note,
 $\dim P^{(2j)} = \binom{2j+2-1}{2-1} = 2j + 1$

Exercise 21. [8] *spin-0* Consider the trivial representation τ :

$$\begin{array}{ccc} \mathbb{C} & \xrightarrow{T} & P^{(0)}(V) \\ \tau(g) \downarrow & & \downarrow \rho^{(0)}(g) \\ \mathbb{C} & \xrightarrow{T} & P^{(0)}(V) \end{array}$$

$\tau : G \rightarrow \text{End}(\mathbb{C})$
 $\tau(g) : \mathbb{C} \rightarrow \mathbb{C}$
 $\tau(g) = 1_{\mathbb{C}}$

Clearly, $P^{(0)}(V) = \mathbb{C}$, since $P^{(0)}(V)$ consists of polynomials of constants in \mathbb{C} .
Consider $c_0 \in \mathbb{C}$, $f = k_0 \in P^{(0)}(V)$
 $\rho^{(0)}(g)(f) = f \circ \rho(g^{-1}) = k_0$
 $\implies \rho^0(g)T(c_0) = T \circ \tau(g)c_0 = T(c_0)$. Let $T = 1_{\mathbb{C}} = 1_{P^0(V)}$
So $\rho^{(0)}(g) = \tau(g) = 1$. $T = 1$. So representations $\rho^{(0)}$ and trivial representation τ on G are equivalent.

Exercise 22. [8] *spin- $\frac{1}{2}$* For $\text{spin-}\frac{1}{2}$, $j = \frac{1}{2}$, $2j = 1$.

$\forall f \in P^{(1)}(V)$, $V = \mathbb{C}^2$. So in general form, $f(x, y) = ax + by \in P^{(1)}(V)$, $\begin{pmatrix} x \\ y \end{pmatrix} \in V = \mathbb{C}^2$

Recall the fundamental representation

$$\begin{array}{l} \rho : G \rightarrow GL(2, \mathbb{C}) \equiv GL(\mathbb{C}^2) \\ \rho(g) : \mathbb{C}^2 \rightarrow \mathbb{C}^2 \\ \rho(g) = g \end{array}$$

So consider T such that

$$\begin{array}{ccc} \mathbb{C}^2 & \xrightarrow{T} & P^{(1)}(V) \\ \rho(g) \downarrow & & \downarrow \rho^{(1)}(g) \\ \mathbb{C}^2 & \xrightarrow{T} & P^{(1)}(V) \end{array}$$

Consider $\forall v \in \mathbb{C}^2$, $v = \begin{pmatrix} x \\ y \end{pmatrix}$, then

$$\rho(g)v = gv = \begin{bmatrix} ax + by \\ cx + dy \end{bmatrix}$$

```
sage: g*X
[a*x + b*y]
[c*x + d*y]
```

For notation, let $U \in G = SU(2)$ s.t. $UU^\dagger = 1$.
Consider $(\rho^{(2j)}(U)(f))(x) = f(U^{-1}x)$, $\forall x \in \mathbb{C}^2$.

Choose $f(x, y) = x$. So for $f(x, y) = Ax + By$, $A = 1, B = 0$. Choose $U = \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix}$ so $U^{-1} = \begin{pmatrix} \bar{a} & -b \\ \bar{b} & a \end{pmatrix}$. Then

$$U^{-1}x = \begin{pmatrix} \bar{a}x - by \\ \bar{b}x + ay \end{pmatrix}$$

So

$$\begin{aligned} (\rho^{(1)}(U)(f))(x) &= f(U^{-1}x) = \bar{a}x - by \\ (\rho^{(1)}(U)(f))(x) &= f(U^{-1}x) = \bar{b}x + ay \text{ for } f(x, y) = y \end{aligned}$$

Let $f(x, y) = Ax + By$

$$(\rho^{(1)}(U)(f))(x) = f(U^{-1}x) = (A\bar{a} + B\bar{b})x + (Ba - Ab)y = (\bar{a}x - by)A + (\bar{b}x + ay)B = (A\bar{a} + B\bar{b})x + (Ba - Ab)y$$

which was calculated with the assistance of Sage Math:

```
sage: U_try1 = Matrix( [[a.conjugate(), -b],[b.conjugate(), a] ] )
sage: f1( U_try1*X).coefficient(x)
A*conjugate(a) + B*conjugate(b)
sage: f1( U_try1*X).coefficient(y)
B*a - A*b
```

Treating $P^{(1)}(\mathbb{C}^2)$ as a vector space, in its matrix formulation, then $f(x, y) = Ax + By \in P^{(1)}(\mathbb{C}^2)$ is treated as $\begin{bmatrix} A \\ B \end{bmatrix}$, then $(\rho^{(1)}(U)f)$ is

$$\implies \begin{bmatrix} \bar{a} & \bar{b} \\ -b & a \end{bmatrix} \begin{bmatrix} A \\ B \end{bmatrix} = \begin{bmatrix} A\bar{a} + B\bar{b} \\ -Ab + Ba \end{bmatrix}$$

so conclude in general that $\rho^{(1)}(U) = (U^\dagger)^T$.
Now, as Kosmann-Schwarzbach (2010) [11] says, on pp. 13, Chapter 2 Representations of Finite Groups, “Two representations (E_1, ρ_1) and (E_2, ρ_2) are equivalent if and only if there is a basis B_1 of E_1 and a basis B_2 of E_2 such that for every $g \in G$, the matrix of $\rho_1(g)$ in the basis B_1 is equal to the matrix of $\rho_2(g)$ in the basis B_2 . In particular, if the representations (E_1, ρ_1) and (E_2, ρ_2) are equivalent, then E_1 is isomorphic to E_2 .” So we need a change of basis between $\rho(U) = U$ and $\rho^{(1)}(U)$. What’s the linear transformation T s.t.

$$T^{-1}\rho^{(1)}(U)T = U?$$

By intuition,

$$T = \sigma_x \sigma_z \equiv \sigma_1 \sigma_3$$

where σ_i ’s are Pauli matrices.
Indeed,

```
sage: Paulimat[3] * Paulimat[1]*U_try*Paulimat[1] * Paulimat[3]
[conjugate(a) conjugate(b)]
[      -b      a]
```

Then $\rho^{(1)}(U) \circ T = TU$, so this $T = \sigma_1 \sigma_3$ is an “intertwining operator” between $\rho^{(1)}(U)$ and fundamental representation $\rho(U) = U$, with $T = \begin{bmatrix} & -1 \\ 1 & \end{bmatrix}$, and $T^{-1} = \begin{bmatrix} & 1 \\ -1 & \end{bmatrix}$.
 T is an isomorphism between \mathbb{C}^2 and $P^{(1)}(\mathbb{C}^2)$. So fundamental representation ρ of $G = SU(2)$ is equivalent to $\rho^{(1)}(U)$ on $P^{(1)}(\mathbb{C}^2)$.

Exercise 23. [8] (Also from Exercise 2.6 of Kosmann-Schwarzbach (201) [11])

Let (E, π) representation of group G .
 $\forall g \in G$, $\xi \in E^*$, $x \in E$, set $\langle \pi^*(g)(\xi), x \rangle = \langle \xi, \pi(g^{-1})(x) \rangle$
dual (or *contragredient*) of π , $\pi^* : G \rightarrow \text{End}(E^*)$, π^* is a representation, since

$$\begin{aligned} \langle \pi^*(gh)(\xi), x \rangle &= \langle \xi, \pi((gh)^{-1})(x) \rangle = \langle \xi, \pi(h^{-1}g^{-1})(x) \rangle = \langle \xi, \pi(h^{-1})\pi(g^{-1})(x) \rangle = \langle \xi, \pi(h^{-1})(\pi(g^{-1})(x)) \rangle = \\ &= \langle \pi^*(h)(\xi), \pi(g^{-1})(x) \rangle = \langle \pi^*(g)\pi^*(h)(\xi), x \rangle \end{aligned}$$

since this is true, $\forall x \in E$, $\forall \xi \in E^*$, $\pi^*(gh) = \pi^*(g)\pi^*(h)$.
dual π^* of π is a representation.

7.2. Adjoint Representation. I will first follow Sec. 7.3 The Adjoint Representation of Ch. 4 Lie Groups and Lie Algebras of Kosmann-Schwarzbach (201) [11]).

The *conjugation action* $\mathcal{C}_g : G \rightarrow G$ is defined as

$$\begin{aligned}\mathcal{C}_g &: G \rightarrow G \\ \mathcal{C}_g &: h \mapsto ghg^{-1}\end{aligned}$$

So

$$\begin{aligned}\mathcal{C} &: G \rightarrow \text{Aut}(G) \\ \mathcal{C}g &= \mathcal{C}_g\end{aligned}$$

Now define the *adjoint action* of g as the differential or push forward of \mathcal{C}_g :

$$\text{Ad}_g := D_1\mathcal{C}_g \equiv (\mathcal{C}_g)_*|_1 \equiv (\mathcal{C}_g)_*|_{g=1} \quad (\text{adjoint action of } g)$$

Now $\text{Ad}_g : \mathfrak{g} \rightarrow \mathfrak{g}$, so $\text{Ad} : G \rightarrow \text{End}(\mathfrak{g})$

$\text{Ad}(g) \equiv \text{Ad}_g$
Note $\mathcal{C}_{gg'} = \mathcal{C}_g\mathcal{C}_{g'} \equiv \mathcal{C}(gg') = \mathcal{C}(g) \circ \mathcal{C}(g')$ and so

$$\xrightarrow{D_1} \text{Ad}_{gg'} = \text{Ad}_g \circ \text{Ad}_{g'}$$

Kosmann-Schwarzbach (201) [11]) claims, because $\text{Ad}_g = 1_{\mathfrak{g}}$ when $g = 1$,

$\text{Ad} : G \rightarrow GL(\mathfrak{g})$ is a representation of G on \mathfrak{g} . (EY : 20160505 ???)

$\text{Ad} : g \mapsto \text{Ad}_g$

Definition 27. *representation* Ad of G on $V = \mathfrak{g}$ is called *adjoint representation* of Lie group G .

Denote adjoint representation of Lie algebra \mathfrak{g} , ad .

By definition, $\text{Ad}_{\exp(tX)} = \exp(t\text{ad}_X)$

cf. Prop. 7.8 of Kosmann-Schwarzbach (201) [11])

Proposition 12. (1) *Let A invertible matrix, $A \in \text{Lie group } G$.*

Let X matrix s.t. $X \in \mathfrak{g}$. Then

$$\text{Ad}_A(X) = AXA^{-1}$$

(2) *Let $X, Y \in \mathfrak{g}$. Then*

$$\text{ad}_X(Y) = [X, Y]$$

(3) *Let $X, Y \in \mathfrak{g}$. Then*

$$\text{ad}_{[X, Y]} = [\text{ad}_X, \text{ad}_Y]$$

Proof. (1) By def., $\forall B \in G$, $\mathcal{C}_A(B) = ABA^{-1}$, and thus

$$\text{Ad}_A(X) = \left. \frac{d}{dt} A \exp(tX) A^{-1} \right|_{t=0} = AXA^{-1}$$

(2)

$$\begin{aligned}\text{ad}_X(Y) &= \left. \frac{d}{dt} \text{Ad}_{\exp(tX)}(Y) \right|_{t=0} = \left. \frac{d}{dt} \exp(tX) Y \exp(tX) \right|_{t=0} = \\ &= XY - YX = [X, Y]\end{aligned}$$

(3) Use Jacobi identity:

$$[A, [B, C]] + [B, [C, A]] + [C, [A, B]] = 0 \text{ or}$$

$$[[A, B], C] = [A, [B, C]] - [B, [A, C]]$$

$$\text{ad}_{[X, Y]}C = [[X, Y], C] = [X, [Y, C]] - [Y, [X, C]] = [X, \text{ad}_Y C] - [Y, \text{ad}_X C] \text{ and that}$$

$$\text{ad}_X \text{ad}_Y C = [X, [Y, C]] \implies \text{ad}_{[X, Y]} C = [\text{ad}_X, \text{ad}_Y] C$$

Part 5. Cohomology; Stoke's Theorem

8. STOKE'S THEOREM

Theorem 14 (Stoke's Theorem). *Let M be oriented, smooth n -manifold with boundary, let ω be a compactly supported smooth $(n-1)$ -form on M , or if $\omega \in A_c^{n-1}(M)$, Then*

$$(20) \quad \int_M d\omega = \int_{\partial M} \omega$$

*If $\partial M = \emptyset$, then $\int_{\partial M} \omega = 0$
 $\int_{\partial M} \omega$ interpreted as $\int_{\partial M} i_{\partial M}^* \omega = \int_{\partial M} i^* \omega$ so*

$$(21) \quad \int_M d\omega = \int_{\partial M} i^*(\omega)$$

where inclusion $i : \partial M \hookrightarrow M$

Proof. Begin with very special case:

Suppose $M = \mathbb{H}^n$ (upper half space), $\partial M = \mathbb{R}^{n-1}$

ω has compact support, so $\exists R > 0$ s.t. $\text{supp } \omega \subseteq \text{rectangle } A = [-R, R] \times \cdots \times [-R, R] \times [0, R]$.

$$\forall \omega \in A_c^{n-1}(\mathbb{H}^n)$$

$$(22) \quad \omega = \sum_{j=1}^n (-1)^{j-1} f_j dx^1 \wedge \cdots \wedge \widehat{dx}^j \wedge \cdots \wedge dx^n \equiv \sum_{i=1}^n \omega_i dx^1 \wedge \cdots \wedge \widehat{dx}^i \wedge \cdots \wedge dx^n$$

with Conlon (2008) [16] and John Lee (2012) [3]'s notation, respectively, and where f_j has compact support.

$$i^* \omega = (f_1 \circ i) dx^2 \wedge \cdots \wedge dx^n \in A_c^{n-1}(\partial \mathbb{H}^n)$$

$$\begin{aligned}d\omega &= \sum_{i=1}^n d\omega_i \wedge dx^1 \wedge \cdots \wedge \widehat{dx}^i \wedge \cdots \wedge dx^n = \sum_{i,j=1}^n \frac{\partial \omega_i}{\partial x^j} dx^j \wedge dx^1 \wedge \cdots \wedge \widehat{dx}^i \wedge \cdots \wedge dx^n = \\ &= \sum_{i=1}^n (-1)^{i-1} \frac{\partial \omega_i}{\partial x^i} dx^1 \wedge \cdots \wedge dx^n\end{aligned}$$

i.e. (for another notation)

$$d\omega = \left(\sum_{j=1}^n \frac{\partial f_j}{\partial x^j} \right) dx^1 \wedge \cdots \wedge dx^n \in A_c^n(\mathbb{H}^n)$$

$$d\omega = \left(\sum_{j=1}^n \frac{\partial f_j}{\partial x^j} \right) dx^1 \wedge \cdots \wedge dx^n \in A_c^n(\mathbb{H}^n)$$

$$\int_{\mathbb{H}^n} d\omega = \sum_{i=1}^n (-1)^{i-1} \int_A \frac{\partial \omega_i}{\partial x^i} dx^1 \wedge \cdots \wedge dx^n = \sum_{i=1}^n (-1)^{i-1} \int_0^R \int_{-R}^R \cdots \int_{-R}^R dx^1 \cdots dx^n \frac{\partial \omega_i}{\partial x^i}(x)$$

We can change order of integration in each term so to do x^i integration first.

By fundamental thm. of calculus, terms for which $i \neq n$ reduce to

$$\begin{aligned} \sum_{i=1}^{n-1} (-1)^{i-1} \int_0^R \int_{-R}^R \cdots \int_{-R}^R \frac{\partial \omega_i}{\partial x^i}(x) dx^1 \dots dx^n &= \sum_{i=1}^{n-1} (-1)^{i-1} \int_0^R \int_{-R}^R \cdots \int_{-R}^R \frac{\partial \omega_i}{\partial x^i}(x) dx^i dx^1 \dots \widehat{dx}^i \dots dx^n = \\ &= \sum_{i=1}^{n-1} (-1)^{i-1} \int_0^R \int_{-R}^R \cdots \int_{-R}^R [\omega_i(x)]_{x^i=-R}^{x^i=R} dx^1 \dots \widehat{dx}^i \dots dx^n = 0 \end{aligned}$$

because we've chosen R large enough that $\omega = 0$ when $x^i = \pm R$.

Part 6. Prástaro

Prástaro (1996) [12]

8.0.1. *Affine Spaces.* cf. Sec. 1.2 - *Affine Spaces* of Prástaro (1996) [12]

Definition 28 (affine space).

$$(23) \quad \begin{aligned} &\text{affine space} \quad (M, \mathbf{M}, \alpha) \\ &\quad \text{with} \\ &M \equiv \text{set (set of pts.)} \\ &\mathbf{M} \equiv \text{vector space (space of free vectors)} \\ &\alpha \equiv \mathbf{M} \times M \rightarrow M \equiv \text{translation operator} \\ &\alpha : (v, p) \mapsto p' \equiv p + v \end{aligned}$$

Note: α is a **transitive** action and without fixed pts. (free).

i.e. $\forall p \in M$,

$$\begin{aligned} &\forall \text{ pt. } O \in M, \alpha : (v, O) \mapsto O' \equiv O + v, \alpha(\cdot, O) \equiv \alpha_O \equiv \alpha(O). \quad \alpha_O(v) = O' = O + \mathbf{v} \quad \forall O' \in M, \exists \mathbf{v} \in \mathbf{M} \text{ s.t. } O' = O + \mathbf{v} \\ &\implies M \equiv \mathbf{M}. \\ &\forall (O, \{e_i\})_{1 \leq i \leq n}, \text{ where } \{e_i\} \text{ basis of } \mathbf{M}, M \equiv \mathbf{M} = \mathbb{R}^n \text{ so isomorphism } M \simeq \mathbb{R}^n \end{aligned}$$

i.e. α is **without fixed pts.**, meaning,

Given pointed space (M, O) , where base pt. $O \in M$, we can associate $\forall p \in M$, vector $\mathbf{x} \in \mathbf{M}$, by 1-to-1 mapping $M \rightarrow \mathbf{M}$.

So for

$$\alpha : \mathbf{M} \times M \rightarrow M$$

$$\alpha(\mathbf{x}, p) = p' = p + \mathbf{x}$$

Consider

$$\alpha(\mathbf{x}, O) = p = \alpha_O(\mathbf{x}) = p \implies \exists \alpha_O^{-1}(p) = \mathbf{x} \in \mathbf{M}$$

(1) tangent space of M in $p \in M$ is vector space $T_p M \equiv (\mathbf{M}, p) \cong M$

(2) If \mathbf{M} Euclidean space, affine space (M, \mathbf{M}, α) is Euclidean

(3) Call dim. of affine space (M, \mathbf{M}, α) , dim. of $\mathbf{M} \equiv \dim \mathbf{M}$

$\{\mathbf{e}_i\}$ basis of \mathbf{M}

Definition 29. $(O, \{e_i\}) \equiv \text{affine frame}$.

$\forall \text{ affine frame } (O, \{e_i\}), \exists \text{ coordinate system } x^\alpha : M \rightarrow \mathbb{R}$,

where $x^\alpha(p)$ is α th component, in basis $\{e_i\}$, of vector $p - O$

Proposition 13 (1.6, Prástaro (1996) [12]). $\forall O \in M$, we have canonical identification $M \equiv \mathbf{M}$, since

$$\alpha_O^{-1} : M \rightarrow \mathbf{M}$$

$$\alpha_O : \mathbf{M} \rightarrow M$$

$$\alpha_O^{-1}(p) = \mathbf{x}$$

$$\alpha_O : \mathbf{x} = \alpha(\mathbf{x}, O) = p$$

Furthermore,

\forall **affine frame** $(O, \{\mathbf{e}_i\})_{1 \leq i \leq d}$, where $\{\mathbf{e}_i\}$ basis of \mathbf{M} ,

\exists isomorphism $M \cong \mathbb{R}^d$,

Then, $\forall (O, \{\mathbf{e}_i\})_{1 \leq i \leq d}$,

\exists coordinate system $x^\alpha : M \rightarrow \mathbb{R}$,

where $x^\alpha(p) = \alpha$ th component, in basis $\{\mathbf{e}_i\}$, of vector $p - O$.

Theorem 15 (1.4 Prástaro (1996) [12]). Let $(x^\alpha), (\bar{a}^\alpha)$ 2 coordinate systems correspond to affine frames $(O, \{e_i\}), (\bar{O}, \{\bar{e}_i\})$, respectively.

□

$$(24) \quad \bar{x}^\alpha = A^\alpha_\beta x^\beta + y^\alpha$$

where

$$y^\alpha \in \mathbb{R}^n, \quad A^\alpha_\beta \in GL(n; \mathbb{R})$$

Definition 30 (1.10 Prástaro (1996) [12]).

$$(25) \quad A(n) \equiv GL(n, \mathbb{R}) \times \mathbb{R}^n$$

affine group of dim. n

Theorem 16 (1.5). symmetry group of n -dim. affine space, called affine group $A(M)$ of M . \exists isomorphism,

$$(26) \quad A(M) \simeq A(n), \quad f \mapsto (f^\alpha_\beta, y^\alpha); \quad f^\alpha \equiv x^\alpha \circ f = f^\alpha_\beta x^\beta + y^\alpha$$

cf. Eq. 1.4 Prástaro (1996) [12]

Definition 31 (metric). Let smooth manifold M , $\dim M = n$, $\forall p \in M$, \exists vector space $T_p M$, and so for

$$(27) \quad \begin{aligned} &g_p(T_p M)^2 \rightarrow \mathbb{R} \\ &g_p : (X_p, Y_p) \mapsto g_p(X_p, Y_p) \in \mathbb{R} \end{aligned}$$

with g_p being bilinear, symmetric (in X_p, Y_p), nondegenerate (i.e. if $g_p(X_p, Y_p) = 0$, then X_p or $Y_p = 0$)

Note that

$$g \in \Gamma((TM \otimes TM)^*)$$

and that for $X = X^i \frac{\partial}{\partial x^i}$ so

$$Y = Y^i \frac{\partial}{\partial x^i}$$

$$g(X, Y) = g_{ij} X^i Y^j$$

Now for

$$F : M \rightarrow N \quad DF \equiv F_* : T_p M \rightarrow T_{F(p)} N$$

$$F : x \mapsto y = y(x) \quad DF : X_p \mapsto (DF)(X^j \frac{\partial}{\partial x^j}) = X^j \frac{\partial y^i}{\partial x^j} \frac{\partial}{\partial y^i}$$

$$\begin{aligned} (F^* g')(X, Y) &= (F^* g')(X^i \frac{\partial}{\partial x^i}, Y^j \frac{\partial}{\partial x^j}) = (F^* g')_{ij} X^i Y^j = g'(F_* X, F_* Y) = \\ &= \end{aligned}$$

Part 7. Connections

9. CONNECTIONS OF VECTOR BUNDLES

[24]

Definition 32 (Connection in a vector bundle). ***connection in a vector bundle** $\pi : E \rightarrow M$ over C^∞ manifold M , is a bilinear map*

$$\nabla : \mathfrak{X}(M) \times \Gamma(E) \rightarrow \Gamma(E)$$

satisfying

- (i) $\nabla_{fX}s = f\nabla_X s$
- (ii) $\nabla_X(fs) = f\nabla_X s + (Xf)s$ where $f \in C^\infty(M)$, $X \in \mathfrak{X}(M)$, $s \in \Gamma(E)$

$\nabla_X s$ is covariant derivative of s relative to X (for Morita (2001)[24])

Claim: Any vector bundle admits a connection.

e.g. product bundle $M \times \mathbb{R}^n$. Let x_1, \dots, x_n be canonical coordinates in \mathbb{R}^n . Take frame field (s_1, \dots, s_n) , where $s_i(p) = \frac{\partial}{\partial x^i}$. Set $\nabla_X s_i = 0$ ($i = 1, \dots, m$) \forall vector space X , $\forall s = \sum_i a_i s_i$, $\forall X \in \mathfrak{X}(M)$, set

$$\nabla_X s = \sum_{i=1}^n (Xa_i) s_i$$

For this connection $\nabla_X s$ is the partial derivative in direction of X if s is considered \mathbb{R}^n -valued function on M . Call it **trivial connection** in product bundle.

Indeed,

$$\begin{aligned} \nabla_X s &= \nabla_{X^i \frac{\partial}{\partial x^i}} \left(s^m \frac{\partial}{\partial x^m} \right) = X^i \nabla_{\frac{\partial}{\partial x^i}} \left(s^m \frac{\partial}{\partial x^m} \right) = X^i \left(\nabla_{\frac{\partial}{\partial x^i}} s^m \right) \frac{\partial}{\partial x^m} + X^i s^m \nabla_{\frac{\partial}{\partial x^i}} \frac{\partial}{\partial x^m} = \\ &= X^i \left(\frac{\partial}{\partial x^i} s^m \right) \frac{\partial}{\partial x^m} + X^i s^m \Gamma_{mi}^q \frac{\partial}{\partial x^q} = X^i \frac{\partial s^m}{\partial x^i} \frac{\partial}{\partial x^m} + 0 \end{aligned}$$

if $\Gamma_{mi}^q = 0$ at a chosen point p .

For arbitrary vector bundle $\pi : E \rightarrow M$, take locally finite open covering $\{U_\alpha\}_{\alpha \in A}$ s.t. $\pi^{-1}(U_\alpha)$ trivial. Denote ∇^α trivial connection $\forall \pi^{-1}(U_\alpha)$. Let $\{f_\alpha\}$ be a partition of unity for covering U_α , define

$$\nabla_X s := \sum_\alpha f_\alpha \nabla_X^\alpha s$$

Verify this defines connection in E :

$$\begin{aligned} \nabla_X(gs) &= \sum_\alpha f_\alpha \nabla_X^\alpha(gs) = \sum_\alpha f_\alpha \left[X^i \left(\frac{\partial g}{\partial x^i} \right) s + X^i g \frac{\partial s^m}{\partial x^i} \frac{\partial}{\partial x^m} \right] = \\ &= g \sum_\alpha f_\alpha \nabla_X^\alpha s + \sum_\alpha f_\alpha (Xg)s \end{aligned}$$

Proposition 14 (5.18 Morita (2001)[24]). *Let ∇_i ($1 \leq i \leq k$) be k connections in a given vector bundle. Then \forall linear combination $\sum_{i=1}^k t_i \nabla_i$, where $t_1 + \dots + t_k = 1$ is a connection.*

Proof. TODO: Ex. 5.5

Part 8. Holonomy

Definition 33 (Conlon, 10.1.2). *If $X, Y \in \mathfrak{X}(M)$, $M \subset \mathbb{R}^m$, **Levi-Civita connection** on $M \subset \mathbb{R}^m$*

$$(28) \quad \begin{aligned} \nabla : \mathfrak{X}(M) : \mathfrak{X}(M) &\rightarrow \mathfrak{X}(M) \\ \nabla_X Y &:= p(D_X Y) \end{aligned}$$

with

$$\begin{aligned} D_X Y &:= \sum_{j=1}^m X(Y^j) \frac{\partial}{\partial x^j} = \sum_{i,j=1}^m X^i \frac{\partial Y^j}{\partial x^i} \frac{\partial}{\partial x^j} & \forall X &= \sum_{i=1}^m X^i \frac{\partial}{\partial x^i}, \\ \forall Y &= \sum_{i=1}^m Y^i \frac{\partial}{\partial x^i} \end{aligned}$$

$$\nabla_{fX} Y = f(D_{fX} Y) = p(f D_X Y) = fp D_X Y = f \nabla_X Y$$

$$\nabla_X fY = p(D_X fY) = p \left(\sum_{i,j=1}^m \left(X^i f \frac{\partial Y^j}{\partial x^i} + X^i Y^j \frac{\partial f}{\partial x^i} \right) \frac{\partial}{\partial x^j} \right) = f \nabla_X Y + p \sum_{j=1}^m X(f) Y^j \frac{\partial}{\partial x^j} = f \nabla_X Y + X(f) p(Y)$$

Definition 34 (Conlon, 10.1.4; Christoffel symbols).

$$(29) \quad \begin{aligned} \nabla_{\frac{\partial}{\partial x^i}} \frac{\partial}{\partial x^j} &= \Gamma_{ij}^k \frac{\partial}{\partial x^k} & (\text{Conlon's notation}) \\ \nabla_{\frac{\partial}{\partial x^j}} \frac{\partial}{\partial x^i} &= \Gamma_{ij}^k \frac{\partial}{\partial x^k} & (F. Schuller's notation) \end{aligned}$$

Definition 35 (torsion).

$$(30) \quad \begin{aligned} T : \mathfrak{X}(M) \times \mathfrak{X}(M) &\rightarrow \mathfrak{X}(M) \\ T(X, Y) &= \nabla_X Y - \nabla_Y X - [X, Y] \end{aligned}$$

If $T = 0$, ∇ torsion-free or symmetric.

$$T(fX, Y) = f \nabla_X Y - (f \nabla_Y X + Y(f)X) - \{(fXY - (Y(f)X + fYX))\} = fT(X, Y)$$

$$T(X, fY) = f \nabla_X Y + X(f)Y - f \nabla_Y X - \{((X(f)Y + fXY) - fYX)\} = fT(X, Y)$$

Thus, $T(X, Y)$ $C^\infty(M)$ -bilinear.

$T \in \tau_1^2(M)$.

$T(v, w) \in T_x M$ defined, $\forall v, w \in T_x M$, $\forall x \in M$.

Thus, torsion is a **tensor**.

Exercise 10.1.7 Conlon (2008)[16] . .

If $T(X, Y) = 0$,

$$T(e_i, e_j) = \Gamma_{ji}^k e_k - \Gamma_{ij}^k e_k - 0 = 0 \implies \Gamma_{ji}^k = \Gamma_{ij}^k$$

If $\Gamma_{ij}^k = \Gamma_{ji}^k$, $T(e_i, e_j) = 0$.

Exercise 10.1.8, Conlon (2008)[16].

If $M \subset \mathbb{R}^m$ smoothly embedded submanifold, $\forall \frac{\partial}{\partial x^j}, \frac{\partial}{\partial x^i} \in T_x M$, spanning $T_x M$, consider $\frac{\partial}{\partial x^j} = X_j^k \frac{\partial}{\partial \tilde{x}^k}$, $\frac{\partial}{\partial x^i} = X_i^k(\tilde{x}) \frac{\partial}{\partial \tilde{x}^k}$

$$\nabla_{\frac{\partial}{\partial x^j}} \frac{\partial}{\partial x^i} = p D_{X_j^k \frac{\partial}{\partial \tilde{x}^k}} X_i^l \frac{\partial}{\partial \tilde{x}^l} = p \left(X_j^k \frac{\partial X_i^l}{\partial \tilde{x}^k} \frac{\partial}{\partial \tilde{x}^l} \right) = X_j^k p \left(\frac{\partial X_i^l}{\partial \tilde{x}^k} \frac{\partial}{\partial \tilde{x}^l} \right)$$

$$\nabla_{\frac{\partial}{\partial x^i}} \frac{\partial}{\partial x^j} = X_i^k p \left(\frac{\partial X_j^l}{\partial \tilde{x}^k} \frac{\partial}{\partial \tilde{x}^l} \right)$$

□

If $X \in \mathfrak{X}(M)$, smooth $s : [a, b] \rightarrow M$, then $\forall s(t)$,

$$X'_{s(t)} = \nabla_{\dot{s}(t)} X \in T_{s(t)} M$$


In fact, it's often natural to consider fields $X_{s(t)}$ along s , parametrized by parameter t , allowing

$$X_{s(t_1)} \neq X_{s(t_2)}$$

each of $s(t_1) = s(t_2)$.

Definition 36 (10.1.9). *Let smooth $s : [a, b] \rightarrow M$.*

Vector field along s is smooth $v : [a, b] \rightarrow TM$ s.t.



commutes.

Note that $v \in \mathfrak{X}(s) \subset \mathfrak{X}(M)$

e.g. $(Y|s)(t) = Y_{s(t)}$, restriction of $Y \in \mathfrak{X}(M)$ to s .

e.g. $\dot{s}(t) \in \mathfrak{X}(M)$.

$\forall v, w \in \mathfrak{X}(s)$, $v + w \in \mathfrak{X}(s)$,

$$(fv + gv)(t) := (f(s(t)) + g(s(t)))v(t) = f(s(t))v(t) + g(s(t))v(t) = (f + g)v(t)$$

Likewise,

$$f(v + w) = fv + fw$$

$\mathfrak{X}(s)$ is a real vector space and $C^\infty[a, b]$ -module.

Definition 37 (10.1.10). *Let conection ∇ on M .*

Associated covariant derivative is operator

$$\frac{\nabla}{dt} \mathfrak{X}(s) \rightarrow \mathfrak{X}(s)$$

\forall smooth s on M , s.t.

- (1) $\frac{\nabla}{dt}$ \mathbb{R} -linear
- (2) $\left(\frac{\nabla}{dt}\right)(fv) = \frac{df}{dt}v + f\frac{\nabla}{dt}v$, $\forall f \in C^\infty[a, b]$, $\forall v \in \mathfrak{X}(s)$
- (3) If $Y \in \mathfrak{X}(M)$, then

$$\frac{\nabla}{dt}(Y|s)(t) = \nabla_{\dot{s}(t)}Y \in T_{s(t)}M, \quad a \leq t \leq b$$

Theorem 17 (Conlon Thm. 10.1.11[16]). \forall connection ∇ on M , $\exists!$ associated covariant derivative $\frac{\nabla}{dt}$

Proof. Consider arbitrary coordinate chart $(U, x^1 \dots x^n)$.

Consider smooth curve $s : [a, b] \rightarrow U$.

Let $v \in \mathfrak{X}(s)$, $v(t) = v^i(t) \frac{\partial}{\partial x^i}$; $\dot{s}(t) = s^j \frac{\partial}{\partial x^j}$.

$$\frac{\nabla v}{dt} = \frac{dv^i(t)}{dt} \frac{\partial}{\partial x^i} + v^i(t) \frac{\nabla}{dt} \frac{\partial}{\partial x^i} = \frac{dv^i}{dt} \frac{\partial}{\partial x^i} + v^i \nabla_{\dot{s}(t)} \frac{\partial}{\partial x^i} = \dot{v}^i \frac{\partial}{\partial x^i} + v^i \dot{s}^j \Gamma_{ij}^k \frac{\partial}{\partial x^k} = (\dot{v}^k + v^i \dot{s}^j \Gamma_{ij}^k) \frac{\partial}{\partial x^k}$$

This is an explicit, local formula in terms of connection, proving uniqueness.

Existence: \forall coordinate chart $(U, x^1 \dots x^n)$, $(\dot{v}^k + v^i \dot{s}^j \Gamma_{ij}^k) \frac{\partial}{\partial x^k} =: \frac{\nabla v}{dt}$.

$$\frac{\nabla}{dt}(fv) = \dot{f}v^k + f\dot{v}^k + fv^i \dot{s}^j = \dot{f}v + f \frac{\nabla v}{dt}$$

If f constant, then $\frac{\nabla}{dt}$ is \mathbb{R} -linear.

Definition 38 (10.1.12 Conlon (2008)[16]). *Let (M, ∇) . Let $v \in \mathfrak{X}(s)$ for smooth $s : [a, b] \rightarrow M$.*

*If $\frac{\nabla v}{dt} \equiv 0$ on s , then v is **parallel** along s .*

Theorem 18 (10.1.13). *Let (M, ∇) , smooth $s : [a, b] \rightarrow M$, $c \in [a, b]$, $v_0 \in T_{s(c)}M$.*

Then $\exists!$ parallel field $v \in \mathfrak{X}(s)$ s.t. $v(c) = v_0$.

v parallel transport along s .

Proof.

$$\begin{aligned} \dot{s}(t) &= \dot{s}^j(t) e_j \\ v(t) &= v^i(t) e_i \\ v_0 &= a^i e_i \\ 0 &= \left(\frac{dv^k}{dt}(t) + v^i(t) \dot{s}^j(t) \Gamma_{ij}^k(s(t)) \right) e_k \end{aligned}$$

or equivalently

$$(31) \quad \frac{dv^k}{dt} = -v^i \dot{s}^j \Gamma_{ij}^k, \quad 1 \leq k \leq n \quad (10.1)$$

with initial conditions $v^k(c) = a^k$, $1 \leq k \leq n$.

By existence and uniqueness of solutions of O.D.E.

$\exists \epsilon > 0$ s.t. $\exists!$ solutions $v^k(t)$. For $c - \epsilon < t < c + \epsilon$.

In fact, these ODEs being linear in v^k , by ODE theory (Appendix C, Thm. C.4.1).

\nexists restriction on ϵ , so $\exists! v^k(t) \forall t \in [a, b]$, $1 \leq k \leq n$

□

9.1. Principal bundle, vector bundle case for parallel transport. Recall the 2 different forms or viewpoints for Lie-algebra valued 1-forms, or vector-valued 1-forms, or sections of 1-form-valued endomorphisms:

$$\omega_{i\mu}^k dx^\mu \equiv \omega_i^k \in \Omega^1(M, \mathfrak{gl}(n, \mathbb{F})) = \Gamma(\mathfrak{gl}(n, \mathbb{R} \otimes T^*M|_U))$$

for $i, k = 1 \dots n = \dim E$.

$\mu = 1 \dots d = \dim E$

Now

$$D_X \mu = X^\mu D_{\frac{\partial}{\partial x^\mu}} \mu = X^\mu \left[\left(\frac{\partial}{\partial x^\mu} \mu^k \right) e_k + \mu^i \omega_{i\mu}^k e_k \right] = (X(\mu^k) + \mu^i \omega_i^k(X)) e_k = (d\mu^k(X) + \mu^i \omega_i^k(X)) e_k$$

So then define

$$(32) \quad \begin{aligned} D : \Gamma(E) &\rightarrow \Gamma(E) \otimes \Gamma(T^*M) \\ D\mu &= D(\mu^i e_i) = e_k (d\mu^k + \mu^i \omega_i^k) \equiv (d + A)\mu \end{aligned}$$

Also, D can be defined for this case:

$$D : \Gamma(\text{End}(E)) \rightarrow \Gamma(\text{End}E) \otimes \Gamma(T^*M)$$

Let $\sigma = \sigma_j^i e_i \otimes e^j \in \Gamma(\text{End}(E))$

$$(33) \quad \begin{aligned} D\sigma &= D(\sigma_j^i e_i) \otimes e^j + \sigma_j^i e_i \otimes D^* e^j = (d\sigma_j^k + \sigma^i A_i^k) e_k \otimes e^j + \sigma_j^i e_i \otimes (A^*)_k^j e^k = \\ &= (d\sigma_j^k + \sigma_j^i A_i^k) e_k \otimes e^j + \sigma_i^k e_j \otimes (-A_j^i) e^j = (d\sigma_j^k + [A, \sigma]_j^k) e_k \otimes e^j \end{aligned}$$

cf. Def. 4.1.4 of Jost (2011), pp. 138.

For $\mu \in \Gamma(E)$, smooth $s : [a, b] \rightarrow M$, $X(t) = \dot{s}(t)$,

$$(34) \quad D_{\dot{s}(t)} \mu = \dot{s}^\mu D_{\frac{\partial}{\partial x^\mu}} \mu = \dot{s}^\mu \left[\frac{\partial \mu^k}{\partial x^\mu} e_k + \mu^i \omega_{i\mu}^k e_k \right] = \left[\dot{s}^\mu \frac{\partial \mu^k}{\partial x^\mu} + \dot{s}^\mu \mu^i \omega_{i\mu}^k \right] e_k = \frac{d}{dt} \mu(s(t)) + \mu^i \dot{s}^\mu \omega_{i\mu}^k e_k$$

□

Let $D_{\dot{s}(t)}\mu = 0$. Then,

$$(35) \quad \frac{d}{dt}\mu(s(t)) = -\mu^i \dot{s}^\mu \omega_{i\mu}^k e_k$$

Recall, given vector bundle $E \xrightarrow{\pi} N$, given $\varphi : M \rightarrow N$, then pullback

$$(36) \quad \varphi^* E \rightarrow M$$

i.e.

$$\begin{array}{ccc} \varphi^* E & \xleftarrow{\varphi^*} & E \\ \downarrow \psi & & \downarrow \pi \\ M & \xrightarrow{\varphi} & N \end{array} \quad \begin{array}{c} (\varphi^* E)_x = E_{\varphi(x)} \\ \uparrow \\ x \in M \end{array}$$

i.e. if $s \in \Gamma(E)$,

$$\varphi^* s = s \circ \varphi \in \Gamma(\varphi^* E)$$

Thus,

$$\begin{array}{ccc} \gamma^* E & \xleftarrow{\gamma^*} & E \\ \downarrow & & \downarrow \pi \\ [a, b] & \xrightarrow{\gamma} & M \end{array} \quad \begin{array}{c} (\varphi^* E)_c = E_{\gamma(c)} \\ \uparrow \\ c \in [a, b] \end{array}$$

For

$$\begin{aligned} \dot{v}^k &= -v^i \dot{s}^j \Gamma_{ij}^k \\ v^k(c) &= v_0^k \quad 1 \leq k \leq m \\ \dot{v} &= -v^i \dot{s}^j \Gamma_{ij} \end{aligned}$$

$$(v + w) = -(v^i + w^i) \dot{s}^j \Gamma_{ij}(v + w)(c) = v(c) + w(c) = v_0 + w_0$$

so $v + w \in \mathfrak{X}(s)$ is parallel transport of $v_0 + w_0$.

Likewise, $\forall a \in \mathbb{F}$, $av \in \mathfrak{X}(s)$ is the parallel transport of av_0 .

$$\dot{\mu}^k = -\mu^i \dot{s}^\mu \omega_{i\mu}^k = -\mu^i \omega_i^k(\dot{s}^\mu)$$

Suppose $\gamma^* E$ trivialized over $[a, b]$.

Closed interval is contractible, so this is always possible.

For chart (U, φ) ,

$$\begin{array}{ccc} \gamma^* E & \xleftarrow{\gamma^*} & E \\ \downarrow & & \downarrow \pi \\ [a, b] & \xrightarrow{\gamma} & M \end{array} \quad \begin{array}{ccc} E|_U & \xrightarrow{\psi} & U \times V \\ \uparrow \pi^{-1} & \nearrow & \\ U \subset M & & \end{array}$$

Consider

$$\begin{aligned} \varphi : [a, b] \times V &\rightarrow \gamma^* E \\ \varphi(t, \cdot) &= \gamma^* \circ \psi^{-1}(\gamma(t), \cdot) \end{aligned}$$

$$\forall \mu \in \Gamma(E|_{x \in M}),$$

$$\mu = \mu^i e_i.$$

$$\varphi(t, e_i) = \epsilon_i \text{ is a basis for } \gamma^* E.$$

$$\forall \sigma \in \Gamma(\gamma^* E),$$

$$\begin{aligned} \sigma &= \sigma^i \epsilon_i, \quad \sigma^i : [a, b] \rightarrow \mathbb{F} \\ \nabla_{\frac{\partial}{\partial x^\mu}} \sigma &= \frac{\partial \sigma^k}{\partial x^\mu} \epsilon_k + \omega_{j\mu}^k \sigma^j \epsilon_k = \left(\frac{\partial \sigma^k}{\partial x^\mu} + \omega_{j\mu}^k \sigma^j \right) \epsilon_k \\ \nabla \sigma &= \epsilon_k \otimes (d\sigma^k + \omega_{j\mu}^k dx^\mu \sigma^j) = \epsilon_k \otimes (d\sigma^k + \omega_{j\mu}^k \sigma^j) \\ \nabla_{\frac{d}{dt}} \sigma &= \epsilon_k \otimes \left(\frac{d\sigma^k}{dt} + \omega_{j\mu}^k \dot{x}^\mu \sigma^j \right) \end{aligned}$$

Now

$$\frac{d}{dt} = \dot{x}^\nu \frac{\partial}{\partial x^\nu}$$

Then σ parallel along γ if

$$\frac{d\sigma^k}{dt} + \omega_{j\mu}^k \dot{x}^\mu \sigma^j = 0$$

Definition 39 (3.1.4 [17]). *Parallel transport along γ is*

$$(37) \quad \begin{aligned} P_\gamma : E_{\gamma(a)} &\rightarrow E_{\gamma(b)} \\ P_\gamma(v) &\mapsto \sigma(b) \end{aligned}$$

where $\sigma \in \Gamma(\gamma^* E)$, σ unique and s.t. $\sigma(a) = v$.

Lemma 1 (10.1.16[16]). *holonomy*

$$h_s : T_x M \rightarrow T_{x_0} M$$

if ∇ around piecewise smooth loop s is a linear transformation.

Lemma 2 (10.1.18 Conlon (2008)[16]). *Let piecewise smooth loop $s : [a, b] \rightarrow M$ at x_0 .*

Let weak reparametrization $\tilde{s} = s \circ r : [c, d] \rightarrow M$.

If reparametrization is orientation-preserving, then $h_{\tilde{s}} = h_s$,

If reparametrization is orientation-reversing, then $h_{\tilde{s}} = h_s^{-1}$,

Proof. Without loss of generality, assume smooth s, r

$$\tilde{s}(\tau) = s(r(\tau))$$

$$\tilde{v}(\tau) = v(r(\tau))$$

$$\tilde{u}^j(\tau) = \frac{dt}{d\tau}(\tau) u^j(r(\tau))$$

$$\frac{d\tilde{v}^k}{d\tau}(\tau) = \frac{dr}{d\tau}(\tau) \frac{dv^k}{dt}(r(\tau))$$

$$\frac{d\tilde{v}^k}{d\tau} = -\tilde{v}^i \tilde{u}^j \Gamma_{ij}^k$$

since

$$\frac{dv^k}{dt} = -v^i u^j \Gamma_{ij}^k; \quad 1 \leq k \leq n$$

$$v^k(c) = a^k; \quad 1 \leq k \leq a$$

$$\frac{dr}{d\tau} \frac{dv^k}{dt} = -v^i \frac{dr}{d\tau} u^j \Gamma_{ij}^k = \frac{d\tilde{v}^k}{d\tau} = -\tilde{v}^i \tilde{u}^j \Gamma_{ij}^k$$

Thus, if $r(c) = a$, $r(d) = b$

$$h_{\tilde{s}}(v_0) = \tilde{v}(d) = v(b) = h_s(v_0)$$

If $r(c) = a$, $r(d) = b$, then

$$\tilde{v}(c) = v(b) = h_s(v_0)$$

and

$$h_{\tilde{s}}(h_s(v_0)) = h_{\tilde{s}}(v(b)) = \tilde{v}(d) = v(a) = v_0$$

At this point, I will switch to my notation because it clarified to me, at least, what was going on, in that a holonomy h_s is *invariant* under orientation-preserving reparametrization, and its inverse is well-defined.

For $\tilde{s} = s \circ t : [c, d] \rightarrow M$,
piecewise smooth t is reparametrized, i.e.

$$(38) \quad t : [c, d] \rightarrow [a, b]$$

Now,

$$\begin{aligned} \frac{d}{d\tau} \tilde{s}(\tau) &= \frac{d}{d\tau} \tilde{s}(t(\tau)) = \dot{s}(t) \frac{dt}{d\tau}(\tau) \equiv \dot{s} \frac{dt}{d\tau} \\ v^k(t) &= v^k(t(\tau)) = v^k(\tau) \\ \frac{dv^k}{d\tau}(t(\tau)) &= \frac{dv^k}{dt} \frac{dt}{d\tau} = \frac{dt}{d\tau} (-v^i(\tau) \dot{s}^j(t) \Gamma_{ij}^k) = -v^i(\tau) \frac{d\tilde{s}^j}{d\tau} \Gamma_{ij}^k \end{aligned}$$

Consider

$$h_s(v_0) = v(b)$$

If $t(c) = a$,
 $t(d) = b$

$$h_{\tilde{s}}(v_0) = \tilde{v}(d) = v(t(d)) = v(b) = h_s(v_0)$$

If $t(c) = b$,
 $t(d) = a$

$$\begin{aligned} h_{\tilde{s}}(h_s(v_0)) &= h_{\tilde{s}}(v(b)) = h_{\tilde{s}}(v(t(c))) = h_{\tilde{s}}(\tilde{v}(c)) = \\ &= \tilde{v}(d) = v(t(d)) = v(a) = v_0 \end{aligned}$$

Thus,

$$\boxed{h_{\tilde{s}} = h_s^{-1}}$$

□

I am working through Conlon (2008) [16], Clarke and Santoro (2012) [17], and Schreiber and Waldorf (2007)[18], concurrently, for holonomy.

10. PATH GROUPOID OF A SMOOTH MANIFOLD; GENERALIZATION OF PATHS

cf. Schreiber and Waldorf (2007)[18].

Definition 40 (path). ***path** is a smooth map $\gamma : [0, 1] \rightarrow M$, between 2 pts. $x, y \in M$, which has a sitting instant; i.e. number $0 < \epsilon < \frac{1}{2}$ s.t.*

$$(39) \quad \gamma(t) = \begin{cases} x & \text{for } 0 \leq t < \epsilon \\ y & \text{for } 1 - \epsilon < t \leq 1 \end{cases}$$

Denote the set of such paths by PM ,

$$(40) \quad PM \equiv \{\gamma \in \Gamma(M) \mid \text{smooth } \gamma : [0, 1] \rightarrow M \text{ s.t. } \exists 0 < \epsilon < \frac{1}{2} \text{ s.t. } \begin{cases} x & \text{for } 0 \leq t < \epsilon \\ y & \text{for } 1 - \epsilon < t \leq 1 \end{cases}\}$$

cf. Def. 2.1. of Schreiber and Waldorf (2007)[18]

Define *composition*:

Given paths γ_1, γ_2 ; $\gamma_1(0) = x$, $\gamma_2(0) = y$,

$$\gamma_1(1) = y \quad \gamma_2(1) = z$$

define composition to be path

$$(41) \quad \begin{aligned} &\gamma_2 \circ \gamma_1 \\ (\gamma_2 \circ \gamma_1)(t) &:= \begin{cases} \gamma_1(2t) & \text{for } 0 \leq t \leq \frac{1}{2} \\ \gamma_2(2t - 1) & \text{for } \frac{1}{2} \leq t \leq 1 \end{cases} \end{aligned}$$

$\gamma_2 \circ \gamma_1$ smooth since γ_1, γ_2 both constant near gluing pt., due to sitting instants ϵ_1, ϵ_2 , respectively.

Define *inverse*:

$$(42) \quad \begin{aligned} &\gamma^{-1} : [0, 1] \rightarrow M \\ \gamma^{-1}(t) &:= \gamma(1 - t) \end{aligned}$$

$$(\text{so that } \gamma^1(t) = \begin{cases} y & \text{for } 1 - \epsilon < 1 - t \leq 1 \text{ or } 0 \leq t < \epsilon \\ x & \text{for } 0 \leq 1 - t < \epsilon \text{ or } 1 - \epsilon < t \leq 1 \end{cases})$$

Definition 41 (thin homotopy equivalent). 2 paths γ_1, γ_2 s.t. $\gamma_1(0) = \gamma_2(0) = x$, γ_1, γ_2 are thin homotopy equivalent,

$$\gamma_1(1) = \gamma_2(1) = y$$

if \exists smooth $h : [0, 1] \times [0, 1] \rightarrow M$ s.t.

$$(1) \quad \exists 0 < \epsilon < \frac{1}{2} \text{ with}$$

$$(a) \quad h(s, t) = x \text{ for } 0 \leq t < \epsilon$$

$$h(s, t) = y \text{ for } 1 - \epsilon < t \leq 1$$

$$(b)$$

$$(c) \quad h(s, t) = \gamma_1(t) \text{ for } 0 \leq s < \epsilon$$

$$h(s, t) = \gamma_2(t) \text{ for } 1 - \epsilon < s \leq 1$$

$$(2) \quad \text{differential of } h \text{ has at most rank 1 everywhere, i.e.}$$

$$(43) \quad \text{rank}(dh|_{(s,t)}) \leq 1 \quad \forall (s, t) \in [0, 1] \times [0, 1]$$

cf. Def. 2.2. of Schreiber and Waldorf (2007)[18]

$h(s, t) = \gamma_1(t)$ for $0 \leq s < \epsilon$ is the homotopy from γ_1 to γ_2 , i.e. $h(0, t) = \gamma_1(t)$

$h(s, t) = \gamma_2(t)$ for $1 - \epsilon < s \leq 1$ $h(1, t) = \gamma_2(t)$

and define an equivalence relation on PM .

Note that for $h : [0, 1] \times [0, 1] \rightarrow M$,

$$(Dh)|_{(s,t)} = \left[\frac{\partial h^i}{\partial s}, \frac{\partial h^i}{\partial t} \right]$$

$P^1M \equiv$ set of thin homotopy classes of paths, i.e.

$$(44) \quad P^1M = \{[\gamma] \mid \gamma_1 \in PM, \text{ if } \exists \text{ smooth } h : [0, 1] \times [0, 1] \rightarrow M \text{ s.t. } h \text{ thin homotopy of } \gamma_1 \text{ and } \gamma_2, \gamma_1 \sim \gamma_2\}$$

$\text{pr} : PM \rightarrow P^1M$ is projection to classes.

Denote thin homotopy class of path γ , $\gamma(0) = x$, by $\overline{\gamma}$, or $[\gamma]$.

$$\gamma(1) = y$$

10.1. **Reparametrization of thin homotopies.** Let $\beta : [0, 1] \rightarrow [0, 1]$, $\beta(0) = 0$,
 $\beta(1) = 1$

Then \forall path γ , $\gamma(0) = x$, $\gamma \circ \beta$ is also a path $\gamma \circ \beta(0) = x$ and

$$\gamma(1) = y$$

$$\gamma \circ \beta(1) = y$$

$$(45) \quad h(s, t) := \gamma(t\beta(1 - s) + \beta(t)\beta(s))$$

defines a homotopy from γ to $\gamma \circ \beta$.

$$\gamma_1 \circ \gamma_2 \in PM \xrightarrow{\text{pr}} [\gamma_1 \circ \gamma_2] = [\gamma_1][\gamma_2] \in P^1M$$

Composition of thin homotopy classes of paths obeys following rules:

Lemma 3. \forall path γ , $\gamma(0) = x$

$$\gamma(1) = y$$

$$(1) \quad \overline{\gamma} \circ \overline{id_x} = \overline{\gamma} = \overline{id_y} \circ \overline{\gamma} \equiv [\gamma]1_x = [\gamma] = 1_y[\gamma]$$

$$(2) \quad \text{for paths } \gamma'; \gamma'(0) = y, \quad \gamma''(0) = z$$

$$\gamma'(1) = z \quad \gamma''(1) = w$$

$$(46) \quad (\overline{\gamma''} \circ \overline{\gamma'}) \circ \overline{\gamma} = \overline{\gamma''} \circ (\overline{\gamma'} \circ \overline{\gamma}) \equiv ([\gamma''][\gamma'])[\gamma] = [\gamma'']([\gamma'][\gamma])$$

$$(3) \quad \overline{\gamma} \circ \overline{\gamma}^{-1} = \overline{id_y} \text{ and } \overline{\gamma}^{-1} \circ \overline{\gamma} = \overline{id_x} \equiv [\gamma][\gamma^{-1}] = 1_y \text{ and } [\gamma^{-1}][\gamma] = 1_x$$

cf. Lemma 2.3. of Schreiber and Waldorf (2007)[18]

Definition 42 (path groupoid). \forall smooth manifold M ,
consider category whose set of objects is M ,

whose set of morphisms is P^1M , where class $[\gamma]$, $[\gamma](0) = x$ is a morphism from x to y and

$$[\gamma](1) = y$$

composition $[\gamma_1][\gamma_2] = [\gamma_1 \circ \gamma_2] \in P^1M$ Lemma 3 are axioms of a category, 3rd. property says \forall morphism is invertible.
Hence, we've defined a groupoid, called **path groupoid** of M , $\mathcal{P}_1(M)$.

So

$$\text{Obj}(\mathcal{P}_1(M)) = M$$

$$\text{Mor}(\mathcal{P}_1(M)) = P^1M$$

\forall smooth $f : M \rightarrow N$, denote functor f_*

$$(47) \quad f_* : \mathcal{P}_1(M) \rightarrow \mathcal{P}_1(N)$$

with

$$f_*(x) = f(x)$$

$$(f_*)([\gamma]) := [f \circ \gamma]$$

If $\gamma \sim \gamma'$, for $f \circ \gamma$, $f \circ \gamma'$,

$f \circ h(s, t)$ with $f \circ h(0, t) = f \circ \gamma(t)$,

$$f \circ h(1, t) = f \circ \gamma'(t)$$

so $f \circ h$ is a thin homotopy between $f \circ \gamma$, $f \circ \gamma'$ and so $[f \circ \gamma]$ well-defined.

Part 9. Complex Manifolds

EY : 20170123 I don't see many good books on Complex Manifolds for physicists other than Nakahara's. I will supplement this section on Complex Manifolds with external links to the notes of other courses that I found useful to myself.

[Complex Manifolds - Lecture Notes](#) Koppensteiner (2010) [13]

[Lectures on Riemannian Geometry, Part II: Complex Manifolds by Stefan Vandoren](#)

Vandoren (2008) [14]

Part 10. Jets, Jet bundles, h -principle, h -Prinzipien

cf. Eliashberg and Misahchev (2002) [19]

cf. Ch. 1 Jets and Holonomy, Sec. 1.1 Maps and sections of Eliashberg and Misahchev (2002) [19].

Visualize $f : \mathbb{R}^n \rightarrow \mathbb{R}^q$ as graph $\Gamma_f \subset \mathbb{R}^n \times \mathbb{R}^q$.

Consider this graph as image of $\mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^q$, i.e.

$$x \mapsto (x, f(x))$$

$\mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^q$ is called section (by mathematicians),

$$x \mapsto (x, f(x))$$

is called *field* or \mathbb{R}^q -valued field (by physicists).

cf. Ch. 1 Jets and Holonomy, Sec. 1.2 Coordinate definition of jets of Eliashberg and Misahchev (2002) [19].

Definition 43 (r -jet). Given (smooth) $f : \mathbb{R}^n \rightarrow \mathbb{R}^q$, given $x \in \mathbb{R}^n$.

r -jet of f at x - sequence of derivatives of f , up to order r , \equiv

$$(48) \quad J_f^r(x) = (f(x), f'(x) \dots f^{(r)}(x))$$

$f^{(q)}$ consists of all partial derivatives $D^\alpha f$, $\alpha = (\alpha_1 \dots \alpha_n)$, $|\alpha| = \alpha_1 + \dots + \alpha_n = s$, ordered lexicographically.

e.g. $q = 1$,

$f : \mathbb{R}^n \rightarrow \mathbb{R}$.

1-jet of f at $x = J_f^1(x) = (f(x), f^{(1)}(x))$.

$$f^{(1)}(x) = \{D^\alpha f | \alpha = (\alpha_1 \dots \alpha_n), |\alpha| = \alpha_1 + \dots + \alpha_n = 1\} = \left(\frac{\partial f}{\partial x^1}, \frac{\partial f}{\partial x^2}, \dots, \frac{\partial f}{\partial x^n} \right)$$

Let $d_r = d(n, r)$ = number of all partial derivatives D^α of order r of function $\mathbb{R}^n \rightarrow \mathbb{R}$.

Consider r -jet $J_f^r(x)$ of map $f : \mathbb{R}^n \rightarrow \mathbb{R}^q$ as pt. of space $\mathbb{R}^q \times \mathbb{R}^{qd_1} \times \mathbb{R}^{qd_2} \times \dots \times \mathbb{R}^{qd_r} = \mathbb{R}^{qN_r}$, where $N_r = N(n, r) = 1 + d_1 + d_2 + \dots + d_r$, i.e.

$$J_f^r(x) = (f(x), f^{(1)}(x), \dots, f^{(r)}(x)) \in \mathbb{R}^q \times \mathbb{R}^{qd_1} \times \dots \times \mathbb{R}^{qd_r} = \mathbb{R}^{qN_r}$$

Exercise 1.

Given order r , consider n -tuple of (positive) integers $(r_1, r_2 \dots r_n)$ s.t. $r_1 + r_2 + \dots + r_n = r$, and $r_k \geq 0$.

Imagine r_k = occupancy number, num ber of balls in k th cell. $(r_1 \dots r_n)$ describes a positive ocnfiguration of occupancy numbers, with indistinguishable balls; 2 distributions are distinguishable only if corresponding n -tuples $(r_1 \dots r_n)$ not identical.

Represent balls by stars, and indicate n cells by n spaces between $n + 1$ bars.

With $n+1$ bars, r stars, 2 bars are fixed. $n-1$ bars and r stars to arrange linearly, so a total of $n-1+r$ objects to arrange. r stars indistinguishable amongst themselves, so choose r out of $n-1+r$ to be stars.

$$(49) \quad \implies d_r = d(n, r) = \binom{n-1+r}{r}$$

Use *induction* (cf. **Ch. 4 Binomial Coefficients**).

$$N_0 = N(n, 0) = \binom{n-1+0}{0} = 1$$

$$N_1 = N(n, 1) = 1 + \binom{n-1+1}{1} = 1 + n = \frac{(n+1)!}{n!1!}$$

Induction step:

$$N_{r-1} = N(n, r-1) = \sum_{k=1}^{r-1} d_k + 1 = \binom{n+r-1}{r-1}$$

and so

$$N_r = N(n, r) = \sum_{k=1}^r d_k + 1 = \sum_{k=1}^r \binom{n-1+k}{k} + 1 = \sum_{k=1}^{r-1} \binom{n-1+k}{k} + \binom{n-1+r}{r} + 1 =$$

$$= \binom{n+r-1}{r-1} + \binom{n-1+r}{r} = \frac{(n+r-1)!}{(r-1)!n!} + \frac{(n-1+r)!}{r!(n-1)!} = \frac{(n+r)!}{n!r!} = \binom{n+r}{r}$$

$$\begin{array}{ccc} \mathbb{R}^{qN_r} & & J_f^r(x) \\ \uparrow J_f^r & & \uparrow J_f^r \\ \mathbb{R}^n & \xrightarrow{f} & \mathbb{R}^q \\ & & \downarrow \perp \\ & & x \end{array} \quad \begin{array}{ccc} & & f \\ & & \downarrow \perp \\ & & f(x) \end{array}$$

Definition 44 (space of r -jets). *space of r -jects of maps $\mathbb{R}^n \rightarrow \mathbb{R}^q$ or space of r -jets of sections $\mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^q \equiv$*

$$(50) \quad J^r(\mathbb{R}^n, \mathbb{R}^q) = \mathbb{R}^n \times \mathbb{R}^{qN_r} \equiv \mathbb{R}^n \times \mathbb{R}^q \times \mathbb{R}^{qd_1} \times \mathbb{R}^{qd_2} \times \dots \times \mathbb{R}^{qd_r}$$

e.g. $J^1(\mathbb{R}^n, \mathbb{R}^q) = \mathbb{R}^n \times \mathbb{R}^q \times M_{q \times n}$, where $M_{q \times n} = \mathbb{R}^{qn}$ is the space of $(q \times n)$ -matrices.

Part 11. Morse Theory

11. MORSE THEORY INTRODUCTION FROM A PHYSICIST

I needed some physical motivation to understand Morse theory, and so I looked at Hori, et. al. [15].
cf. pp. 43, Sec. 3.4 Morse Theory, from Ch. 3. Differential and Algebraic Topology of Hori, et. al. [15].
Consider smooth $f: M \rightarrow \mathbb{R}$, with non-degenerate critical points.

If no critical values of f between a and b ($a < b$), then subspace on which f takes values less than a is deformation retract of subspace where f less than b , i.e.

$$\{x \in M | f(x) < b\} \times [0, 1] \xrightarrow{F} \{x \in M | f(x) < b\}$$

$\forall x \in M$ s.t. $f(x) < b$,

$$\begin{array}{l} F(x, 0) = x \\ F(x, 1) \in \{x \in M | f(x) < a\} \end{array} \quad \text{and} \quad F(a', 1) = a' \quad \forall a' \in M \text{ s.t. } f(a') < a$$

To show this, consider $-\nabla f / |\nabla f|^2$

Morse lemma: \forall critical pt. p s.t. \exists choice of coordinates s.t.

$$(51) \quad f = -(x_1^2 + x_2^2 + \dots + x_\mu^2) + x_{\mu+1}^2 + \dots + x_n^2$$

where $f(p) = 0$ and p is at origin of these coordinates.

- difference between

$$f^{-1}(\{x \leq -\epsilon\}), f^{-1}(\{x \leq +\epsilon\})$$

can be determined by local analysis and only depends on μ , $\mu \equiv$ “Morse index” = number of negative eigenvalues of Hessian of f at critical pt.

Answer:

$f^{-1}(\{x \leq +\epsilon\})$ can be obtained from $f^{-1}(\{x \leq -\epsilon\})$ by “attaching μ -cell” along boundary $f^{-1}(0)$

- “attaching μ -cell to X mean, take μ -ball $B_\mu = \{|x| \leq 1\}$ in μ -dim. space, identity pts. on boundary $S^{\mu-1}$ with pts. in the space X , through cont. $f: S^{\mu-1} \rightarrow X$, i.e. take

$$X \coprod B_\mu$$

with $x \sim f(x) \quad \forall x \in \partial B_\mu = S^{\mu-1}$.

- find homology of M ,
 f defines chain complex C_f^* , k th graded piece C^{α_k} , α_k is number of critical pts. with index k .

$$(52) \quad \begin{array}{l} \partial: C_p^k \rightarrow C_p^{k-1} \\ \partial x_a = \sum_b \Delta_{a,b} x_b \end{array}$$

where $\Delta_{a,b} :=$ signed number of lines of gradient flow from x_a to x_b , b labels pts. of index $k-1$.

Gradient flow line is path $x(t)$ s.t. $\dot{x} = \nabla(f)$, with $x(-\infty) = x_a$

$$x(+\infty) = x_b$$

- To define this number ($\Delta_{a,b}?$), construct moduli space of such lines of flow (???)
by intersecting outward and inward flowing path spaces from each critical point, and then show this moduli space is oriented, 0-dim. manifold (pts. with signs)
- $\partial^2 = 0$ proof
 ∂ , boundary of space of paths connecting critical points, whose index differs by 2 = union over compositions of paths between critical pts. whose index differs by 1.
 \implies coefficients of ∂^2 are sums of signs of pts. in 0-dim. space, which is boundary of 1-dim. space.
These signs must therefore add to 0, so $\partial^2 = 0$.

Hori, et. al. [15] is good for physics, but there isn’t much thorough, step-by-step explanations of the math. I will look at Hirsch (1997) [6] and Shastri (2011) [5] at the same time.

11.1. Introduction, definitions of Morse Functions, for Morse Theory. cf. Ch. 6, Morse Theory of Hirsch (1997) [6], Section 1. Morse Functions, pp. 143-

Recall for TM , $T_x M \xrightarrow{\varphi} \mathbb{R}^n$.

Cotangent bundle T^*M defined likewise:

$$T_x^* M \xrightarrow{\varphi^*} \text{dual vector space } (\mathbb{R}^n)^* = L(\mathbb{R}^n, \mathbb{R})$$

i.e.

$$T^*M = \bigcup_{x \in M} (M_x^*) \quad M_x^* = L(M_x, \mathbb{R})$$

If chart (φ, U) on M , natural chart on T^*M is

$$\begin{array}{l} T^*U \rightarrow \varphi(U) \times (\mathbb{R}^n)^* \\ \lambda \in M_x^* \mapsto (\varphi(x), \lambda \varphi_x^{-1}) \end{array}$$

Projection map

$$\begin{aligned} p &: T^* \rightarrow M \\ M_x^* &\mapsto x \end{aligned}$$

Let C^{r+1} map, $1 \leq r \leq \omega$, $f : M \rightarrow \mathbb{R}$, $\forall x \in M$, linear map $T_x f : M_x \rightarrow \mathbb{R}$ belongs to M_x^*

$$T_x f = Df_x \in M_x^*$$

Then

$$\begin{aligned} Df &: M \rightarrow T^* M \\ x &\mapsto Df_x = Df(x) \end{aligned}$$

is C^r section of T^*M .

Definition 45. critical point x of f is zero of Df , i.e.

$$(53) \qquad Df(x) = 0$$

of vector space M_x^* .

Thus, set of critical pts. of f is counter-image of submanifold $Z^* \subset T^*M$ of zeros. Note $Z^* \approx M$, codim. of Z^* is $n = \dim M$.

Definition 46. Morse function f if \forall critical pts. of f are nondegenerate.

Note set of critical pts. closed discrete subset of M .

Let open $U \subset \mathbb{R}^n$, let C^2 map $g : U \rightarrow \mathbb{R}$,

critical pt. $p \in U$ nondegenerate iff

- linear $D(Dg)(p) : \mathbb{R}^n \rightarrow (\mathbb{R}^n)^*$ bijective
- identify $L(\mathbb{R}^n, (\mathbb{R}^n)^*)$ with space of bilinear maps $\mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$, \implies equivalent to condition that symmetric bilinear $D^2g(p) : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ non-degenerate
- $n \times n$ Hessian matrix

$$\left[\frac{\partial^2 g}{\partial x^i \partial x^j}(p) \right]$$

has rank n

Hessian of g at critical pt. p is quadratic form $H_p f$ associated to bilinear form $D^2g(p)$

$$\implies H_p f(y) = D^2g(p)(y, y) = \sum_{i,j} \frac{\partial^2 g}{\partial x^i \partial x^j}(p) y^i y^j$$

Let open $V \subset \mathbb{R}^n$, suppose C^2 diffeomorphism $h : V \rightarrow U$.

Let $q = h^{-1}(p)$, so q is critical pt. of $gh : V \rightarrow \mathbb{R}$.

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{H_q(gh)} & \mathbb{R} \\ \downarrow Dh(q) & \nearrow H_p g & \\ \mathbb{R}^n & & \end{array}$$

(quadratic) form $(H_p f)$ invariant under diffeomorphisms.

Let C^2 $f : M \rightarrow \mathbb{R}$.

\forall critical pt. x of f , define

Hessian quadratic form

$$H_x f : M_x \rightarrow \mathbb{R}$$

$$H_x f : M_x \xrightarrow{D\varphi_x} \mathbb{R}^n \xrightarrow{H_{\varphi(x)}(f\varphi^{-1})} \mathbb{R}$$

where φ is any chart at x .

Thus, critical pt. of a C^2 real-valued function nondegenerate iff associated Hessian quadratic form is nondegenerate.

Let Q nondegenerate quadratic form on vector space E .

Q negative definite on subspace $F \subset E$ if $Q(x) < 0$ whenever $x \in F$ nonzero.

Index of $Q \equiv \text{Ind}Q$, is largest possible dim. of subspace on which Q is negative definite.

cf. 1.1. Morse's Lemma of Ch. 6, pp. 145, Morse Theory of Hirsch (1997) [6]

Lemma 4 (Morse's Lemma). *Let $p \in M$ be nondegenerate critical pt. of index k of C^{r+2} map $f : M \rightarrow \mathbb{R}$, $1 \leq r \leq \omega$. Then $\exists C^r$ chart (φ, U) at p s.t.*

$$(54) \qquad f\varphi^{-1}(u_1 \dots u_n) = f(p) - \sum_{i=1}^k u_i^2 + \sum_{i=k+1}^n u_i^2$$

Let ${}^TQ \equiv Q^T$ denote tranpose of matrix Q .

Lemma 5. *Let $A = \text{diag}\{a_1, \dots, a_n\}$ diagonal $n \times n$ matrix, with diagonal entries ± 1 .*

Then \exists neighborhood N of A in vector space of symmetric $n \times n$ matrices, C^∞ map

$$(55) \qquad P : N \rightarrow GL(n, \mathbb{R})$$

s.t. $P(A) = I$, and if $P(B) = Q$, then $Q^T B Q = A$

Proof. Let $B = [b_{ij}]$ be symmetri matrix near A s.t. $b_{11} \neq 0$ and b_{11} has same sign as a_1 .

Consider $x = Ty$ where

$$\begin{aligned} x_1 &= \left[y_1 - \frac{b_{12}}{b_{11}} y_2 - \dots - \frac{b_{1n}}{b_{11}} y_n \right] / \sqrt{|b_n|} \\ x_k &= y_k \text{ for } k = 2, \dots n \end{aligned}$$

□

12. LAGRANGE MULTIPLIERS

From *wikipedia:Lagrange multiplier*, https://en.wikipedia.org/wiki/Lagrange_multiplier, find local minima (maxima), pt. $a \in N$, s.t. \exists neighborhood U s.t. $f(x) \geq f(a)$ ($f(x) \leq f(a)$) $\forall x \in U$.

For $f : U \rightarrow \mathbb{R}$, open $U \subset \mathbb{R}^n$, find $x \in U$ s.t. $D_x f \equiv Df(x) = 0$, check if Hessian $H_x f < 0$.

Maxima may not exit since U open.

References:

[Relative Extrema and Lagrange Multipliers](#)

Other interesting links:

[The Lagrange Multiplier Rule on Manifolds and Optimal Control of nonlinear systems](#)

Part 12. Classical Mechanics

13. CLASSICAL MECHANICS

13.1. Structure of Galilean Space-Time. cf. Sec. 3.1 - *Structure of Galilean Space-Time* of Prástaro (1996) [12].

Mechanics assumes a particular simple formulation if formulated with respect to some spacetime manifold.

In Galilean spacetime, it's possible to naturally recognize absolute objects, and others that depend on frames.

cf. Def. 3.1 of Prástaro (1996) [12]

Definition 47 (Galilean spacetime structure). (1) *Galilean spacetime structure* $:= (\mathcal{G}, g)$ where \mathcal{G} is (fiber bundle space-time)

$$(56) \qquad \mathcal{G} \equiv \{ \tau : M \rightarrow T \}$$

where

$M = 4\text{-dim. affine manifold}$ (**space-time**); corresponding structure is (M, \mathbf{M}, α) ,

(2) $T = 1$ -dim. affine space (**time**), corresponding affine structure is (T, \mathbf{T}, β)
 (3) $\tau =$ surjective affine mapping, of constant rank 1, s.t. $\forall p \in M$ associates its time $\tau(p) \in T$
 Put $\mathbf{S} = \ker(\underline{\tau}) \equiv \ker(D\tau) \in M$,
 where $\underline{\tau} \equiv D\tau$, D is symbol of derivative.
 Define

$$g : M \rightarrow vS_2^0(M) \equiv M \times S_2^0(\mathbf{S})$$

$$g(p) = (p, \underline{g}) \equiv (p, Dg), \forall p \in M$$

where $\underline{g} \equiv Dg$ is a Euclidean structure on \mathbf{S} . \underline{g} is called vertical metric field.

Thus, given (M, \mathbf{M}, α) , $\forall (O, \{\mathbf{e}_i\}_{1 \leq i \leq d}, \{\mathbf{e}_i\}_{i=1 \dots d})$, is basis of \mathbf{M} ,

$$M \cong \mathbb{R}^4, \text{ and } \exists \{x^\alpha : M = \mathbb{R}^4 \rightarrow \mathbb{R}\}_{\alpha=1 \dots 4}$$

13.2. Fundamental Theorems of (Classical) Dynamics. cf. Sec. 3.4 - *Fundamental Theorems of Dynamics* of Prástaro (1996) [12].

cf. Thm. 3.20 of Prástaro (1996) [12]

Theorem 19 (Momentum Theorem). *Variation of the free part of momentum of the observed motion of 1 body, in time interval $\Delta t \equiv [0, t]$ is equal to the corresponding impulse:*

$$(57) \quad I[0, t] \equiv \int_0^t F dt \equiv \left(\int_0^t F^j dt \right) \mathbf{e}_j$$

where $\{\mathbf{e}_j\}_{1 \leq j \leq 3}$ is a fixed basis of \mathbf{S}

$$(58) \quad \bar{p}_\psi(t) - \bar{p}_\psi(0) = I[0, t]$$

Proof.

$$\bar{p}_\psi = \mu \ddot{m}_\psi = \bar{f}_\psi \implies \dot{\bar{p}}_\psi = \dot{p}^j \mathbf{e}_j = F^j \mathbf{e}_j \implies \int_{[0, t]} \dot{p}^j dt = \int_{[0, t]} F^j dt$$

14. FLUID MECHANICS, FLUID FLOW

14.1. Mass Conservation for Fluid Flow, Continuum media. The mass of fluid in some volume $V_0 \subset N$ is $\int_{V_0} \rho \text{vol}^n$, where ρ is fluid density, $\rho \in C^\infty(N)$.

The total mass of fluid flowing out of volume V_0 is

$$\frac{d}{dt} \int_{V_0} \rho \text{vol}^n = \int_{V_0} \mathcal{L}_{\frac{\partial}{\partial t} + \mathbf{u}}(\rho \text{vol}^n) = \int_{V_0} \frac{\partial}{\partial t} \rho \text{vol}^n + \int_{V_0} \mathcal{L}_u \rho \text{vol}^n$$

$$\int_{V_0} \mathcal{L}_u \rho \text{vol}^n = \int_{V_0} di_{\mathbf{u}} \rho \text{vol}^n + i_{\mathbf{u}} d\rho \text{vol}^n = \int_{V_0} di_{\mathbf{u}} \rho \text{vol}^n + 0 = \int_{V_0} di_{\mathbf{u}} \rho \text{vol}^n = \int_{\partial V_0} i_{\mathbf{u}} \rho \text{vol}^n$$

Now

$$i_u \text{vol}^n = i_u \frac{\sqrt{g}}{n!} \epsilon_{i_1 \dots i_n} dx^{i_1} \wedge \dots \wedge dx^{i_n}$$

$$i_u dx^{i_1} \wedge \dots \wedge dx^{i_n} = u^{i_1} dx^{i_2} \wedge \dots \wedge dx^{i_n} - dx^{i_1} \wedge u^{i_2} dx^{i_3} \wedge \dots \wedge dx^{i_n} + \dots + (-1)^{n+1} dx^{i_1} \wedge \dots \wedge dx^{i_{n-1}} u^{i_n} = \epsilon_{j_1 \dots j_n}^{i_1 \dots i_n} u^{j_1} dx^{j_2} \wedge \dots \wedge dx^{j_n}$$

$$(59) \quad \implies i_u \text{vol}^n = \frac{\sqrt{g}}{(n-1)!} \epsilon_{j_1 \dots j_n} u^{j_1} dx^{j_2} \wedge \dots \wedge dx^{j_n}$$

We can also rewrite Eq. 59 to be a "surface differential":

$$(60) \quad \int_{V_0} \mathbf{d}i_{\mathbf{u}} \rho \text{vol}^n = \int_{\partial V_0} i_{\mathbf{u}} \rho \text{vol}^n =$$

$$\int_{\partial V_0} \rho \frac{\sqrt{g}}{(n-1)!} u^{j_1} \epsilon_{j_1 j_2 \dots j_n} dx^{j_2} \wedge \dots \wedge dx^{j_n} \equiv \int_{\partial V_0} \rho \mathbf{u} \cdot d\mathbf{S} \equiv \int_{\partial V_0} \rho \langle \mathbf{u}, d\mathbf{S} \rangle$$

If $\sqrt{g} = 1$, $n = 2$,

$$i_u \text{vol}^2 = (u^1 dx^2 - u^2 dx^1) = u \cdot n_1 dx^2 + u \cdot n_2 dx^1 = u \cdot ndS$$

with $n_1 = e_1$ and $n_2 = -e_2$.

Now

$$di_u \rho \text{vol}^n =$$

$$= \frac{\partial(\sqrt{g} \rho u^{j_1})}{\partial x^k} \frac{\epsilon_{j_1 \dots j_n}}{(n-1)!} dx^k \wedge dx^{j_2} \wedge \dots \wedge dx^{j_n} = \frac{\partial(\sqrt{g} \rho u^k)}{\partial x^k} \frac{\epsilon_{j_1 \dots j_n}}{n!} dx^{j_1} \wedge \dots \wedge dx^{j_n} = \frac{1}{\sqrt{g}} \frac{\partial(\sqrt{g} \rho u^k)}{\partial x^k} \text{vol}^n =$$

$$= \frac{\partial(\rho u^k)}{\partial x^k} \text{vol}^n + \rho u^k \frac{\partial \ln \sqrt{g}}{\partial x^k} \text{vol}^n = \text{div}(\rho u) \text{vol}^n + \rho u^k \frac{\partial \ln \sqrt{g}}{\partial x^k} \text{vol}^n$$

Now if $\sqrt{g} = 1$, then

$$\frac{d}{dt} \int_{V_0} \rho \text{vol}^n = \int_{V_0} \frac{\partial \rho}{\partial t} \text{vol}^n + \int_{V_0} di_u \rho \text{vol}^n = \int_{V_0} \frac{\partial \rho}{\partial t} \text{vol}^n + \int_{V_0} \text{div}(\rho u) \text{vol}^n \implies \frac{\partial \rho}{\partial t} + \text{div}(\rho u) = 0$$

which is the so-called mass continuity equation. $j = \rho u$ is the mass flux density.

Thus,

$$(61) \quad \square \quad \begin{aligned} & \text{(mass conservation)} \\ & m = m(t) := \int_{V_0} \rho \text{vol}^n, V_0 \subset N \\ & \dot{m} \equiv \frac{d}{dt} m(t) = \int_{V_0} \left(\frac{\partial \rho}{\partial t} \text{vol}^n + \mathbf{d}i_{\mathbf{u}} \rho \text{vol}^n \right) = \boxed{\int_{V_0} \frac{\partial \rho}{\partial t} \text{vol}^n + \int_{\partial V_0} \rho \mathbf{u} \cdot d\mathbf{S}} \\ & \text{if } \sqrt{g} = 1, \text{ and } \dot{m} = 0, \text{ then} \\ & \frac{\partial \rho}{\partial t} + \text{div}(\rho \mathbf{u}) = 0 \end{aligned}$$

TODO: 20190804 Frankel (2012) [23] in pp. 138 and onwards, for Sec. 4.3. Differentiation of Integrals posed the rightful question, "How does one compute the rate of change of an integral when the domain of integration is also changing?" Revisit the derivation from a Lie derivative and 1-parameter flow point of view.

Force should not be represented by a vector but rather by a 1-form. Then,

$$f \in \Omega^1(N), f = f(t, \mathbf{x}) = f_j dx^j \quad j = 1, \dots, n, \mathbf{x} = (x^1, \dots, x^n), \dim N = n$$

Indeed, the reason for f to be a 1-form is that we integrate differential forms, we don't integrate vectors.

$$(62) \quad W = \int_C f \quad (\text{line integral})$$

If f conservative, \exists scalar $U = U(x)$; $x \in N$ s.t. $f = -dU$.

Let $\mathbf{u} = u^j(t, \mathbf{x}) \frac{\partial}{\partial x^j} \in \mathfrak{X}(N)$ generate flow ϕ_t .

The Lie derivative along vector field \mathbf{u} , $\mathcal{L}_{\mathbf{u}}$, can be calculated in at least 2 ways: from the definition:

$$\mathcal{L}_{\mathbf{u}} \omega = \frac{d}{dt} \big|_{t=0} \phi_t^* \omega$$

or Cartan’s magic formula:

(63)

$$\begin{aligned}\mathcal{L}_{\mathbf{u}}\mathrm{vol}^n &= (i_{\mathbf{u}}\mathbf{d} + \mathbf{d}i_{\mathbf{u}})\mathrm{vol}^n = 0 + \mathbf{d}i_{\mathbf{u}}\mathrm{vol}^n = \mathbf{d}i_{\mathbf{u}}\left(\frac{\sqrt{g}}{n!}\epsilon_{i_1\dots i_n}dx^{i_1}\wedge\dots\wedge dx^{i_n}\right) = \\ &= \mathbf{d}\left(\frac{\sqrt{g}}{(n-1)!}\epsilon_{j_1\dots j_n}u^{j_1}dx^{j_2}\wedge\dots\wedge dx^{j_n}\right) = \frac{\partial(\sqrt{g}u^{j_1})}{\partial x^k}\frac{\epsilon_{j_1\dots j_n}}{(n-1)!}dx^k\wedge dx^{j_2}\wedge\dots\wedge dx^{j_n} = \\ &= \frac{\partial(\sqrt{g}u^{j_1})}{\partial x^k}\frac{\epsilon_{j_1\dots j_n}}{(n-1)!}\epsilon_{k_1,\dots,k_n}^{kj_2\dots j_n}dx^{k_1}\wedge\dots\wedge dx^{k_n}\underbrace{\left(\frac{\sqrt{g}}{\sqrt{g}}\right)}_{=1} = \frac{1}{\sqrt{g}}\frac{\partial(\sqrt{g}u^k)}{\partial x^k}\mathrm{vol}^n = \\ &= (\mathrm{div}\mathbf{u})\mathrm{vol}^n\end{aligned}$$

cf. <https://math.stackexchange.com/questions/2566381/lie-derivative-of-volume-form?rq=1>
Note that

$$\begin{aligned}\mathrm{div} : \mathfrak{X}(N) &\rightarrow C^\infty(N) \\ \mathcal{L}_{\mathbf{u}} : \Omega^n(N) &\rightarrow \Omega^n(N)\end{aligned}$$

Frankel (2012) [23] on pp. lvii of the ”Elasticity and Stresses” section offered this analogy: ”While work in particle mechanics pairs a force covector (f_i) with a contravariant tangent vector (dx^i/dt) to a curve, work done by traction in elasticity pairs the contravariant stress force 2-form \mathcal{S} with the covector valued deformation 1-form \mathcal{E} , to yield a scalar valued 3-form.

Instead of pushing along a curve (line) particle to do work, what does it mean to do work on a moving fluid element? One could possibly try to consider the deformation of a volume element under fluid flow. Frankel (2012) [23] in Sec. 4.2. The Lie Derivative of a Form, on pp. 132, asks ”If a flow deforms some attribute, say volume, how does one measure the deformation? $\mathcal{L}_{\mathbf{u}}\mathrm{vol}^n$ ”is the n -form that reads off the rate of change of volume of a parallelopiped spanned by n vectors that are pushed forward by the flow ϕ_t .” So, ”in other words, $\mathcal{L}_{\mathbf{u}}\mathrm{vol}^n$ measures how volumes are changing under the flow ϕ_t generated by \mathbf{X} ” on pp. 133 on Frankel (2012) [23].

Then rate of work done on a fluid element could be the following:

$$W = \int_V \mathcal{L}_{\mathbf{u}}(p\mathrm{vol}^n) = \int_V \mathbf{d}i_{\mathbf{u}}p\mathrm{vol}^n = \int_{\partial V} i_{\mathbf{u}}p\mathrm{vol}^n = \int_{\partial V} p\mathbf{u} \cdot d\mathbf{S}$$

TODO: work on a fluid element?

15. THERMODYNAMICS

Let Σ be a (topological) manifold.
Suppose U is a global coordinate on Σ :

(64)

(First Law of Thermodynamics (energy conservation))

$$\boxed{dU = Q - W \text{ or } Q = dU + W}$$

where $dU, Q, W \in \Omega^1(\Sigma)$ (i.e. dU, Q, W are 1-forms over manifold Σ).

Consider a path in Σ , $\gamma, \gamma : \mathbb{R} \rightarrow \Sigma$,
and using a chart (U, S^1, \dots, S^n) (e.g. $n = 1, S^1 = v$ for volume)

$$\begin{aligned}\gamma(t) &= (U(t), S^1(t), \dots, S^n(t)) \\ \dot{\gamma} \in \mathfrak{X}(\Sigma), \dot{\gamma} &= \dot{U}\frac{\partial}{\partial U} + \dot{S}^i\frac{\partial}{\partial S^i}\end{aligned}$$

Now

$$\begin{aligned}dU(\dot{\gamma}) &= \dot{\gamma}(U) = \dot{U}\frac{\partial}{\partial U}U + 0 = \dot{U} \\ Q(\dot{\gamma}) &= Q(t)dt\left(\dot{\gamma}\frac{\partial}{\partial t}\right) = Q(t)\dot{\gamma} \\ \implies \dot{U} &= Q(t)\dot{\gamma} - W(t)\dot{\gamma}\end{aligned}$$

Recall that for enthalpy $H, H := U + pV, H = H(\sigma, p)$
TODO 20190804 Derive and check convection form of enthalpy against both Kittle and Kroemer plus thermodynamics and Sonntag, et. al.
Incomplete:

$$\begin{aligned}dH &= Q + Vdp \\ W = pdV &= pdV + Vdp - Vdp = d(pV) - Vdp = \\ dE &= Q - W + \mu dH \\ dE &= Q - W + \hat{h}dN\end{aligned}$$

$U \rightarrow E$ notation is to promote the internal energy U to include kinetic and potential energies, so that possibly, $E = U + \text{K.E.} + \text{P.E.}$, or, i.e., E includes internal energy and mechanical energy.

$$\begin{aligned}G &= \mu N \\ G = U - \tau\sigma + pV &\xrightarrow{U \rightarrow E} G = E - \tau\sigma + pV \implies \tau\check{h} \\ dE &= Q + W + \mu dN = Q + W + (\check{h} - \tau\check{\sigma})dN \\ Q &= \tau d\sigma = \tau d(N\check{\sigma}) = \tau Nd\check{\sigma} + \tau\check{\sigma}dN\end{aligned}$$

$\tau Nd\check{\sigma}$ is the entropy change due to change in entropy per particle; i.e. **conduction term**
 $\tau\check{\sigma}dN$ is entropy change due to change in number of particles, i.e. **convection term**

$$dE = Q + W + \check{h}dN - \tau\check{\sigma}dN = \tau Nd\check{\sigma} + \tau\check{\sigma}dN + W + \check{h}dN - \tau\check{\sigma}dN = \tau Nd\check{\sigma} + W + \check{h}dN$$

$m(t) = MN$
Assume only 1 chemical species:

$$\begin{aligned}\check{Q} &:= \tau Nd\check{\sigma} \quad \hat{h} := \frac{H}{m} = \frac{H}{MN} \\ \implies dE &= \check{Q} + W + \check{h}dN\frac{M}{M} = \check{Q} + W + \hat{h}dm\end{aligned}$$

The Gibbs free energy equilibrium is given by,

$$dG = \mu dN - \sigma d\tau + Vdp$$

For a throttling valve (don’t all valves throttle?) the pressure drop is accounted for by the Gibbs free energy, *not* by an isenthalpy condition.

Part 13. Classical Mechanics applications

cf. Arnold, Kozlov, Neishtadt (2006) [21].
If known forces $\mathbf{F}_1 \dots \mathbf{F}_n$ acts on points, then

(65)

$$\sum_{i=1}^n \langle m_i \ddot{\mathbf{r}}_i - \mathbf{F}_i, \xi_i \rangle = 0$$

cf. Eq. (1.26) of Arnold, Kozlov, Neishtadt (2006) [21], where $\xi_1, \dots \xi_n$ are arbitrary tangent vectors to $M, \xi_i, \dots \xi_n \in TM$.
 $\sum_{i=1}^n \langle m_i \ddot{\mathbf{r}}_i - \mathbf{F}_i, \xi_i \rangle$ called ”general equation of dynamics” or d’Alembert-Lagrange principle.

16. ORBITAL MECHANICS

16.1. **Conic sections.** The following is an amalgamation or summary of the development in Ch. 13, Sections 13.18-22, from pp. 497, Apostol (1991) [25]

Definition 48 (General definition of conic sections (vector-formulated definition)). *Given line L , called the directrix, pt. F , the focus, such that $F \notin L$,*

eccentric $e > 0$,

$d =$ distance of L from F , then

*the **conic section** $C = \{X\}$ (set of all points) such that*

$$(66) \quad \|X - F\| = ed(X, L)$$

Now,

$e < 1$ ellipse,

$e = 1$ parabola,

$e > 1$ hyperbola.

Further, if N unit normal to L , then $\forall P \in L$,

$$(67) \quad d(X, L) = |(X - P) \cdot N|$$

If $(X - P) \cdot N \geq 0$, $X \in$ positive half plane (negative half-plane).

Theorem 20 (Further conic section development). *Building upon Def. 48,*

If $F \in$ positive half plane determined by N , then $F - dN \in L$.

If $F \in$ negative half plane determined by N , then $F + dN \in L$.

Choose this convenient, specific point: $F + dN = P \in L \implies \|X - F\| = e|(X - (F + dN)) \cdot N|$

If $F = 0$, $\|X\| = e|X \cdot N - d| = r = e(d - r \cos \theta)$. Then

$$(68) \quad \frac{ed}{r} = 1 + e \cos \theta \text{ or } \boxed{r = \frac{ed}{1 + e \cos \theta}} \text{ for } F = 0$$

Consider symmetry about the origin (i.e. of $X, \exists -X$).

As a caveat, let $A = e(d + F \cdot N)$. Then

$$\|X - F\| = e|(X - (F + dN)) \cdot N| = ed(X, L) = e|X \cdot N - \frac{A}{e}|$$

Then

$$(69) \quad \|X - F\|^2 = \|X\|^2 - 2X \cdot F + \|F\|^2 = e^2((X \cdot N)^2 - 2\frac{A}{e}X \cdot N + \left(\frac{A}{e}\right)^2)$$

Using the symmetry about the origin, $X \rightarrow -X$,

$$\implies X \cdot F = AeX \cdot N \text{ or } X \cdot (F - AeN) = 0$$

Then $F = AeN$. Note that $A = e(F \cdot N \pm d)$ (for F in negative (positive) half-plane).

Then

$$\begin{aligned} Ae &= e^2(Ae + d) \text{ or } \frac{A}{e} = Ae + d \text{ or } A\left(\frac{1}{e} - e\right) = d \text{ or } A = \frac{ed}{1 - e^2} \\ \implies \|X - AeN\| &= e|(X - (AeN + dN)) \cdot N| = e|X \cdot N - (Ae + d)| \end{aligned}$$

Let $X = aN$. Plug this into Eq. 69.

$$\begin{aligned} a^2 - 2aAe + A^2e^2 &= e^2(a^2 - \frac{2A}{e}a + \left(\frac{A}{e}\right)^2) = e^2a^2 - 2Aae + A^2 \implies a = A \\ \implies a &= \frac{ed}{1 - e^2} \end{aligned}$$

Thus, we've shown that A is equal to the semi-major axis a and that

$$(70) \quad \boxed{F = aeN}$$

Thus, the semi-major axis is obtained when $X = \pm aN$ (vertices), so that $\|X\|^2 + (ae)^2 = e^2(X \cdot N)^2 + a^2$ is satisfied. Likewise, the semi-minor axis is obtained if $X = \pm bN_\perp$, so that

$$(71) \quad b^2 + (ae)^2 = 0 + a^2 \implies \boxed{b = a\sqrt{1 - e^2}}$$

Compare this against (26.6) of Greiner (2004) [20]:

$$(72) \quad r = \frac{ed}{1 + e \cos \theta}, \quad e = \frac{C\hat{L}_0^2}{GM}, \quad d = 1/C$$

cf. Eq. (26.32) of Greiner (2004) [20].

Investigate which physical quantities (e.g. E, L) eccentricity e depends on.

$$\begin{aligned} u &= \frac{GM}{\hat{L}_0^2} + C \cos(\theta - \theta_0) \implies \frac{du}{d\theta} = -C \sin(\theta - \theta_0) \\ \frac{1}{2}mL_0^2 \left[C^2 \sin^2(\theta - \theta_0) + \left(\frac{GM}{L_0^2}\right)^2 + 2C\frac{GM}{L_0^2} \cos(\theta - \theta_0) + C^2 \cos^2(\theta - \theta_0) \right] &= \\ \implies &= \frac{1}{2}m\hat{L}_0^2 \left[C^2 + \left(\frac{GM}{\hat{L}_0^2}\right)^2 + 2C\frac{GM}{\hat{L}_0^2} \cos(\theta - \theta_0) \right] = \\ &= E - V\left(\frac{1}{u}\right) = E + GMmu = E + \frac{(GM)^2}{\hat{L}_0^2}m + CGMm \cos(\theta - \theta_0) \\ \implies \frac{1}{2}m\hat{L}_0^2 C^2 &= E + \frac{1}{2}\frac{(GM)^2}{\hat{L}_0^2}m \end{aligned}$$

Solve for the dummy constant C :

$$(73) \quad C = \sqrt{\frac{2E}{m\hat{L}_0^2} + \frac{(GM)^2}{\hat{L}_0^4}}$$

cf. Eq. (26.36) of Greiner (2004) [20].

From Eq. (26.32) of Greiner (2004) [20] or Eq. 72,

$$e = \frac{C\hat{L}_0^2}{GM} = \sqrt{\frac{2E}{m\hat{L}_0^2} + \frac{(GM)^2}{\hat{L}_0^4}} \left(\frac{\hat{L}_0^2}{GM}\right)$$

So

$$(74) \quad \boxed{e = \sqrt{\frac{2E\hat{L}_0^2}{m(GM)^2} + 1}}$$

cf. Eq. (26.37) of Greiner (2004) [20].

Hence, shape of the path depends on total energy E , angular momentum $L_0 = m\hat{L}_0$ of moving body (recall that $\hat{L} := \mathbf{r} \times \mathbf{v}$). Given Eq. [76](#) for a , plug in for E :

It holds that for a

$$\begin{aligned} \text{parabola } e = 1, & & E = 0 \\ \text{ellipse } 0 < e < 1, & & E < 0, \quad \frac{(GM)^2 m}{-2\hat{L}_0^2} < E < 0 \\ \text{circle } e = 0 & & E = \frac{-(GM)^2 m}{2\hat{L}_0^2} \\ \text{hyperbola } e = 1, & & E > 0 \end{aligned}$$

The general form of initial energy E_0 is, recall, give by

$$E_0 = \frac{1}{2}mv^2 + V(r) = \frac{1}{2}mv^2 - \frac{GMm}{r_0} = m \left(\frac{v^2}{2} - \frac{GM}{r_0} \right)$$

Notice that for the case of the parabola that $E_0 = 0$ when

$$(75) \quad v = \sqrt{\frac{2GM}{r_0}}$$

Eq. [75](#) is the expression for the **escape velocity**.

Knowing what we know now, let's rewrite some of the expressions from before:

$$\begin{aligned} r(\theta) &= \frac{\hat{L}_0^2/GM}{1 + \frac{C\hat{L}_0^2}{GM} \cos(\theta - \theta_0)} \\ \hat{L}_0^2 &:= r_0^2 \dot{\theta}_0 \\ C &= \sqrt{\frac{2E}{m\hat{L}_0^2} + \frac{(GM)^2}{\hat{L}_0^4}} \end{aligned}$$

16.2. Path parameters (i.e. orbital parameters). cf. pp. 262, Greiner (2004) [\[20\]](#).

$$a = \frac{1}{2}(r(\theta = 0) + r(\theta = \pi)) = \frac{1}{2} \left[\frac{\hat{L}_0^2/GM}{1+e} + \frac{\hat{L}_0^2/GM}{1-e} \right] = \frac{\hat{L}_0^2/GM}{1-e^2}$$

The addition above for a is needed because we center origin at a focus, for this expression.

$$(76) \quad a = \frac{\hat{L}_0^2/GM}{\frac{-2E\hat{L}_0^2}{m(GM)^2}} = \frac{GMm}{-2E}$$

cf. Eq. (26.39) of Greiner (2004) [\[20\]](#).

$$(77) \quad b = \sqrt{a^2 - f^2} = \sqrt{a^2 - e^2 a^2} = a\sqrt{1 - e^2} = a\sqrt{\frac{-2E\hat{L}_0^2}{m(GM)^2}} = \hat{L}_0\sqrt{\frac{-m}{2E}}$$

cf. Eq. (26.40) of Greiner (2004) [\[20\]](#).

We can also calculate the perigee (point of closest approach to mass M):

$$(78) \quad r(\theta = 0) = \frac{\hat{L}_0^2/GM}{1+e} = \frac{\hat{L}_0^2/GM}{1 + \sqrt{\frac{2E\hat{L}_0^2}{m(GM)^2} + 1}}$$

$$(79) \quad a = \frac{-GMm}{2E} = -\frac{GMm}{2\left(\frac{1}{2}mv_0^2 - \frac{GMm}{r}\right)} = \frac{-GM}{v_0^2 - \frac{2GM}{r}} = \frac{-GM}{v_0^2\left(1 - \frac{2GM}{rv_0^2}\right)}$$

Notice when the denominator is 0. It is when v_0 is the the escape velocity (cf. Eq. [75](#)).

16.2.1. Eccentricity from r_0, \mathbf{v}_0 . Consider being given \mathbf{r}, \mathbf{v} , with \mathbf{r} in a coordinate frame with its origin at the center of mass (or, if M large enough, from the focus where M is centered at).

The first step is to use a coordinate *frame* (set of orthonormal unit vectors) with \mathbf{r}, \mathbf{v} forming a *plane*. Note this important fact from geometry: a plane is spanned by 2 non-collinear vectors. So \mathbf{r}, \mathbf{v} better not be parallel to each other.

Let

$$\begin{aligned} \mathbf{v} &= v^x \mathbf{e}_x + v^y \mathbf{e}_y + v^z \mathbf{e}_z = v^r \mathbf{e}_r + v^\varphi \mathbf{e}_\varphi \\ \mathbf{r} &= x \mathbf{e}_x + y \mathbf{e}_y + z \mathbf{e}_z = r \mathbf{e}_r \end{aligned}$$

Now

$$(\mathbf{r} \times \mathbf{v})_i = \epsilon_{jki} r_j v_k$$

To get the vector *orthogonal* to the plane that is formed by \mathbf{r}, \mathbf{v} ,

$$\begin{aligned} ((\mathbf{r} \times \mathbf{v}) \times \mathbf{r})_m &= \epsilon_{ijm} \epsilon_{jki} r_j v_k r_l = \epsilon_{ilm} \epsilon_{ijk} r_j v_k r_l = \text{or } ((\mathbf{r} \times \mathbf{v}) \times \mathbf{r}) = r^2 \mathbf{v} - (\mathbf{v} \cdot \mathbf{r}) \mathbf{r} \\ &= r^2 v_m - v_k r_k r_m = r^2 v_m - v_k r_k r_m \end{aligned}$$

Then

$$\hat{L}_0 = r^2 \dot{\theta} = r \left(\mathbf{v} \cdot \frac{((\mathbf{r} \times \mathbf{v}) \times \mathbf{r})}{|(\mathbf{r} \times \mathbf{v}) \times \mathbf{r}|} \right) = r \left(\frac{r^2 v^2 - (\mathbf{v} \cdot \mathbf{r})^2}{|r^2 \mathbf{v} - (\mathbf{v} \cdot \mathbf{r}) \mathbf{r}|} \right)$$

Thus, we can find the following:

$$e^2 = 1 + \frac{2E\hat{L}_0^2}{m(GM)^2} = 1 + \frac{2\left(\frac{1}{2}mv_0^2 - \frac{GMm}{r}\right)\hat{L}_0^2}{m(GM)^2} = 1 + \frac{(v_0^2 - \frac{2GMm}{r})\hat{L}_0^2}{(GM)^2}$$

$$a = \frac{-GMm}{2E} = -\frac{GMm}{2\left(\frac{1}{2}mv_0^2 - \frac{GMm}{r}\right)} = \frac{-GM}{v_0^2 - \frac{2GM}{r}} = \frac{-GM}{v_0^2\left(1 - \frac{2GM}{rv_0^2}\right)}$$

TODO: Calculate period of revolution.

Here are some calculations (forms) that can help with further calculations:

$$(80) \quad \hat{L}_0^2 = b^2 \left(\frac{2E}{-m} \right) \text{ or } \frac{\hat{L}_0^2}{GM} = b^2 \left(\frac{2E}{-GMm} \right) = \frac{b^2}{a} = a(1 - e^2)$$

using Eq. [71](#), for $e^2 = 1 - \left(\frac{b}{a}\right)^2$.

16.3. Path in coordinates. For \mathbf{r} centered at focus F and \mathbf{r}, \mathbf{v} expressed in coordinates in the plane of the conic section path, i.e. in the plane formed by \mathbf{r}, \mathbf{v} , with the axes expressed as $\frac{\partial}{\partial x}, \frac{\partial}{\partial y}$,

$$(81) \quad \mathbf{r} = r \cos \theta \frac{\partial}{\partial x} + r \sin \theta \frac{\partial}{\partial y}$$

Compare this to Eq. (2.5-1) of Sec. 2.5.1 "Expressing r and v in the Perifocal System" of Bate, Mueller, White (1971) [26].

Recall the equation for the conic section in polar coordinates, with the coordinates centered at focus F :

$$(82) \quad r = \frac{a(1 - e^2)}{1 + e \cos \theta}$$

Compare this to Eq. (1.5-4) of Bate, Mueller, White (1971) [26].

Now

$$(83) \quad \mathbf{v} = \dot{\mathbf{r}} = (\dot{r} \cos \theta - r \dot{\theta} \sin \theta) \frac{\partial}{\partial x} + (\dot{r} \sin \theta + r \dot{\theta} \cos \theta) \frac{\partial}{\partial y}$$

Compare this to the equation on pp. 72, of Sec. 2.5.1 "Expressing r and v in the Perifocal System" of Bate, Mueller, White (1971) [26].

Now

$$(84) \quad \begin{aligned} \dot{r} &= \frac{a(1 - e^2)}{(1 + e \cos \theta)^2} e \sin \theta \dot{\theta} = \frac{r^2}{a(1 - e^2)} e \sin \theta \frac{\hat{L}_0}{r^2} = \frac{\hat{L}_0 e \sin \theta}{a(1 - e^2)} = \frac{\hat{L}_0 e \sin \theta}{\frac{-GMm}{2E} \left(\frac{L_0^2}{\left(\frac{(GM)^2 m}{-2E} \right)} \right)} \\ &\implies \dot{r} = \frac{GM}{\hat{L}_0} e \sin \theta \end{aligned}$$

Compare this to Eq. (2.5-2) of Sec. 2.5.1 "Expressing r and v in the Perifocal System" of Bate, Mueller, White (1971) [26].

Then

$$(85) \quad \begin{aligned} &\implies \frac{a(1 - e^2)}{(1 + e \cos \theta)^2} \dot{\theta} = \frac{GM}{\hat{L}_0} \text{ or} \\ &r \dot{\theta} = \frac{GM}{\hat{L}_0} (1 + e \cos \theta) \end{aligned}$$

Compare this to Eq. (2.5-3) on pp. 73 of Sec. 2.5.1 "Expressing r and v in the Perifocal System" of Bate, Mueller, White (1971) [26].

Then

$$(86) \quad \begin{aligned} \mathbf{v} &= \frac{GM}{\hat{L}_0} \left[(e \sin \theta \cos \theta - (1 + e \cos \theta) \sin \theta) \frac{\partial}{\partial x} + (e \sin^2 \theta + (1 + e \cos \theta) \cos \theta) \frac{\partial}{\partial y} \right] = \\ &\mathbf{v} = \frac{GM}{\hat{L}_0} \left[\sin \theta \frac{\partial}{\partial x} + (e + \cos \theta) \frac{\partial}{\partial y} \right] = \frac{GM}{\hat{L}_0} \left[\frac{y}{r} \frac{\partial}{\partial x} + \left(e + \frac{x}{r} \right) \frac{\partial}{\partial y} \right] \end{aligned}$$

Compare this to Eq. (2.5-4) on pp. 73 of Sec. 2.5.1 "Expressing r and v in the Perifocal System" of Bate, Mueller, White (1971) [26].

Consider this ansatz from subsection 4.4.3 "The f and g expressions from Ch. 4 section on "Position and velocity - A function of time" in Bate, Mueller, White (1971) [26].

Because \mathbf{r}, \mathbf{v} lie in the plane (i.e. coplanar) containing the entire conic section trajectory, then make the following ansatz:

$$(87) \quad \begin{aligned} \mathbf{r} &= f \mathbf{r}_0 + g \mathbf{v}_0 \\ \dot{\mathbf{r}} &=: \mathbf{v} = \dot{f} \mathbf{r}_0 + \dot{g} \mathbf{v}_0 \end{aligned}$$

Solve for f and g by taking the appropriate cross products:

$$(88) \quad \begin{aligned} \mathbf{r} \times \mathbf{v}_0 &= f \mathbf{r}_0 \times \mathbf{v}_0 \\ \mathbf{r} \times \mathbf{r}_0 &= g \mathbf{v}_0 \times \mathbf{r} \end{aligned}$$

16.4. Eccentricity anomaly; Eccentricity anomaly coordinates. Let the point of interest be P. Choose the origin for the coordinates (x, y) to have *its origin* at the *center of the ellipse*.

Let point C be the point at the center of the ellipse (origin), let point A be the projection of point P onto the x -axis, point F be the point at the focus.

Recall that the length from the center C to the focus, \overline{CF} , is ae , where a is the semi-major axis, and eccentricity is e .

The eccentricity anomaly E (it's just an angle) is given by

$$(89) \quad \cos E = \frac{x}{a}$$

Remember E is *not* the angle formed by triangle CAP, but rather, first take the line through P and A. This line is perpendicular to the x-axis. Then intersect that line with a circle, centered at C, of radius a . Then, the line from C to that intersection forms an angle E with the x -axis.

By using the ellipse equation and trigonometry,

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \implies \cos^2 E + \frac{y^2}{b^2} = 1 \text{ or } \frac{y^2}{b^2} = \sin^2 E \text{ or}$$

$$(90) \quad \sin E = \frac{y}{b}$$

Now

$$\overline{AF} = \overline{CF} - \overline{CA} = ae - a \cos E$$

Use the right triangle PAF and the hypotenuse being r (distance from the focus F to point P) to find

$$\overline{AF} = r \cos(\pi - \theta) = -r \cos \theta$$

So

$$\cos \theta = \frac{a(\cos E - e)}{r}$$

Then, plug into the conic section equation, Eq. 68,

$$\begin{aligned} r &= \frac{ed}{1 + e \cos \theta} = \frac{a(1 - e^2)}{1 + e \cos \theta} = \frac{b^2/a}{1 + e \cos \theta} = \frac{b^2/a}{1 + e \left(\frac{a(\cos E - e)}{r} \right)} \\ &\implies 1 = \frac{b^2/a}{r + e(a(\cos E - e))} \\ \implies r &= \frac{b^2}{a} - e(a(\cos E - e)) = \frac{b^2}{a} + ae^2 - ea \cos E = a(1 - e \cos E) \end{aligned}$$

$$(91) \quad \boxed{r = a(1 - e \cos E)}$$

Compare this to Eq. (4.6-9) on pp. 215 of Sec. 4.6 in "The Kepler Problem" of Bate, Mueller, White (1971) [26].

For the coordinates with the origin at the *focus* F , it is given by

$$(92) \quad \begin{aligned} x &= a(\cos E - e) \\ y &= b \sin E = a\sqrt{1 - e^2} \sin E \end{aligned}$$

Compare this to Eqns. (4.6-7), (4.6-8) on pp. 215 of Sec. 4.6 in "The Kepler Problem" of Bate, Mueller, White (1971) [26].

Indeed, for $E = 0$, $x = a - ae$ (the distance from the focus F to the vertex is $a - ae$), $E = \pi/2$, $x = -ae$ (the distance from the focus F and back to the center C) and for $E = \pi$, $x = -a - ae$ (the distance to the other vertex).

16.4.1. *Path in terms of eccentric anomaly.* Take the time derivative of Eq. 92:

$$(93) \quad \begin{aligned} \dot{x} &= -a \sin E \dot{E} \\ \dot{y} &= a \sqrt{1 - e^2} \cos E \dot{E} \end{aligned}$$

To determine \dot{E} , consider both expressions for the radial distance r for Eq. 91 and Eq. 68:

$$\begin{aligned} r &= \frac{a(1 - e^2)}{1 + e \cos \theta} = \frac{b}{1 + e \cos \theta} \\ \implies \dot{r} &= \frac{-b}{(1 + e \cos \theta)^2} (-e \sin \theta) \dot{\theta} = \frac{be \sin \theta \dot{\theta}}{(1 + e \cos \theta)^2} = \\ \dot{r} &= \frac{r^2 e}{b} \sin \theta \dot{\theta} \\ r &= a(1 - e \cos E) \\ \implies \dot{r} &= +ae \sin E \dot{E} \\ \implies ae \sin E \dot{E} &= \frac{r^2 e}{b} \sin \theta \dot{\theta} \text{ or } a \sin E \dot{E} = \frac{r^2}{b} \sin \theta \dot{\theta} = \frac{\hat{L}_0 \sin \theta}{b} = \frac{\hat{L}_0}{b} \frac{y}{r} \\ \implies \dot{E} &= \frac{\hat{L}_0}{ab} \frac{y}{r \sin E} = \frac{\hat{L}_0}{ar} = \frac{b}{a^{3/2} r} \end{aligned}$$

Compare the expression for \dot{E} with Eq. (4.6-11) on pp. 216 of Bate, Mueller, White (1971) [26].

Armed with an expression for \dot{E} , plug this back into the expression for \dot{r} :

$$(94) \quad \dot{r} = ae \sin E \left(\frac{\hat{L}_0}{ar} \right) = \frac{e \sin E \hat{L}_0}{r}$$

Compare this to the equation on pp. 216, "Position and Velocity - a function of time" in Ch. 4 of Bate, Mueller, White (1971) [26].

Use the expressions above in Eqns. 88. For \mathbf{r}, \mathbf{r}_0 centered at focus F:

$$\begin{aligned} xv_{0,y} - yv_{0,x} &= a(\cos E - e)b \cos E_0 \dot{E}_0 - b \sin E(-a \sin E_0 \dot{E}_0) = \\ &= ab \dot{E}_0 [(\cos E - e) \cos E_0 + \sin E \sin E_0] = \frac{b^2}{\sqrt{ar}} [\cos(E - E_0) - e \cos E_0] = \frac{\hat{L}^2 \left(\frac{-m}{2E} \right)}{\sqrt{GM} \sqrt{\frac{m}{-2E}} r} [\cos(E - E_0) - e \cos E_0] = \\ &= \frac{\hat{L}_0 b}{r \sqrt{GM}} [\cos(E - E_0) - e \cos E_0] \end{aligned}$$

Comparing this against the "right-hand side (RHS)" with $f(\mathbf{r}_0 \times \mathbf{v}_0)$, the RHS has a magnitude of $f \hat{L}_0$ in the same direction as $(\mathbf{r} \times \mathbf{v}_0)$. Thus

$$(95) \quad f = \frac{b}{r \sqrt{GM}} [\cos(E - E_0) - e \cos E_0]$$

For the expression involving g in Eqns. 88,

$$\mathbf{r} \times \mathbf{r}_0 = xy_0 - yx_0 = a(\cos E - e)b \sin E_0 - b \sin E a(\cos E_0 - e) = -ab [(\cos E_0) \sin E - (\cos E - e) \sin E_0]$$

with

$$ab = a \hat{L}_0 \sqrt{\frac{-m}{2E}} = a \hat{L}_0 \left(\frac{a}{GM} \right)^{1/2} = \left(\frac{a^3}{GM} \right)^{1/2} \hat{L}_0$$

For the RHS with $g\mathbf{v}_0 \times \mathbf{r}_0$, $\mathbf{v}_0 \times \mathbf{r}_0$, by the right-hand rule or the right-hand orientation, it is in the *opposite* direction as $\mathbf{r} \times \mathbf{r}_0$. Thus, apply a negative side on the RHS.

$$(96) \quad \implies g = \left(\frac{a^3}{GM} \right)^{1/2} [(\cos E_0 - e) \sin E - (\cos E - e) \sin E_0]$$

Part 14. General Relativity

Part 15. WE Heraeus International Winter School on Gravity and Light

INTRODUCTION (FROM EY)

The International Winter School on Gravity and Light held *central lectures* given by Dr. Frederic P. Schuller. These lectures on General Relativity and Gravity are unequivocally and undeniably, the best and most lucid and well-constructed lecture series on General Relativity and Gravity. The mathematical foundation from topology and differential geometry from which General Relativity arises from is solid, well-selected in rigor. The lectures themselves are well-thought out and clearly explained.

Even more so, the International Winter School provided accompanying Tutorial Sessions for each of the lectures. I had given up hopes in seeing this component of the learning process ever be put online so that anyone and everyone in the world could learn through the Tutorial process as well. I was afraid that nobody would understand how the Tutorial or "Office Hours" session was important for students to digest and comprehend and work out-doing exercises-the material presented in the lectures. This International Winter School gets it and shows how online education has to be done, to do it in an excellent manner, moving forward.

For anyone who is serious about learning General Relativity and Gravity, I would simply point to these video lectures and tutorials.

What I want to do is to build upon the material presented in this International Winter School. Why it's important to me, and to the students and practicing researchers out there, is that the material presented takes the student from an introduction to the research frontier. That is the stated goal of the International Winter School. I want to dig into and help contribute to the cutting edge in research and this entire program with lectures and tutorials appears to be the most direct and sensible route directly to being able to do research in General Relativity and Gravity. -EY 20150323

20200321 update: Someone collected all the lecture videos and tutorials:

<http://mathswithphysics.blogspot.com/2016/07/the-we-heraeus-international-winter.html>

17. LECTURE 1: TOPOLOGY

17.1. Lecture 1: Topological Spaces.

Definition 49. *Let M be a set.*

A **topology** \mathcal{O} is a subset $\mathcal{O} \subseteq \mathcal{P}(M)$, $\mathcal{P}(M)$ power set of M : set of all subsets of M . satisfying

- (i) $\emptyset \in \mathcal{O}$, $M \in \mathcal{O}$
- (ii) $U \in \mathcal{O}$, $V \in \mathcal{O} \implies U \cap V \in \mathcal{O}$
- (iii) $U_\alpha \in \mathcal{O}$, $\alpha \in \mathcal{A} \implies \left(\bigcup_{\alpha \in \mathcal{A}} U_\alpha \right) \in \mathcal{O}$

\mathcal{O} utterly useless

Definition 50. $\mathcal{O}_{\text{standard}} \subseteq \mathcal{P}(\mathbb{R}^d)$

EY : 20150524

I'll fill in the proof that $\mathcal{O}_{\text{standard}}$ is a topology.

Proof. $\emptyset \in \mathcal{O}_{\text{standard}}$
since $\forall p \in \emptyset$, $\exists r \in \mathbb{R}^+$: $\mathcal{B}_r(p) \subseteq \emptyset$ (i.e. satisfied "vacuously")

Suppose $U, V \in \mathcal{O}_{\text{standard}}$.

Let $p \in U \cap V$. Then $\exists r_1, r_2 \in \mathbb{R}^+$ s.t. $\mathcal{B}_{r_1}(p) \subseteq U$
 $\mathcal{B}_{r_2}(p) \subseteq V$

Let $r = \min \{r_1, r_2\}$.
Clearly $\mathcal{B}_r(p) \subseteq U$ and $\mathcal{B}_r(p) \subseteq V$. Then $\mathcal{B}_r(p) \subseteq U \cap V$. So $U \cap V \in \mathcal{O}_{\text{standard}}$.

Suppose, $U_\alpha \in \mathcal{O}_{\text{standard}}, \forall \alpha \in \mathcal{A}$.
Let $p \in \bigcup_{\alpha \in \mathcal{A}} U_\alpha$. Then $p \in U_\alpha$ for at least 1 $\alpha \in \mathcal{A}$.
 $\exists r_\alpha \in \mathbb{R}^+$ s.t. $\mathcal{B}_{r_\alpha}(p) \subseteq U_\alpha \subseteq \bigcup_{\alpha \in \mathcal{A}} U_\alpha$. So $\bigcup_{\alpha \in \mathcal{A}} U_\alpha \in \mathcal{O}_{\text{standard}}$

17.2. **2. Continuous maps.**

17.3. **3. Composition of continuous maps.**

17.4. **4. Inheriting a topology.** EY : 20150524
I'll fill in the proof that given f continuous (cont.), then the restriction of f onto a subspace S is cont. If you want a reference, check out Klaus Jänich [?, pp. 13, Ch. 1 Fundamental Concepts, Sec. Continuous Maps]
If cont. $f : M \rightarrow N, S \subseteq M$, then $f|_S$ cont.

Proof. Let open $V \subseteq N$, i.e. $V \in \mathcal{O}_N$ i.e. V in the topology \mathcal{O}_N of N .

$$f|_S^{-1}(V) = \{m \in M \mid f|_S(m) \in V\}$$

Now $f^{-1}(V) = \{m \in M \mid f(m) \in V\}$.
So $f^{-1}(V) \cap S = f|_S^{-1}(V)$
Now f cont. So $f^{-1}(V) \in \mathcal{O}_N$.
and recall $\mathcal{O}_S := \{U \cap S \mid U \in \mathcal{O}_M\}$.
so $f^{-1}(V) \cap S = f|_S^{-1}(V) \in \mathcal{O}_S$ i.e. $f|_S^{-1}(V)$ open.
 $\implies f|_S$ cont.

TOPOLOGY TUTORIAL SHEET

filename : **main.pdf**
The WE-Heraeus International Winter School on Gravity and Light: Topology

EY : 20150524
What I won't do here is retype up the solutions presented in the Tutorial (cf. https://youtu.be/_XkhZQ-hNLs): the presenter did a very good job. If someone wants to type up the solutions and copy and paste it onto this LaTeX file, in the spirit of open-source collaboration, I would encourage this effort.

Instead, what I want to encourage is the use of as much CAS (Computer Algebra System) and symbolic and numerical computation because, first, we're in the 21st century, second, to set the stage for further applications in research. I use Python and Sage Math alot, mostly because they are open-source software (OSS) and fun to use. Also note that the structure of Sage Math modules matches closely to Category Theory.

In checking whether a set is a topology, I found it strange that there wasn't already a function in Sage Math to check each of the axioms. So I wrote my own; see my code snippet, which you can copy, paste, edit freely in the spirit of OSS here, titled **topology.sage**:

[gist github ernestyalumni topology.sage](#)
[Download topology.sage](#)

Loading **topology.sage**, after changing into (with the usual Linux terminal commands, cd, ls) by

```
sage: load('‘topology.sage’')
```

Exercise 2: Topologies on a simple set.

Question Does $\mathcal{O}_1 := \dots$ constitute a topology ...?.

Solution: Yes, since we check by typing in the following commands in Sage Math:

```
emptyset in O_1
Axiom2check(O_1) # True
Axiom3check(O_1) # True
```

□

Question What about $\mathcal{O}_2 \dots$?

Solution: No since the 3rd. axiom fails, as can be checked by typing in the following commands in Sage Math:

```
emptyset in O_2
Axiom2check(O_2) # True
Axiom3check(O_2) # False
```

18. LECTURE 2: TOPOLOGICAL MANIFOLDS

Lecture 2: Manifolds. Topological spaces: \exists so man that mathematicians cannot even classify them.
For spacetime physics, we may focus on topological spaces (M, \mathcal{O}) that can be charted, analogously to how the surface of the earth is charted in an atlas.

18.1. **Topological manifolds.**

Definition 51. A topological space (M, \mathcal{O}) is called a ***d-dimensional topological method*** if
 $\forall p \in M : \exists U \in \mathcal{O}, U \ni p : \exists x : U \subseteq M \rightarrow x(U) \subseteq \mathbb{R}^d \quad (M, \mathcal{O}), (\mathbb{R}^d, \mathcal{O}_{std})$

(i) x ***invertible*** :

$$x^{-1} : x(U) \rightarrow U$$

- (ii) x ***continuous***
- (iii) x^{-1} ***continuous***

□

18.2. **Terminology.**

18.3. **3. Chart transition maps.** Imagine 2 charts (U, x) and (V, y) with overlapping regions.

18.4. **4. Manifold philosophy.** Often it is desirable (or indeed the way) to define properties (“continuity”) of real-world object (“ $\mathbb{R} \xrightarrow{\gamma} M$ ”) by judging suitable coordinates not on the “real-world” object itself, but on a chart-representation of that real world object.

EY’s add-ons. This lecture gives me a good excuse to review Topology and Topological Manifolds from a mathematician’s point of view. I find John M. Lee’s **Introduction to Topological Manifolds** book good because it’s elementary and thorough and it’s fairly recent (2010) so it’s up to date [?]. See my notes and solutions for the book; it’s a file titled LeeJM_IntroTopManifolds_sol.pdf of which I’ll try to keep the pdf and LaTeX file available for download on my ernestyalumni Google Drive (so try to search for it on Google).

$$\begin{aligned} y_+ : V_+ &\rightarrow (-1, 1) \subseteq \mathbb{R} \\ (m, n) &\mapsto n \\ y_- : V_- &\rightarrow (-1, 1) \subseteq \mathbb{R} \\ (m, n) &\mapsto n \end{aligned}$$

filename: Sheet_1.2.pdf

Exercise 4: Before the invention of the wheel.

Another one-dimensional topological manifold. Another one?

Consider set $F^1 := \{(m, n) \in \mathbb{R}^2 \mid m^4 + n^4 = 1\}$, equipped with subset topology $\mathcal{O}_{\text{std}}|_{F^1}$

Question $x : F^1 \rightarrow \mathbb{R}$ is what?.

Solution . EY : 20150525 The tutorial video https://youtu.be/ghfEQ3u_B6g is really good and this solution is how I'd write it, but it's really the same (I needed the practice).

$$\begin{aligned} x : F^1 &\rightarrow \mathbb{R} \\ (m, n) &\mapsto m \end{aligned}$$

If $m = 0$, $n^4 = 1$ so $n = \pm 1$ so it's not injective.

Let the closed n -dim. upper half-space $\mathbb{H}^n \subseteq \mathbb{R}^1$. Then

$$\begin{aligned} \mathbb{H}^n &= \{(x_1 \dots x_n) \in \mathbb{R}^n \mid x_n \geq 0\} \\ \text{int}\mathbb{H}^n &= \{(x_1 \dots x_n) \in \mathbb{R}^n \mid x_n > 0\} \\ -\mathbb{H}^n &= \{(x_1 \dots x_n) \in \mathbb{R}^n \mid x_n \leq 0\} \\ -\text{int}\mathbb{H}^n &= \{(x_1 \dots x_n) \in \mathbb{R}^n \mid x_n < 0\} \end{aligned}$$

Question This map x may be made injective by restricting its domain to either of 2 maximal open subsets of F^1 . Which ones?.

Solution .

Let

$$\begin{aligned} U_+ &= F^1 \cap \text{int}\mathbb{H}^2 \\ U_- &= F^1 \cap -\text{int}\mathbb{H}^2 \end{aligned}$$

Look at

$$\begin{aligned} x^4 &= 1 - n^4 \\ \implies x &= \pm(1 - n^4)^{1/4} \end{aligned}$$

Then for

$$\begin{aligned} x_+^{-1} : (-1, 1) &\subseteq \mathbb{R} \rightarrow U_+ \\ m &\mapsto (m, (1 - m^4)^{1/4}) \\ x_-^{-1} : (-1, 1) &\subseteq \mathbb{R} \rightarrow U_- \\ m &\mapsto (m, -(1 - m^4)^{1/4}) \end{aligned}$$

x_+, x_- injective (since left inverse exists).

Question Construct injective y .

Solution .

Let

$$\begin{aligned} V_+ &= F^1 \cap \text{int}\mathbb{H}^1 \\ V_- &= F^1 \cap -\text{int}\mathbb{H}^1 \end{aligned}$$

Question Construct inverse y^{-1} . Solution .

For

$$\begin{aligned} y_+^{-1} : (-1, 1) &\subseteq \mathbb{R} \rightarrow V_+ \\ n &\mapsto ((1 - n^4)^{1/4}, n) \\ y_-^{-1} : (-1, 1) &\subseteq \mathbb{R} \rightarrow V_- \\ n &\mapsto (-(1 - n^4)^{1/4}, n) \end{aligned}$$

y_+, y_- injective (since left inverse exists).

Note $(-1, 0) \notin U_+, U_-$

$(1, 0) \notin U_+, U_-$

and

$(0, 1) \notin V_+, V_-$

$(0, -1) \notin V_+, V_-$

Question construct *transition map* $x \circ y^{-1}$.

Solution .

$$\begin{aligned} x_+ y_+^{-1} : (0, 1) &\subseteq \mathbb{R} \rightarrow (0, 1) \subseteq \mathbb{R} \\ n &\mapsto (1 - n^4)^{1/4} \\ x_- y_+^{-1} : (-1, 0) &\subseteq \mathbb{R} \rightarrow (0, 1) \subseteq \mathbb{R} \\ n &\xrightarrow{y_+^{-1}} ((1 - n^4)^{1/4}, n) \xrightarrow{x_-} (1 - n^4)^{1/4} \\ x_+ y_-^{-1} : (0, 1) &\subseteq \mathbb{R} \rightarrow (-1, 0) \subseteq \mathbb{R} \\ n &\mapsto -(1 - n^4)^{1/4} \\ x_- y_-^{-1} : (-1, 0) &\subseteq \mathbb{R} \rightarrow (-1, 0) \subseteq \mathbb{R} \\ n &\mapsto -(1 - n^4)^{1/4} \end{aligned}$$

Question ... Does the collection of these domains and maps form an atlas of F^1 ?.

Yes, with atlas

$$\mathcal{A} = \left\{ \begin{pmatrix} U_+, x_+ \\ U_-, x_- \end{pmatrix}, \begin{pmatrix} V_+, y_+ \\ V_-, y_- \end{pmatrix} \right\}$$

Clearly

$$\begin{aligned} U_+ \cup U_- \cup V_+ \cup V_- &= (F^1 \cap \text{int}\mathbb{H}^2) \cup (F^1 \cap -\text{int}\mathbb{H}^2) \cup (F^1 \cap \text{int}\mathbb{H}^1) \cup (F^1 \cap -\text{int}\mathbb{H}^1) = \\ &= F^1 \cap \mathbb{R}^2 \setminus \{(0,0)\} = F^1 \end{aligned}$$

and (the point is that) x_{\pm}, y_{\pm} are homeomorphisms of open sets of F^1 onto open sets of 1 dim. \mathbb{R}^1 (namely $(-1,1) \subseteq \mathbb{R}^1$), and so \mathcal{A} is an atlas of F^1 .

19. LECTURE 3: MULTILINEAR ALGEBRA

Lecture 3: Multilinear Algebra (International Winter School on Gravity and Light 2015)

We will **not** equip space(time) with a vector space structure. Do you know where

$$\begin{array}{ccc} 5 \cdot \text{Paris} & =? & \\ \text{lie ?} & & \text{Paris} + \text{Vienna} =? \end{array}$$

Moreover, the tangent spaces T_pM (lecture 5) smooth manifolds (Lecture 4)

Beneficial to first study vector spaces abstractly for two reason

- (i) for construction of T_pM one needs an intermediate vector space $C^\infty(M)$
- (ii) tensor technique are most easily understood in an abstract setting.

19.1. Vector spaces.

Definition 52. A vector space $(V, +, -)$ is

- (i) a set V
- (ii) $+: V \times V \rightarrow V$ “addition”
- (iii) $\cdot: \mathbb{R} \times V \rightarrow V$ “s-multiplication” EY : 20160317 s for “scalar”

satisfying:

CANIADDU

$$\begin{array}{ll} C^+ : & v + w = w + v \\ A^+ : & (u + v) + w = u + (v + w) \\ N^+ : & \exists 0 \in V : \forall v \in V : v + 0 = v \\ I^+ : & \forall v \in V : \exists (-v) \in V : v + (-v) = 0 \\ A : & \lambda \cdot (\mu + v) = (\lambda \cdot \mu) \cdot v \quad (\forall \lambda, \mu \in \mathbb{R}) \\ D : & (\lambda + \mu) \cdot v = \lambda \cdot v + \mu \cdot v \\ D : & \lambda \cdot v + \lambda \cdot w = \lambda \cdot (v + w) \\ U : & 1 \cdot v = v \end{array}$$

Terminology. An element of a vector space is often referred to, informally as a vector.

Example. def. **set** of polynomials (fixed) degree $\mathcal{P} := \{p : (-1, +1) \rightarrow \mathbb{R} | p(x) = \sum_{n=0}^N p_n \cdot x^n\}$

Thought bubble: is \square a vector?

$$\square(x) = x^2$$

No $\square \in \mathcal{P}$.

$$+: \mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$$

$$(p, q) \mapsto p + q$$

where $(p + q)(x) = p(x) +_{\mathbb{R}} q(x)$

$$\cdot: \mathbb{R} \times \mathcal{P} \rightarrow \mathcal{P}$$

$$(\lambda, p) \mapsto \lambda \cdot p$$

where $(\lambda \cdot p)(x) := \lambda \cdot_{\mathbb{R}} p(x)$

Thought bubble: \square a vector?

Yes, but who cares?

$(\mathcal{P}, +, \cdot)$ is a vector space.

$$\square \in \mathcal{P}$$

19.2. Linear maps. These are the structure-respecting maps between vector spaces.

EY : 20160316 out of tradition, they’re called “linear” maps

Definition 53. $(V, +_V, \cdot_V)$ and $(W, +_W, \cdot_W)$ vector spaces

Then a map

$$\varphi: V \rightarrow W$$

is called **linear** if

- (i) $\varphi(v +_V \tilde{v}) = \varphi(v) +_W \varphi(\tilde{v})$
- (ii) $\varphi(\lambda \cdot_V v) = \lambda \cdot_W \varphi(v)$

$$\delta: \mathcal{P} \rightarrow \mathcal{P}$$

Example. :

$$p \mapsto \delta(p) := p'$$

linear:

- (i) $\delta(p + q) = (p +_{\mathcal{P}} q)'$ $\underbrace{\quad}_{\text{sum rule}} p' +_{\mathcal{P}} q' = \delta(p) +_{\mathcal{P}} \delta(q)$
- (ii) $\delta(\lambda p) = (\lambda p)' = \lambda \cdot p' = \lambda \cdot \delta(p)$

Notation: $\varphi: V \rightarrow W$ linear $\iff \varphi: V \xrightarrow{\sim} W$

$$\begin{array}{ccccc} V & \xrightarrow[\sim]{\psi} & W & \xrightarrow[\sim]{\varphi} & U \\ & & \searrow \scriptstyle \sim & \nearrow \scriptstyle \sim & \\ & & \varphi \circ \psi & & \end{array}$$

19.2.1. Example*. $\delta \circ \delta: \mathcal{P} \xrightarrow{\sim} \mathcal{P}$

19.3. Vector space of Homomorphisms. fun fact: $(V, +, \cdot)$ $(W, +, \cdot)$ vector spaces

def. $\text{Hom}(V, W) := \{\varphi: V \xrightarrow{\sim} W\}$ set.

We can make this into a vector spaces.

$$\begin{array}{l} \oplus: \text{Hom}(V, W) \times \text{Hom}(V, W) \rightarrow \text{Hom}(V, W) \\ (\varphi, \psi) \mapsto \varphi \oplus \psi \end{array}$$

where $(\varphi \otimes \psi)(v) := \varphi(v) +_W \psi(v)$

$\otimes: \dots$ similarly.

$(\text{Hom}(V, W), \oplus, \otimes)$ **is** a vector space.

19.3.1. Example*. $\text{Hom}(\mathcal{P}, \mathcal{P})$ is a vector space.

$$\delta \in \text{Hom}(\mathcal{P}, \mathcal{P})$$

$$\delta \circ \delta \in \text{Hom}(\mathcal{P}, \mathcal{P})$$

\vdots

$$\underbrace{\delta \circ \dots \circ \delta}_M \in \text{Hom}(\mathcal{P}, \mathcal{P})$$

$$\implies 5 \circ \delta \oplus_{\text{Hom}(\mathcal{P}, \mathcal{P})} \delta \circ \delta \in \text{Hom}(\mathcal{P}, \mathcal{P})$$

19.4. **Dual vector space.** heavily used special case:

$(V, +, \cdot)$ vector space:

Definition 54.

$$V^* := \{\varphi : V \xrightarrow{\sim} \mathbb{R}\} = Hom(V, \mathbb{R})$$

$\underbrace{(V^*, \oplus, \otimes)}_{\text{dual vector space (to } V)}$ is a vector space

Terminology: $\varphi \in V^*$ is called, informally, a covector.

Example. $I : \mathcal{P} \xrightarrow{\sim} \mathbb{R}$

i.e. $I \in \mathcal{P}^*$

def. $I(p) := \int_0^1 dx p(x)$

linear:
$$I(p+q) = \int_0^1 dx \underbrace{(p+q)(x)}_{p(x)+q(x)} \\ = \dots = I(q) + I(p)$$

$$I(\lambda p) = \lambda \cdot I(p)$$

i.e. $I = \int_0^1 dx$

19.5. **Tensors.**

Definition 55. Let $(V, +, \cdot)$ be a vector space.

An $\underline{(r, s)}$ -tensor T over V $r, s \in \mathbb{N}_0$

is a multi-linera map

$$T : \underbrace{V^* \times \dots \times V^*}_r \times \underbrace{V \times \dots \times V}_s \xrightarrow[\sim]{\sim} \mathbb{R}$$

19.5.1. *Example.* : T (1,1)-tensor

$$T(\varphi + \psi, v) = T(\varphi, v) + T(\psi, v)$$

$$T(\varphi, v + w) = T(\varphi, v) + T(\varphi, w)$$

$$T(\lambda \varphi, v) = \lambda \cdot T(\varphi, v)$$

$$T(\varphi, \lambda \cdot v) = \lambda T(\varphi, v)$$

$$T(\varphi + \psi, v + w) = \\ = t(\varphi, v) + T(\varphi, w) + T(\psi, v) + t(\psi, w)$$

Excursion: Given $T : V^* \times V \xrightarrow{\sim} \mathbb{R}$

$$\phi_T : V \xrightarrow{\sim} (V^*)^* \underset{\dim V < \infty}{=} V$$

Define $v \mapsto \underbrace{T(\cdot, v)}_{V^* \xrightarrow{\sim} \mathbb{R}}$

Given $\phi : V \xrightarrow{\sim} V$

Construct $T_\phi : V^* \times V \xrightarrow{\sim} \mathbb{R}$

$$(\varphi, v) \mapsto \varphi(\phi(v))$$

\implies given $T : T = T_{\varphi_T}$

given $\phi : \phi = \phi_{T_\phi}$

Example. $g : P \times P \xrightarrow{\sim} \mathbb{R}$

$$(p, q) \mapsto \int_{-1}^1 dx p(x) q(x)$$

is a $(0, 2)$ -tensor over P .

Info: If $T \in \text{Hom}(V, W)$

19.6. **Vectors and covectors as tensors.**

Theorem 21. (including proof)

“covector” $\varphi \in V^* \iff \varphi : V \xrightarrow{\sim} \mathbb{R} \iff \varphi(0, 1)$ -tensor.

Theorem 22. $v \in V \underset{\dim V < \infty}{=} (V^*)^* \iff v : V^* \xrightarrow{\sim} \mathbb{R} \iff v$ is $(1, 0)$ -tensor.

19.7. **Bases.**

Definition 56. $(V, +, \cdot)$ vector space.

A subset $B \subset V$ is called

a basis if

Thought bubble: Hamel (L.A.) EY : 20160316 Hamel basis, Linear Algebra

$$\forall v \in V \quad \exists \text{ finite } \underbrace{F}_{\{f_1, \dots, f_n\}} \subset B : \exists ! \underbrace{v^1, v^2, \dots, v^n}_{\in \mathbb{R}}, \quad v = v^1 f_1 + \dots + v^n f_n$$

Definition 57. If \exists basis \mathcal{B} with finitely many elements, say d many, then we call $d =: \dim V$

This is well-defined.

Remark: $(V, +, \cdot)$ be a finite-dim. vector space.

Having chosen a basis e_1, \dots, e_n of $(V, +, \cdot)$ we may uniquely associate

(Thought bubble: this requires a chosen basis)

$v \mapsto (v^1, \dots, v^n)$ called the components of v w.r.t. chosen basis

where: $v^1 e_1 + \dots + v^n e_n = v$

19.8. **Basis for the dual space.** choose Basis e_1, \dots, e_n for V

can choose Basis $\epsilon^1, \dots, \epsilon^n$ for V^*

However, more economical to require

once e_1, \dots, e_n on V has been chosen, that

$$\epsilon^a(e_b) = \delta_b^a$$

This uniquely determines choice of $\epsilon^1, \dots, \epsilon^n$ from choice of e_1, \dots, e_n

Definition 58. If a basis $\epsilon^1, \dots, \epsilon^n$ of V^* satisfies this, it is called the **dual basis** (of the dual space)

Example: P ($N = 3$)

$$e_0(x) = 1$$

$$e_1(x) = x$$

e_0, e_1, e_2, e_3 basis if $e_2(x) = x^2 \{e_a(x) := x^a$

$$e_3(x) = x^3$$

$$\epsilon^0, \epsilon^1, \epsilon^2, \epsilon^3 \text{ dual basis } \epsilon^a := \frac{1}{a!} \partial^a|_{x=0}$$

Proof. $\epsilon^a(e_b) = \delta_b^a$

□

19.9. Components of tensors. Let T be an (r, s) -tensor on a finite-dim. vs. V . Then define the $(r + s)^{\dim V}$ many real numbers.

$$\underbrace{T^{i_1 \dots i_r}_{j_1 \dots j_s}}_{\in \mathbb{R}} := T(\epsilon^{i_1}, \dots, \epsilon^{i_r}, e_{j_1}, e_{j_2}, \dots, e_{j_s})$$

$i_1 \dots i_r, j_1 \dots j_s \in \{1, \dots, \dim V\}$
Thought bubble: $\underbrace{T^{i_1 \dots i_r}_{j_1 \dots j_s}}_{\in \mathbb{R}}$ are the components of the tensor w.r.t. chosen basis

Useful: Knowing components (and basis) one can reconstruct the entire tensor.

Example. $T(1, 1)$ - tensor

$$T^i_j := T(\epsilon^i, e_j)$$

reconstruct

$$\begin{aligned} T(\varphi, v) &= T\left(\sum_{i=1}^{\dim V} \varphi_i \epsilon^i, \sum_{j=1}^{\dim V} v^j e_j\right) \quad \varphi_i \in \mathbb{R}, v^j \in \mathbb{R} \\ &= \sum_{i=1}^{\dim V} \sum_{j=1}^{\dim V} \varphi_i v^j \underbrace{T(\epsilon^i, e_j)}_{T^i_j} \\ &=: \varphi_i v^j T^i_j \end{aligned}$$


20. LECTURE 4: DIFFERENTIABLE MANIFOLDS

so far: top. mfd. (M, \mathcal{O})

$$\dim M = d$$


- we wish to define a notion of differentiable
- curves $\mathbb{R} \rightarrow M$
- function $M \rightarrow \mathbb{R}$
- maps $M \rightarrow N$

20.1. 1. Strategy. choose a chart (U, x)
 $\gamma : \mathbb{R} \rightarrow M$ portion of curve in chart domain



idea. try to “lift” the undergraduate notion of differentiability of a curve on \mathbb{R}^d to a notion of differentiability of a curve on M

Problem Can this be well-defined under change of chart?



$x \circ \gamma$ undergraduate differentiable (“as a map $\mathbb{R} \rightarrow \mathbb{R}^d$ ”)

$$\underbrace{y \circ \gamma}_{\text{maybe only continuous, but not undergraduate differentiable}} = \underbrace{(y \circ x^{-1})}_{\text{continuous}} \circ \underbrace{(x \circ \gamma)}_{\text{undergrad differentiable}} = y \circ (x^{-1} \circ x) \circ \gamma$$

At first sight, strategy does not work out.

20.2. 2. Compatible charts. In section 1, we used any imaginable charts on the top. mfd. (M, \mathcal{O}) . To emphasize this, we may say that we took U and V from the *maximal atlas* \mathcal{A} of (M, \mathcal{O}) .

Definition 59. Two charts (U, x) and (V, y) of a top. mfd. are called \ast -compatible if either

- (a) $U \cap V = \emptyset$ or
- (b) $U \cap V \neq \emptyset$

chart transition maps have undergraduate \ast property.
EY : 20151109 e.g. since $\mathbb{R}^d \rightarrow \mathbb{R}^d$, can use undergraduate \ast property such as continuity or differentiability.

$$\begin{aligned} y \circ x^{-1} : x(U \cap V) \subseteq \mathbb{R}^d &\rightarrow y(U \cap V) \subseteq \mathbb{R}^d \\ x \circ y^{-1} : y(U \cap V) \subseteq \mathbb{R}^d &\rightarrow x(U \cap V) \subseteq \mathbb{R}^d \end{aligned}$$

Philosophy:

Definition 60. An atlas \mathcal{A}_\ast is a \ast -compatible atlas if any two charts in \mathcal{A}_\ast are \ast -compatible.

Definition 61. A \ast -manifold is a triple $(\underbrace{M, \mathcal{O}}_{\text{top. mfd.}}, \mathcal{A}_\ast)$ $\mathcal{A}_\ast \subseteq \mathcal{A}_{\text{maximal}}$

\ast	undergraduate \ast	
C^0	$C^0(\mathbb{R}^d \rightarrow \mathbb{R}^d) =$	continuous maps w.r.t. \mathcal{O}
C^1	$C^1(\mathbb{R}^d \rightarrow \mathbb{R}^d) =$	differentiable (once) and is continuous
C^k		k -times continuously differentiable
D^k		k -times differentiable
\vdots		
C^∞	$C^\infty(\mathbb{R}^d \rightarrow \mathbb{R}^d)$	
\bigcup		
C^ω	\exists multi-dim. Taylor exp.	
\mathbb{C}^∞	satisfy Cauchy-Riemann equations, pair-wise	


⁵<http://mathoverflow.net/questions/8789/can-every-manifold-be-given-an-analytic-structure>

EY : 20151109 Schuller says: C^k is easy to work with because you can judge k -times cont. differentiability from existence of all partial derivatives **and** their continuity. There are examples of maps that partial derivatives exist but are not D^k , k -times differentiable.

Theorem 23 (Whitney). *Any $C^{k \geq 1}$ -atlas, $\mathcal{A}_{C^{k \geq 1}}$ of a topological manifold contains a C^∞ -atlas.*
Thus we may w.l.o.g. always consider C^∞ -manifolds, “smooth manifolds”, unless we wish to define Taylor expandibility/-complex differentiability ...

EY : 20151109 Hassler Whitney ⁵

Definition 62. A smooth manifold $(\underbrace{M, \mathcal{O}}_{\text{top. mfd.}}, \underbrace{\mathcal{A}}_{C^\infty\text{-atlas}})$



EY: 20151109 Schuller was explaining that the trajectory is real in M ; the coordinate maps to obtain coordinates is $x \circ \gamma$

20.3. 4. Diffeomorphisms. $M \xrightarrow{\phi} N$

If M, N are naked sets, the structure preserving maps are the bijections (invertible maps).
e.g. $\{1, 2, 3\} \rightarrow \{a, b\}$


Definition 63. $M \cong_{\text{set}} N$ (set-theoretically) isomorphic if \exists bijection $\phi : M \rightarrow N$

Examples. $\mathbb{N} \cong_{\text{set}} \mathbb{Z}$
 $\mathbb{N} \cong_{\text{set}} \mathbb{Q}$ (EY: 20151109 Schuller says from diagonal counting)
 $\mathbb{N} \not\cong_{\text{set}} \mathbb{R}$
Now $(M, \mathcal{O}_M) \cong_{\text{top}} (N, \mathcal{O}_N)$ (topl.) isomorphic = “homeomorphic” \exists bijection $\phi : M \rightarrow N$
 ϕ, ϕ^{-1} are continuous.
 $(V, +, \cdot) \cong_{\text{vec}} (W, +_w, \cdot_w)$ (EY: 20151109 vector space isomorphism) if
 \exists bijection $\phi : V \rightarrow W$ linearly
finally

Definition 64. Two C^∞ -manifolds $(M, \mathcal{O}_M, \mathcal{A}_M)$ and $(N, \mathcal{O}_N, \mathcal{A}_N)$ are said to be **diffeomorphic** if \exists bijection $\phi : M \rightarrow N$ s.t.

$$\begin{aligned}\phi &: M \rightarrow N \\ \phi^{-1} &: N \rightarrow M\end{aligned}$$

are both C^∞ -maps



Theorem 24. $\#$ = number of C^∞ -manifolds one can make out of a given C^0 -manifolds (if any) - up to diffeomorphisms.

$dim M$	$\#$	
1	1	Morse-Radon theorems
2	1	Morse-Radon theorems
3	1	Morse-Radon theorems
4	uncountably infinitely many	
5	finite	surgery theory
6	finite	surgery theory
\vdots	finite	surgery theory

EY : 20151109 cf. <http://math.stackexchange.com/questions/833766/closed-4-manifolds-with-uncountably-many-diffeomorphisms>
The wild world of 4-manifolds

TUTORIAL 4 DIFFERENTIABLE MANIFOLDS

EY : 20151109 The gravity-and-light.org website, where you can download the tutorial sheets *and* the full length videos for the tutorials and lectures, are no longer there. = (
Hopefully, the YouTube video will remain: https://youtu.be/FXPdKxOq1KA?list=PLFeEvEPtX_ORQ1ys-7VIsKlBWz7RX-FaL

Exercise 1: True or false?. These basic questions are designed to spark discussion and as a self-test.

Tick the correct statements, but not the incorrect ones!

- (a) The function $f : \mathbb{R} \rightarrow \mathbb{R}$, ...
- -
 - ... , defined by $f(x) = |x^3|$, lies in $C^3(\mathbb{R} \rightarrow \mathbb{R})$.

EY : 20151109 **Solution 1a3.** For $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = |x^3| = \begin{cases} x^3 & \text{if } x \geq 0 \\ -x^3 & \text{if } x < 0 \end{cases}$

$$\begin{aligned}f'(x) &= \begin{cases} 3x^2 & \text{if } x \geq 0 \\ -3x^2 & \text{if } x < 0 \end{cases} \\ f''(x) &= \begin{cases} 6x & \text{if } x \geq 0 \\ -6x & \text{if } x < 0 \end{cases}\end{aligned}$$

Thus,

$f(x) = |x^3| \in C^1(\mathbb{R})$ but $f(x) \notin C^2(\mathbb{R}) \subseteq C^3(\mathbb{R})$

-
-

- (b)
- (c)

Short Exercise 4: Undergraduate multi-dimensional analysis .

A good notation and basic results for partial differentiation.

For a map $f : \mathbb{R}^d \rightarrow \mathbb{R}$ we denote by the map $\partial_i f : \mathbb{R}^d \rightarrow \mathbb{R}$ the partial derivative with respect to the i -th entry.

Question :: Given a function

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}; (\alpha, \beta, \delta) \mapsto f(\alpha, \beta, \delta) := \alpha^3 \beta^2 + \beta^2 \delta + \delta$$

calculate the values of the following derivatives:

Solution ::

- $(\partial_2 f)(x, y, z) =$
- $(\partial_1 f)(\square, \circ, *) =$
- $(\partial_1 \partial_2 f)(a, b, c) =$
- $(\partial_3^2 f)(299, 1222, 0) =$

EY: 20151110

For $f(\alpha, \beta, \delta) := \alpha^3 \beta^2 + \beta^2 \delta + \delta$, or $f(x, y, z) = x^3 y^2 + y^2 z + z$,

$$(\partial_2 f) = 2(x^3 y + yz)$$

$$(\partial_1 f) = 3x^2 y^2$$

$$(\partial_1 \partial_2 f) = 6x^2 y$$

$$(\partial_3^2 f) = 0$$

and so

- $(\partial_2 f)(x, y, z) = 2(x^3 y + yz)$
- $(\partial_1 f)(\square, \circ, *) = 3\square^2 \circ^2$
- $(\partial_1 \partial_2 f)(a, b, c) = 6a^2 b$
- $(\partial_3^2 f)(299, 1222, 0) = 0$

Exercise 5: Differentiability on a manifold.


How to deal with functions and curves in a chart

Let $(M, \mathcal{O}, \mathcal{A})$ be a smooth d -dimensional manifold. Consider a chart (U, x) of the atlas \mathcal{A} together with a smooth curve $\gamma : \mathbb{R} \rightarrow U$ and a smooth function $f : U \rightarrow \mathbb{R}$ on the domain U of the chart.

Question :: Draw a commutative diagram containing the chart domain, chart map, function, curve and the respective represen-

tatives of the function and the curve in the chart.

Solution ::



Question :: Consider, for $d = 2$,

$$(x \circ \gamma)(\lambda) := (\cos(\lambda), \sin(\lambda)) \text{ and } (f \circ x^{-1})((x, y)) := x^2 + y^2$$


Using the chain rule, calculate

$$(f \circ \gamma)'(\lambda)$$

explicitly.

Solution ::

EY : 20151109 Indeed, the domains and codomains of this $f \circ \gamma$ mapping makes sense, from $\mathbb{R} \rightarrow \mathbb{R}$ for



$$(f \circ \gamma)'(\lambda) = (Df) \cdot \dot{\gamma}(\lambda) = \frac{\partial f}{\partial x^j} \dot{\gamma}^j(\lambda) = 2x(-\sin \lambda) + 2y \cos \lambda = 2(-\cos \lambda \sin \lambda + \sin \lambda \cos \lambda) = 0$$

21. LECTURE 5: TANGENT SPACES

lead question: “what is the velocity of a curve γ point p ?

21.1. Velocities.

Definition 65. $(M, \mathcal{O}, \mathcal{A})$ smooth mfd.

curve $\gamma : \mathbb{R} \rightarrow M$ at least C^1 .

Suppose $\gamma(\lambda_0) = p$

The **velocity** of γ p is the linear map


$$v_{\gamma,p} : C^\infty(M) \xrightarrow{\sim} \mathbb{R}$$

$C^\infty(M) := \{f : M \rightarrow \mathbb{R} | f \text{ smooth function} \}$ equipped with $(f \oplus g)(p) := f(p) + g(p)$

$$(\lambda \otimes g)(p) := \lambda \cdot g(p)$$

\sim denotes linear map on top of \rightarrow .

$$f \mapsto v_{\gamma,p}(f) := (f \circ \gamma)'(\lambda_0)$$



intuition

Schuller says: children run around the world. Temperature function as temperature contour lines. You feel the temperature.

You observe the rate of change of temperature as you run around. f is temperature.

past: “ $\underbrace{v^i}_{\text{vector}}(\partial_i f) = (\underbrace{v^i \partial_i}_{\text{vector}})f$

21.2. Tangent vector space.

Definition 66. For each point $p \in M$
def the **set** “tangent space $\neq_0 M$ p “

$v_{\gamma,p} \oplus v_{\delta,p} = v_{\sigma,p}$

$$T_pM := \{v_{\gamma,p} | \gamma \text{ smooth curves} \}$$

□

picture:
rather \overline{M} than (embedded) $p \in T_pM$ EY : 20151109 see https://youtu.be/pepU_7NJSgM?t=12m38s for the picture
Observation: T_pM can be made into a vector space.

$$\begin{aligned} \oplus : T_pM \times T_pM &\rightarrow \\ (v_{\gamma,p} \oplus v_{\delta,p}) \big(\underbrace{f}_{\in C^\infty(M)} \big) &:= v_{\gamma,p}(f) +_{\mathbb{R}} v_{\delta,p}(f) \\ \odot : \mathbb{R} \times T_pM &\rightarrow \text{Hom}(C^\infty(\mathbb{R}), \mathbb{R}) \\ (\alpha \odot v_{\gamma,p})(f) &:= \alpha \cdot_{\mathbb{R}} v_{\gamma,p}(f) \end{aligned}$$

Remains to be shown that

- (i) $\exists \sigma$ curve : $v_{\gamma,p} \oplus v_{\delta,p} = v_{\sigma,p}$
- (ii) $\exists \tau$ curve : $\alpha \odot v_{\gamma,p} = v_{\tau,p}$

Claim: $\tau : \mathbb{R} \rightarrow M$ where $\mu_\alpha : \mathbb{R} \rightarrow \mathbb{R}$, does the trick.
 $\mapsto \tau(\lambda) := \gamma(\alpha\lambda + \lambda_0) = (\gamma \circ \mu_\alpha)(\lambda)$ $r \mapsto \alpha \cdot r + \lambda_0$
 $\tau(0) = \gamma(\lambda_0) = p$

$$\begin{aligned} v_{\tau,p} &:= (f \circ \tau)'(0) = (f \circ \gamma \circ \mu_\alpha)'(0) \\ &= (f \circ \gamma)'(\lambda_0) \cdot \alpha = \\ &= \alpha \cdot v_{\gamma,p} \end{aligned}$$

Now for the sum:
 $v_{\gamma,p} \oplus v_{\delta,p} \stackrel{?}{=} v_{\sigma,p}$
make a choice of chart $(\underbrace{U}_{\ni p}, x)$ In cloud: ill definition alarm bells.

and define:
Claim:

$$\begin{aligned} \sigma : \mathbb{R} &\rightarrow M \\ \sigma(\lambda) &:= x^{-1} \big(\underbrace{(x \circ \gamma)(\lambda_0 + \lambda) + (x \circ \delta)(\lambda_1 + \lambda) - (x \circ \gamma)(\lambda_0)}_{\mathbb{R} \rightarrow \mathbb{R}^d} \big) \end{aligned}$$


does the trick.

Proof. Since:

$$\begin{aligned} \sigma_x(0) &= x^{-1}((x \circ \gamma)(\lambda_0) + (x \circ \delta)(\lambda_1) - (x \circ \gamma)(\lambda_0)) \\ &= \delta(\lambda_1) = p \end{aligned}$$

Now:

$$\begin{aligned} v_{\sigma_x,p}(f) &:= (f \circ \sigma_x)'(0) = \\ &= \underbrace{((f \circ x^{-1}) \circ (x \circ \sigma_x))'(\gamma)}_{\mathbb{R}^d \rightarrow \mathbb{R}} = \underbrace{(x \circ \sigma_x)'(0)}_{(x \circ \gamma)'(\lambda_0) + (x \circ \delta)'(\lambda_1)} \cdot (\partial_i(f \circ x^{-1}))(x(\underbrace{\sigma(0)}_p)) = \\ &= (x \circ \gamma)'(\lambda_0)(\partial_i(f \circ x^{-1}))(x(p)) + (x \circ \delta)'(\lambda_1)(\partial_i(f \circ x^{-1}))(x(p)) \\ &= (f \circ \gamma)'(\lambda_0) + (f \circ \delta)'(\lambda_1) = \\ &= v_{\gamma,p}(f) + v_{\delta,p}(f) \quad \forall f \in C^\infty(M) \end{aligned}$$



picture: (cf. https://youtu.be/pepU_7NJSgM?t=39m5s)

$$\begin{aligned} \gamma : \mathbb{R} &\rightarrow M \\ \delta : \mathbb{R} &\rightarrow M \end{aligned}$$

$(\gamma \oplus)(\lambda) := \gamma(\lambda) + \delta(\lambda)$
EY : 20151109 Schuller says adding trajectories is chart dependent, bad. Adding velocities is good.

21.3. Components of a vector wrt a chart.

Definition 67. Let $(U, x) \in \mathcal{A}_{smooth}$.
 $\gamma : \mathbb{R} \rightarrow U$
Let $\gamma(0) = p$.
Calculate

$$\begin{aligned} v_{\gamma,p}(f) &:= (f \circ \gamma)'(0) = \underbrace{((f \circ x^{-1}) \circ (x \circ \gamma))'(\gamma)}_{\mathbb{R}^d \rightarrow \mathbb{R}} = \underbrace{(x \circ \gamma)^{i'}(0)}_{\dot{\gamma}_x^i(0)} \cdot \underbrace{(\partial_i(f \circ x^{-1}))(x(p))}_{=:(\frac{\partial f}{\partial x^i})_p} \end{aligned}$$

think cloud $f : M \rightarrow \mathbb{R}$

$$= \boxed{\dot{\gamma}_x^i(0) \cdot \left(\frac{\partial}{\partial x^i} \right)_p} f \quad \forall f \in C^\infty(M)$$

\therefore as a map.

$$v_{\gamma,p} \underbrace{=}_{\text{use of chart}} \underbrace{\gamma_x^i(0)}_{\text{“components of the velocity } v_{\gamma,p} \text{”}} \underbrace{\left(\frac{\partial}{\partial x^i} \right)}_{\text{basis elements of the } T_pM \text{ wrt which the components need to be understood. “chart induced basis of } T_pM \text{”}}$$

Picture: https://youtu.be/pepU_7NJSgM?t=1h16s

21.4. 4. Chart-induced basis.

Definition 68. $(U, x) \in \mathcal{A}_{smooth}$
the $\left(\frac{\partial}{\partial x^1}\right)_p, \dots, \left(\frac{\partial}{\partial x^d}\right)_p \in T_p U \subseteq T_p M$
constitute a **basis** of $T_p U$

Proof. remains: linearly independent

$$\begin{aligned} \lambda^i \left(\frac{\partial}{\partial x^i} \right)_p &\stackrel{!}{=} 0 \\ \implies \lambda^i \left(\frac{\partial}{\partial x^i} \right)_p (x^j) &= \lambda^i \partial_i (\underbrace{x^j \circ x^{-1}})(x(p)) = & x^j \circ x^{-1} : \mathbb{R}^d \rightarrow \mathbb{R} \\ & & (\alpha^1, \dots, \alpha^d) \mapsto \alpha^j \\ &= \lambda^i \delta_i^j = \lambda^j & j = 1, \dots, d \end{aligned}$$

in cloud: $x^j : U \rightarrow \mathbb{R}$ differentiable

Corollary 1. $\dim T_p M = d = \dim M$

Terminology: $X \in T_p M \rightarrow \exists \gamma : \mathbb{R} \rightarrow M : X = v_{\gamma, p}$ and
 $\exists \underbrace{X_1^1, \dots, X^d}_\in \mathbb{R} : X = X^i \left(\frac{\partial}{\partial x^i} \right)_p$

21.5. 5. Change of vector components under a change of chart. ✖ vector does **not** change under change of chart.

Let (U, x) and (V, y) be overlapping charts and $p \in U \cap V$.
Let $X \in T_p M$

$$X_{(y)}^i \cdot \left(\frac{\partial}{\partial y^i} \right)_p \underbrace{=}_p \underbrace{X}_{(V, y)} \underbrace{=}_{(U, x)} X_x^i \left(\frac{\partial}{\partial x^i} \right)_p$$

to study change of components formula:

$$\begin{aligned} \left(\frac{\partial}{\partial x^i} \right)_p f &= \partial_i (f \circ x^{-1})(x(p)) = \\ &= \partial_i (\underbrace{(f \circ y^{-1})}_{\mathbb{R}^d \rightarrow \mathbb{R}} \circ \underbrace{(y \circ x^{-1})}_{\mathbb{R}^d \rightarrow \mathbb{R}^d})(x(p)) \\ &= (\partial_i (y^i \circ x^{-1}))(x(p)) \cdot (\partial_j (f \circ y^{-1}))(y(p)) = \\ &= \boxed{\left(\frac{\partial y^p}{\partial x^i} \right)_p \cdot \left(\frac{\partial f}{\partial y^j} \right)_p} f \\ \implies X_{(x)}^i \left(\frac{\partial y^j}{\partial x^i} \right)_p \left(\frac{\partial}{\partial y^j} \right)_p &= X_{(y)}^j \left(\frac{\partial}{\partial y^j} \right)_p \\ \implies \boxed{X_{(y)}^j = \left(\frac{\partial y^j}{\partial x^i} \right)_p X_{(x)}^i} \end{aligned}$$

21.6. 6. Cotangent spaces. $T_p M = V$

trivial $(T_p M)^* := \{\varphi : T_p M \xrightarrow{\sim} \mathbb{R}\}$
Example: $f \in C^\infty(M)$

$$\begin{aligned} (df)_p : T_p M &\xrightarrow{\sim} \mathbb{R} \\ X &\mapsto (df)_p(X) \end{aligned}$$

i.e. $\boxed{(df)_p \in T_p M^*}$

$(df)_p$ called the gradient of $f \quad p \in M$.
Calculate components of gradient w.r.t. chart-induced basis (U, x)

$$\begin{aligned} ((df)_p)_j &:= (df)_p \left(\left(\frac{\partial}{\partial x^j} \right)_p \right) \\ &= \left(\frac{\partial f}{\partial x^j} \right)_p = \partial_j (f \circ x^{-1})(x(p)) \end{aligned}$$

Theorem 25. Consider chart $(U, x) \implies x^i : U \rightarrow \mathbb{R}$

□ Claim: $(dx^1)_p, (dx^2)_p, \dots, (dx^d)_p$ basis of $T_p^* M$
 \implies In fact: dual basis:

$$(dx^a)_p \left(\left(\frac{\partial}{\partial x^b} \right)_p \right) = \left(\frac{\partial x^a}{\partial x^b} \right)_p = \dots = \delta_b^a$$

21.7. 7. Change of components of a covector under a change of chart:

$$\underbrace{T_p^* M}_{\ni \omega} \text{ with } \omega_{(y)}(dy^j)_p = \omega = \omega_{(x)i}(dx^i)_p$$

$$\implies \boxed{\omega_{(y)i} = \frac{\partial x^j}{\partial y^i} \omega_{(x)j}}$$

LECTURE 6: FIELDS

cf. **Lecture 6: Fields (International Winter School on Gravity and Light 2015)**

So far:

$$\begin{array}{c} T_p M \\ \vdots \downarrow \\ T_p^* M \\ \vdots \downarrow \\ \vdots \end{array},$$

now
in Thought Cloud: theory of bundles

21.8. Bundles.

Definition 69. A **bundle** is a triple

$$E \xrightarrow{\pi} M$$

E smooth manifold “**total space**”
 π smooth map (surjective) “*projection map*”
 M smooth manifold “*base space*”

Example $E = \text{cylinder}$ $M = \text{circle}$

Definition 70.

$$E \overset{T}{\longrightarrow} M$$

bundle.

$$\begin{array}{l} p \in M \\ \text{define } \textbf{fibre over } p \\ \qquad := \text{preim}_\pi(\{p\}) \end{array}$$

Definition 71. A *section* σ of a bundle

$$\begin{array}{c} E \\ \pi \downarrow \nearrow \sigma \\ M \end{array}$$

require $\pi \circ \sigma = id_M$

Schuller says: in quantum mechanics, Aside: $\psi : M \rightarrow \mathbb{C}$

21.9. Tangent bundle of smooth manifold. $(M, \mathcal{O}, \mathcal{A})$ smooth manifold

(a) as a **set** $TM := \dot{\bigcup}_{p \in M} T_p M$

(b) surjective $\pi : TM \rightarrow M$ the *unique* point $p \in M, X \in T_p M$

$$\begin{array}{ccccc} X \mapsto p & & & & \\ \text{situation: } \underbrace{TM}_{\text{set}} & \xrightarrow{\pi} & \underbrace{M}_{\text{smooth manifold}} & & \end{array}$$

surjective map

(c) Construct topology on TM that is the coarsest topology such that π (just) continuous. (“initial topology with respect to π ”).

$$\mathcal{O}_{TM} := \{\text{preim}_\pi(U) | U \in \mathcal{O}\}$$

Show: Tutorial \mathcal{O}_{TM} Schuller says this is shown in the tutorial (TM, \mathcal{O}_{TM})

Construction of a C^∞ -atlas on TM from the C^∞ -atlas \mathcal{A} on M .

$$\mathcal{A}_{TM} := \{(TU, \xi_x) | (U, x) \in \mathcal{A}\}$$

where

$$\begin{array}{l} \xi_x : TU \rightarrow \mathbb{R}^{2 \cdot \dim M} \\ X \mapsto \underbrace{((x^1 \circ \pi)(X), \dots, (x^d \circ \pi)(X), (dx^1)_{\pi(X)}(X), \dots, (dx^d)_{\pi(X)}(X))}_{(U,x) - \text{ coords of } \pi(X) \text{ (d many) }} \end{array}$$

where $X \in T_{\pi(X)} M$
 $X = X^i_{(x)} \left(\frac{\partial}{\partial x^i} \right)_{\pi(X)}$

$$\begin{aligned} (dx^j)_{\pi(X)}(X) &= (dx^j)_{\pi(X)} \left(X^i_{(x)} \left(\frac{\partial}{\partial x^i} \right)_{\pi(X)} \right) = \\ &= X^i_{(x)} \delta_i^j = X^j_{(x)} \end{aligned}$$

Write $\xi_x^{-1} : \xi_x(TU) \subseteq \mathbb{R}^{2 \dim M} \rightarrow TU$

$$(\alpha^1, \dots, \alpha^d, \beta^1, \dots, \beta^d) := \beta^i \left(\frac{\partial}{\partial x^i} \right) \underbrace{x^{-1}(\alpha^1, \dots, \alpha^d)}_{\pi(X)}$$

Check:

$$\begin{aligned} (\xi_y \circ \xi_x^{-1})(\alpha^1, \dots, \alpha^d, \beta^1, \dots, \beta^d) &= \\ &= \xi_y \left(\beta^i \left(\frac{\partial}{\partial x^i} \right)_{x^{-1}(\alpha^1, \dots, \alpha^d)} \right) \\ &= \left(\dots, (y^i \circ \pi)(\beta^m \cdot \left(\frac{\partial}{\partial x^m} \right)_{x^{-1}(\alpha^1 \dots \alpha^d)}), \dots, \dots (dy^i)_{x^{-1}(\alpha^1, \dots, \alpha^d)} \left(\beta^m \left(\frac{\partial}{\partial x^m} \right)_{x^{-1}(\alpha^1 \dots \alpha^d)} \right), \dots \right) = \\ &= \underbrace{(\dots, (y^i \circ x^{-1})(\alpha^1, \dots, \alpha^d), \dots, \dots, \beta^m (dy^i)_{x^{-1}(\alpha^1, \dots, \alpha^d)} \left(\left(\frac{\partial}{\partial x^m} \right)_{x^{-1}(\alpha^1 \dots \alpha^d)} \right))}_{\beta^m \left(\frac{\partial y}{\partial x^m} \right)_{x^{-1}(\alpha^1 \dots \alpha^d)}} \end{aligned}$$

Check transition map: $(U, x), (V, y), \quad U \cap V \neq \emptyset$
 $\left(\frac{\partial y}{\partial x^m} \right)_{x^{-1}(\alpha^1 \dots \alpha^d)} = \partial_m (y^i \circ x^{-1})(x \circ (x^{-1}(\alpha^1 \dots \alpha^d))) = \partial_m (y^i \circ x^{-1})(\alpha^1 \dots \alpha^d)$ smooth.
upshot

$$\underbrace{TM}_{\text{smooth manifold}} \xrightarrow{\pi} \underbrace{M}_{\text{smooth manifold}}$$

smooth map

bundle, called the tangent bundle.

3. Vector fields.

Definition 72. A *smooth vector field* χ is a smooth map, (where)

$$\begin{array}{c} TM \\ \pi \downarrow \nearrow \chi \\ M \end{array}$$

Example:

$$\begin{array}{c} TS^1 \\ \downarrow \pi \\ S^1 \end{array}$$

4. The $C^\infty(M)$ -module $\Gamma(TM)$.
 $C^\infty(M)$ -module $\leftarrow (C^\infty(M), +, \cdot)$ (satisfies) $C^+, A^+, N^+, I+, C^+, A^+, N^+, D^+$. Not a field. A ring.

$$\begin{aligned} \text{set } \Gamma(TM) &= \{ \chi \mid M \rightarrow TM \mid \text{smooth section} \} \\ (\chi \oplus \tilde{\chi})(f) &:= (\chi f) \underbrace{+}_{C^\infty(M)} \tilde{\chi}(f) \\ (\underbrace{g}_{C^\infty(M)} \odot \xi)(f) &:= \underbrace{g}_{C^\infty(M)} \cdot \chi(f) \end{aligned}$$

$$\begin{aligned}\chi:M&\rightarrow TM\\p&\mapsto \chi(p)\\\chi f:M&\rightarrow \mathbb{R}\\p&\mapsto \chi(p)f\end{aligned}$$

$(\Gamma(TM), \oplus, \odot)$ $C^\infty(M)$ - **module**

upshot: set of all smooth vector fields can be made into a $C^\infty(M)$ -module.

Fact:

- (1) $\text{ZFC} \implies$ every vector space has a basis. (You have to have C - axiom of choice in set theory)
- (2) no such result exists for modules.

This is a shame, because otherwise, we could have chosen (for any manifolds) vector fields,

$$\chi_{(1)}, \dots, \chi_{(d)} \in \Gamma(TM)$$

and would be able to write every vector field Ξ

$$\chi = \underbrace{\qquad\qquad\qquad}_{\text{component functions}} \cdot \chi_{(i)}$$

Simple counterexample

Schuller says: Take a sphere, Morse Theorem, every smooth vector field must vanish at 2 pts. “mustn’t choose a global basis”

However: $\frac{\partial}{\partial x^i} : U \xrightarrow{\text{smooth}} TU$

$$p \mapsto \left(\frac{\partial}{\partial x^i} \right)_p$$

- 21.10. **Tensor fields.** so far
- $\Gamma(M)$ =”set of vector fields” $C^\infty(M)$ -module
- $\Gamma(T^*M)$ = “covector fields” $C^\infty(M)$ -module

Definition 73. An (r, s) -tensor field T is a multi-linear map

$$T : \underbrace{\Gamma(T^*M) \times \dots \times \Gamma(T^*M)}_r \times \Gamma(TM) \times \dots \times \Gamma(TM) \xrightarrow{\sim} C^\infty(M)$$

Example: $f \in C^\infty(M)$

$$\begin{aligned}df : \Gamma(TM) &\xrightarrow{\sim} C^\infty(M) \\ \chi &\mapsto df(\chi) := \chi[f]\end{aligned}$$

df (0, 1)-T.F. (tensor field)
where $(\chi f) \underbrace{\binom{p}{\in M}} := \underbrace{\chi(p)}_{\in T_p M} f$
can check: df is C^∞ –linear

LECTURE 7: CONNECTIONS

cf. [Lecture 7: Connections \(International Winter School on Gravity and Light 2015\)](#)
So far: saw that a vector field X can be used to provide a directional derivative

$$\nabla_X f := Xf$$

of a function $f \in C^\infty(M)$.

Remark: from now on: consider mostly vector fields.
Notational overkill?

$$\nabla_X f = Xf = (df)(X)$$

In Thought Bubble: $\nabla_X(f \cdot g) = X(fg) = (Xf) \cdot g + fX(g)$ Product rule, because it’s a derivative.
not quite:

$$\begin{aligned}X:C^\infty(M)&\rightarrow C^\infty(M)\\df:\Gamma(TM)&\rightarrow C^\infty(M)\\ \nabla_X:C^\infty(M)&\rightarrow C^\infty(M)\end{aligned}$$

$$\begin{array}{ccc} \nabla_X : C^\infty(M) & \longrightarrow & C^\infty(M) \\ \vdots \downarrow & & \vdots \downarrow \\ \nabla_X : \begin{matrix} TM^p \otimes T^*M^q \text{ i.e.} \\ \binom{p}{q} \text{ tensor field} \end{matrix} & \longrightarrow & \begin{matrix} TM^p \otimes T^*M^q \text{ i.e.} \\ \binom{p}{q} \text{ tensor field} \end{matrix} \end{array}$$

1. Directional derivatives of tensor fields. We formulate a wish list of properties which the ∇_X acting on a tensor field should have.

In Thought Bubble: Any remaining freedom in choosing ∇ , will need to be provided as additional structure beyond $(M, \mathcal{O}, \mathcal{A})$

Definition 74 (connection). In Thought Bubble: linear connection, *covariant derivative*, affine connection
A **connection** ∇ on a smooth manifold $(M, \mathcal{O}, \mathcal{A})$ is a map that takes a pair consisting of a vector (field) X and a (p, q) -tensor field T and sends them to a (p, q) -tensor (field) $\nabla_X T$ satisfying

- (i) $\nabla_X f = Xf \quad \forall f \in C^\infty(M)$
- (ii) $\nabla_X(T + S) = \nabla_X T + \nabla_X S$
- (iii) $\nabla_X(\underbrace{T(\omega, Y)}_{\in C^\infty(M)}) = (\nabla_X T)(\omega, Y) + T(\nabla_X \omega, Y) + T(\omega, \nabla_X Y)$

In Thought Bubble: for $(1, 1)$ -TF T , but analogously for any (p, q) - TF

- ”Leibnitz” rule.
- (iv) $\nabla_{fX+Z} T = f \nabla_X T + \nabla_Z T$
 $f \in C^\infty(M)$

A manifold with connection is quadruple $(M, \mathcal{O}, \mathcal{A}, \nabla)$
Remark: ∇_X is the extension of X .
 $\nabla \text{ — ” — } \text{ of } d$

2. New structure on $(M, \mathcal{O}, \mathcal{A})$ required to fix ∇ . Q: How much freedom do we have in choosing such a structure.

Consider X, Y vector fields

$$\begin{aligned} \nabla_X Y &\stackrel{\text{In Thought Bubble: } (U, x)}{=} \nabla_{X^i \frac{\partial}{\partial x^i}} \left(Y^m \frac{\partial}{\partial x^m} \right) \\ &\stackrel{\text{(iii)}}{=} \underbrace{X^i \left(\nabla_{\frac{\partial}{\partial x^i}} Y^m \right)}_{\text{(iii)}} \frac{\partial}{\partial x^m} + X^i Y^m \underbrace{\nabla_{\frac{\partial}{\partial x^i}} \left(\frac{\partial}{\partial x^m} \right)}_{\text{(iii)}} \\ &\stackrel{\text{(i)}}{=} \underbrace{X^i \left(\frac{\partial}{\partial x^i} Y^m \right)}_{\text{(i)}} \frac{\partial}{\partial x^m} + X^i Y^m \underbrace{\Gamma_{mi}^q(x)}_{\text{connection coefficient functions (on } M) \text{ of } \nabla \text{ wrt } (U, x)} \frac{\partial}{\partial x^q} \end{aligned}$$

$$\underbrace{\left(\underbrace{T}_{(p,q)} \otimes \underbrace{S}_{(r,s)} \right)}_{(p,q) \otimes (r,s)} (\omega, \dots, X, \dots) := T(\omega, \dots, X, \dots) \underbrace{\cdot}_{C^\infty(M)} S(\dots, \dots)$$

$$\nabla_X (T \otimes S) = (\nabla_X T) \otimes S + T \otimes (\nabla_X S)$$

Definition 75 (Connection coefficient functions). $(M, \mathcal{O}, \mathcal{A}, \nabla)$, $(\mathcal{U}, x) \in \mathcal{A}$.

Then the **connection coefficient functions** ("Γ"s) with respect to (wrt) (U, x) on the $(\dim(M))^3$ many functions

$$\begin{aligned} \Gamma_{jk}^i : \mathcal{U} &\rightarrow \mathbb{R} \\ p &\mapsto \left(dx^i \left(\nabla_{\frac{\partial}{\partial x^k}} \frac{\partial}{\partial x^j} \right) \right) (p) \end{aligned}$$

Thus:

$$(\nabla_X Y)^i = X^m \left(\frac{\partial}{\partial x^m} Y^i \right) + \Gamma_{nm}^i \underbrace{\cdot}_{C^\infty(M)} Y^n X^m$$

Remark: On a chart domain U , choice of the $(\dim M)^3$ functions Γ_{jk}^i suffices to fix the action of ∇ on a *vector* field.

Fortunately, the same $(\dim M)^3$ functions fix the action of ∇ on any tensor field.

key observation:

$$\nabla_{\frac{\partial}{\partial x^m}} (dx^i) = \sum_{jm}^i dx^j$$

but now:

$$\begin{aligned} \underbrace{\nabla_{\frac{\partial}{\partial x^m}} \left(dx^i \left(\frac{\partial}{\partial x^j} \right) \right)}_{\delta_j^i} &= \frac{\partial}{\partial x^m} (\delta_j^i) = 0 \\ \underbrace{\parallel}_{\text{(iii)}} & \\ &= \left(\nabla_{\frac{\partial}{\partial x^m}} dx^i \right) \left(\frac{\partial}{\partial x^j} \right) + dx^i \underbrace{\left(\nabla_{\frac{\partial}{\partial x^m}} \frac{\partial}{\partial x^j} \right)}_{\Gamma_{jm}^q \frac{\partial}{\partial x^q}} = 0 \\ \implies \left(\nabla_{\frac{\partial}{\partial x^m}} dx^i \right) \left(\frac{\partial}{\partial x^j} \right) &= -\Gamma_{jm}^i \end{aligned}$$

Summary so far:

$$\begin{aligned} (\nabla_X Y)^i &= X(Y^i) \underbrace{+}_{\text{act on } \mathbf{vector} \text{ field}} \Gamma_{jm}^i Y^j X^m \\ (\nabla_X \omega)_i &= X(\omega_i) + -\Gamma_{im}^j \omega_j X^m \end{aligned}$$

Note that for the immediately above expression for $(\nabla_X Y)^i$, in the second term on the right hand side, Γ_{jm}^i has the last entry at the bottom, m going in the direction of X , so that it matches up with X^m . This is a good mnemonic to memorize the index positions of Γ .

similarly, by further application of Leibnitz

T a $(1, 2)$ -TF (tensor field)

$$(\nabla_X T)^i_{jk} = X(T^i_{jk}) + \Gamma_{sm}^i T^s_{jk} X^m - \Gamma_{jm}^s T^i_{sk} X^m - \Gamma_{km}^s T^i_{js} X^m$$

Question: If in a Euclidean space, the Γ s all vanish in a (then existing) global chart.

Answer: Yes, but: What is a Euclidean space:

$(M = \mathbb{R}^n, \mathcal{O}_{\text{st}}, \mathcal{A})$ smooth manifold.

Assume $(\mathbb{R}^n, \text{id}_{\mathbb{R}^n}) \in \mathcal{A}$ and

$$(\Gamma_{(x)}^i)_{jk} = dx^i \left((\nabla_{\underline{E}})_{\frac{\partial}{\partial x^k}} \frac{\partial}{\partial x^j} \right) \stackrel{!}{=} 0$$

Intuition:

\mathbb{R}^2 : $\nabla_{\text{Euclidean}}$

\mathbb{R}^2 : $\nabla_{\text{Hyperbolic}}$

Definition 76. X vector field on $(M, \mathcal{O}, \mathcal{A}, \nabla)$.

Then divergence of X is the function:

$$\text{div}(X) := \left(\nabla_{\frac{\partial}{\partial x^i}} X \right)^i$$

Claim: chart-independent.

3. Change of Γ 's under change of chart.

$(U, x), (V, y) \in \mathcal{A}$ and $U \cap V \neq \emptyset$

$$\Gamma_{jk}^i(y) := dy^i \left(\nabla_{\frac{\partial}{\partial y^k}} \frac{\partial}{\partial y^j} \right) = \frac{\partial y^i}{\partial x^q} dx^q \left(\nabla_{\frac{\partial x^p}{\partial y^k} \frac{\partial}{\partial x^p}} \frac{\partial x^s}{\partial y^j} \frac{\partial}{\partial x^s} \right)$$

Note ∇_{fX} is C^∞ -linear for fX

covector dy^i is C^∞ -linear in its argument

$$\begin{aligned} \implies \Gamma_{jk}^i(y) &= \frac{\partial y^i}{\partial x^q} dx^q \left(\frac{\partial x^p}{\partial y^k} \left[\left(\nabla_{\frac{\partial}{\partial x^p}} \frac{\partial x^s}{\partial y^j} \right) \frac{\partial}{\partial x^s} + \frac{\partial x^s}{\partial y^j} \left(\nabla_{\frac{\partial}{\partial x^p}} \frac{\partial}{\partial x^s} \right) \right] \right) = \\ &= \frac{\partial y^i}{\partial x^q} \underbrace{\frac{\partial x^p}{\partial y^k} \frac{\partial}{\partial x^p}}_{\frac{\partial}{\partial y^k}} \frac{\partial}{\partial y^j} \delta_s^q + \frac{\partial y^i}{\partial x^q} \frac{\partial x^p}{\partial y^k} \frac{\partial x^s}{\partial y^j} \Gamma_{sp}^q(x) \end{aligned}$$

in summary:

$$(97) \quad \Gamma_{jk}^i(y) = \frac{\partial y^i}{\partial x^q} \frac{\partial^2 x^q}{\partial y^j \partial y^k} + \frac{\partial y^i}{\partial x^q} \frac{\partial x^s}{\partial y^j} \frac{\partial x^p}{\partial y^k} \Gamma_{sp}^q(x)$$

Eq. (97) is the change of connection coefficient function under the change of chart $(U \cap V, x) \rightarrow (U \cap V, y)$

4. **Normal Coordinates.** Let $p \in M$ of $(M, \mathcal{O}, \mathcal{A}, \nabla)$

Then one can construct a chart (U, x) with $p \in U$ such that

$$\Gamma(x)^i_{(jk)}(p) = 0$$

at the point p . **Not** necessarily in any neighborhood.

Proof. Let (V, y) be any chart (with) $p \in V$.

Thus, in general: $\Gamma(y)^i_{jk} \neq 0$

Then consider a new chart (U, x) to which one transits by virtue of

$$(x \circ y^{-1})^i(\alpha^1, \dots, \alpha^d) := \alpha^i - \frac{1}{2}\Gamma(y)^i_{(jk)}(p)\alpha^j\alpha^k$$

$$p = x^{-1}(\alpha^1, \dots, \alpha^d)$$

$$\left(\frac{\partial x^i}{\partial y^j}\right)_p = \partial_j(x^i \circ y^{-1}) = \delta_j^i - \Gamma(y)^i_{mj}(p) \alpha^m|_{\alpha=0} = \delta_j^i$$

$$\frac{\partial x^i}{\partial y^k \partial y^j}(p) = -\Gamma(y)^i_{kj}(p)$$

$$\begin{aligned} \implies \Gamma(x)^i_{jk}(p) &= \Gamma(y)^i_{jk}(p) - \Gamma(y)^i_{kj}(p) = 0 \\ &= \Gamma(y)^i_{[jk]}(p) = T(y)^i_{jk} \end{aligned}$$

Terminology: (U, x) is called a **normal coordinate chart** of ∇ at $\mathbf{p} \in \mathbf{M}$.

Tutorial 7 Connections. Exercise 1. : True or false?

- (a)
- $\nabla_{fX}Y = f\nabla_XY$ by definition so $\nabla_{fX} = f\nabla_X$ i.e. ∇_X is $C^\infty(M)$ -linear in X
 - $f \in C^\infty(M)$ is a $(0,0)$ -tensor field. $\nabla_Xf = Xf \equiv X(f)$ by definition.
 - If the manifold is flat, I'm assuming that means that the manifold is globally a Euclidean space, and by definition, $\Gamma = 0$.

$$\nabla_XY = X^j \frac{\partial}{\partial x^j}(Y^i) \frac{\partial}{\partial x^i} + \Gamma_{jk}^i Y^k X^j \frac{\partial}{\partial x^i} = X^j \frac{\partial Y^i}{\partial x^j} \frac{\partial}{\partial x^i} + 0$$

and similarly for any (p,q) -tensor field, i.e.

$$\nabla_XT = X^j \frac{\partial T^{i_1 \dots i_p}_{j_1 \dots j_q}}{\partial x^j}$$

-

$$\nabla_Xf = X^j \frac{\partial f}{\partial x^j} = X \cdot \text{grad}(f)$$

- $\forall (U, x) \in \mathcal{A}$, locally (after working out the first few cases, and doing induction, one can look up the expression for the local form; I found it in Nakahara's **Geometry, Topology and Physics**, Eq. 7.26, and it needs to be modified for the convention of order of bottom indices for Γ :

$$\nabla_\nu t^{\lambda_1 \dots \lambda_p}_{\mu_1 \dots \mu_q} = \partial_\nu t^{\lambda_1 \dots \lambda_p}_{\mu_1 \dots \mu_q} + \Gamma_{\kappa\nu}^{\lambda_1} t^{\kappa\lambda_2 \dots \lambda_p}_{\mu_1 \dots \mu_q} + \dots + \Gamma_{\kappa\nu}^{\lambda_p} t^{\lambda_1 \dots \lambda_{p-1}\kappa}_{\mu_1 \dots \mu_q} - \Gamma_{\mu_1\nu}^{\kappa} t^{\lambda_1 \dots \lambda_p}_{\kappa\mu_2 \dots \mu_q} - \dots - \Gamma_{\mu_q\nu}^{\kappa} t^{\lambda_1 \dots \lambda_p}_{\mu_1 \dots \mu_{q-1}\kappa}$$

Clearly, ∇_X is uniquely fixed $\forall p \in M$ by choosing each of the $(\dim M)^3$ many connection coefficient functions Γ .

- (b)
- $\nabla : \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$
 $\nabla : (p,q)$ -tensor field $\mapsto (p,q)$ -tensor field
 - By definition, ∇ satisfies the Leibniz rule.
 -

-
-

Exercise 2. : Practical rules for how ∇ acts Torsion-free covariant derivative boils down to a connection coefficient function Γ that is symmetric in the bottom indices.

-

$$\nabla_Xf = X(f) = X^i \frac{\partial f}{\partial x^i}$$

-

$$(\nabla_XY)^a = X^i \frac{\partial Y^a}{\partial x^i} + \Gamma_{jk}^a Y^j X^k$$

-

$$(\nabla_X\omega)_a = X^i \frac{\partial \omega_a}{\partial x^j} - \Gamma_{ak}^i \omega_i X^k$$

-

$$(\nabla_mT)^a_{bc} = \frac{\partial}{\partial x^m}(T^a_{bc}) + \Gamma_{im}^a T^i_{bc} - \Gamma_{bm}^i T^a_{ic} - \Gamma_{cm}^j T^a_{bj}$$

-

$$(\nabla_{[m}A)_{n]} = (\nabla_mA)_n - (\nabla_nA)_m = \frac{\partial A_n}{\partial x^m} - \Gamma_{nm}^i A_i - \left(\frac{\partial A_m}{\partial x^n} - \Gamma_{mn}^i A_i\right) = \frac{\partial A_m}{\partial x^m} - \frac{\partial A_m}{\partial x^n}$$

-

$$(\nabla_m\omega)_{nr} = \frac{\partial \omega_{nr}}{\partial x^m} - \Gamma_{nm}^i \omega_{ir} - \Gamma_{rm}^i \omega_{ni}$$

□

Exercise 3. : Connection coefficients

Question .

The connection coefficient functions Γ in chart $(U \cap V, y)$ is given, in terms of chart $(U \cap V, x)$ as follows:
Recall Eq. (97)

$$\Gamma_{jk}^i(y) = \frac{\partial y^i}{\partial x^q} \frac{\partial^2 x^q}{\partial y^j \partial y^k} + \frac{\partial y^i}{\partial x^q} \frac{\partial x^s}{\partial y^j} \frac{\partial x^p}{\partial y^k} \Gamma_{sp}^q(x)$$

LECTURE 8: PARALLEL TRANSPORT & CURVATURE (INTERNATIONAL WINTER SCHOOL ON GRAVITY AND LIGHT 2015)

Experiment. $\nabla_{v_\gamma}Y \stackrel{!}{=} 0$
(please move in this fashion)

round sphere
 $(S^2, \mathcal{O}, \mathcal{A}, \nabla)$

ellipsoid
 $(S^2, \mathcal{O}, \mathcal{A}, \tilde{\nabla})$

ellisoid embedded in \mathbb{R}^3 with \mathbb{R}^3 Euclidean connection
 (\mathbb{R}^3, ∇_E)

1. Parallelity of vector fields. $(M, \mathcal{O}, \mathcal{A}, \nabla)$ manifold with connection.

Definition 77 (parallelly transported, parallel). (1) A vector field X on M is said to be **parallelly transported** along a smooth curve $\gamma : \mathbb{R} \rightarrow M$ if

$$\nabla_{v_\gamma} X = 0$$

i.e.

$$\left(\nabla_{v_{\gamma, \gamma(\lambda)}} X\right)_{\gamma(\lambda)} = 0$$

(2) A slightly weaker condition than "parallelly transported" is "parallel":

$$(\nabla_{v_{\gamma, \gamma(\lambda)}} X)_{\gamma(\lambda)} = \mu(\lambda) X_{\gamma(\lambda)}$$

for $\mu : \mathbb{R} \rightarrow \mathbb{R}$

Example: Euclidean plane $(\mathbb{R}^2, \mathcal{O}, \mathcal{A}, \nabla_E)$,
parallelly transported X along γ , parallel X along γ , not parallel, not parallelly transported.

2. Autoparallely transported curves. Thought bubble: self- is "Auto"

Child: "How do I get to the toy store." Adult: "Just follow your nose."

(Move in direction of nose means move along curve whose tangent vector is your nose. Means move in the straightest line. We are not talking about shortest line because we do not have a notion of distance.)

Definition 78. A (smooth) curve $\gamma : \mathbb{R} \rightarrow M$ is called **autoparallely transported** if

(98)
$$\nabla_{v_\gamma} v_\gamma = 0$$

$$\iff (\nabla_{v_{\gamma, \gamma(\lambda)}} v_\gamma)_{\gamma(\lambda)} = 0$$

Remark: Define weaker notion of autoparallel curve
Thought bubble: !

$$\nabla_{v_\gamma} v_\gamma = \mu v_\gamma$$

Example. Euclidean plane $(\mathbb{R}^2, \mathcal{O}, \mathcal{A}, \nabla_E)$
autoparallely transported (uniform straight motion, Newton's First law), autoparallel (straight motion) but not autoparallely transported,

3. Autoparallel equation.

Autoparallely transported curve γ

Consider that portion of the curve that lies in U , where $(U, x) \in \mathcal{A}$.

$$\nabla_{v_\gamma} v_\gamma = 0$$

Express $\nabla_{v_\gamma} v_\gamma = 0$ in terms of chart representation.

$$(\nabla_{v_\gamma} v_\gamma) = \left(\nabla_{\dot{\gamma}^m(x) \left(\frac{\partial}{\partial x^m}\right)_{\gamma(\lambda)}} \dot{\gamma}^n(x) \frac{\partial}{\partial x^n}\right)$$

where

$$v_{\gamma, \gamma(\lambda)} = \dot{\gamma}^m(x) \cdot \left(\frac{\partial}{\partial x^m}\right)$$
$$\gamma^m(x) := x^m \circ \gamma$$

$$\implies (\nabla_{v_\gamma} v_\gamma) =$$
$$= \underbrace{\dot{\gamma}^m \frac{\partial \dot{\gamma}^q}{\partial x^m}}_{\ddot{\gamma}^q(x)^m} \cdot \frac{\partial}{\partial x^q} + \dot{\gamma}^m \dot{\gamma}^n \Gamma^q_{nm} \frac{\partial}{\partial x^q}$$

20200306 EY: For further clarification,

$$\dot{\gamma}(t) = v(\gamma(t)) \in T_{\gamma(t)}U$$
$$\dot{\gamma}(t) = v^i(\gamma^1(t), \dots, \gamma^n(t))$$
$$\frac{d}{dt} \dot{\gamma}^i(t) = \frac{d}{dt} v^i(\gamma^1(t), \dots, \gamma^n(t)) = \frac{d}{dt} v^i(x^1, \dots, x^n)$$
$$= \frac{\partial v^i}{\partial x^j}(x) \dot{x}^j = \dot{\gamma}^j \frac{\partial \dot{\gamma}^i}{\partial x^j}$$

in summary:

(99)

$$\ddot{\gamma}^m_{(x)}(\lambda) + (\Gamma^m_{(x)})_{ab}(\gamma(\lambda)) \dot{\gamma}^a_{(x)}(\lambda) \dot{\gamma}^b_{(x)}(\lambda) = 0$$

chart expression of the condition that γ be autoparallely transported.

Examples.

(a) Euclidean plane, $U = \mathbb{R}^2$, $x = \text{id}_{\mathbb{R}}$, $\Gamma(x)^i_{jk} = 0$

$$\implies \ddot{\gamma}(x)^m = 0 \implies \gamma(x)^m(\lambda) = a^m \lambda + b^m a, b \in \mathbb{R}^d$$

(b) round sphere $(S^2, \mathcal{O}, \mathcal{A}, \nabla_{\text{round}})$

Consider a chart:

$$x(p) = (\theta, \varphi)$$
$$\theta \in (0, \pi)$$
$$\varphi \in (0, 2\pi)$$

all other $(\Gamma^1_{11} = 0)$ vanish.

$$\Gamma(x)^1_{22}(x^{-1}(\theta, p)) := -\sin \theta \cos \theta \quad \Gamma(x)^2_{21} = \Gamma(x)^2_{12} := \cot \theta$$

Sloppy notation (classical mechanics)

$$x^1(p) = \theta(p)$$
$$x^2(p) = \varphi(p)$$

autoparallel equation

$$\ddot{\theta} + \Gamma^1_{22} \dot{\varphi} \dot{\varphi} = 0$$
$$\ddot{\varphi} + 2\Gamma^2_{12} \dot{\theta} \dot{\varphi} = 0$$

$$\iff \ddot{\theta} - \sin(\theta) \cos(\theta) \dot{\varphi} \dot{\varphi} = 0$$
$$\ddot{\varphi} + 2 \cot(\theta) \dot{\theta} \dot{\varphi} = 0$$

Solve: e.g. $\theta(\lambda) = \frac{\pi}{2} \quad \varphi(\lambda) = \underline{\omega} \cdot \lambda + \varphi_0$
Similarly,
 $\nabla_{\text{potato}}, (S^2, \mathcal{O}, \mathcal{A})$

3. Torsion. Q: Can one use ∇ to define tensors on $(M, \mathcal{O}, \mathcal{A}, \nabla)$?

Definition 79. The *torsion* of a connection ∇ is the $(1, 2)$ -**tensor field**

(100)
$$T(\omega, X, Y) := \omega(\nabla_X Y - \nabla_Y X - [X, Y])$$

(Inside a cloud)
[X, Y] vector field defined by

$$[X, Y]f := X(Yf) - Y(Xf)$$

Proof. check T is C^∞ -linear in each entry

$$\begin{aligned} T(f \cdot \omega, X, Y) &= f \cdot \omega(\dots) = fT(\omega, X, Y) \\ T(\omega + \psi, X, Y) &= \dots = T(\omega, X, Y) + T(\psi, X, Y) \\ T(\omega, fX, Y) &= \omega(\nabla_{fX} Y - \nabla_Y(fX) - [fX, Y]) = \\ &= \omega(f\nabla_X Y - (Yf)X - f\nabla_Y X - f[X, Y] + (Yf)X) = fT(\omega, X, Y) \end{aligned}$$

since

$$\begin{aligned} [fX, Y]g &= fX(Yg) - Y(fXg) = \\ &= fX(Yg) - (Yf)(Xg) - fY(Xg) = \\ &= f \cdot [X, Y]g - (Yf)Xg = \\ &= (f \cdot [X, Y] - (Yf)X)g \\ T(\omega, X, Y) &= -T(\omega, Y, X) \quad \checkmark \end{aligned}$$

Definition 80. A $(M, \mathcal{O}, \mathcal{A}, \nabla)$ is called *torsion-free* if $T = 0$

In a chart

$$\begin{aligned} T^i_{ab} &:= T\left(dx^i, \frac{\partial}{\partial x^a}, \frac{\partial}{\partial x^b}\right) = dx^i(\dots) \\ &= \Gamma^i_{ab} - \Gamma^i_{ba} = 2\Gamma^i_{[ab]} \end{aligned}$$

From now on, in these lectures, we only use torsion-free connections.

4. Curvature.

Definition 81. The *Riemann curvature* of a connection ∇ is the $(1, 3)$ -tensor field

(101)
$$Riem(\omega, Z, X, Y) := \omega(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z)$$

Proof. do it: C^∞ -linear in each slot.

Tutorials $Riem^i_{jab} = \dots$

Algebraic relevance of Riem.

$$(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z) = R(\cdot, Z, X, Y) + \nabla_{[X, Y]} Z$$

In one chart (U, x)

$$\underbrace{\left(\nabla_a\right)}_{=\nabla\frac{\partial}{\partial x^a}}\nabla_bZ-\nabla_b\nabla_aZ)^m=Riem^m_{nab}Z^n+\nabla\left[\underbrace{\frac{\partial}{\partial x^a},\frac{\partial}{\partial x^b}}_{=0}\right]Z$$

$$\left(\nabla_a\nabla_bZ\right)^m-\left(\nabla_b\nabla_aZ\right)^m=Riem^m_{nab}Z^n$$

Geometric significance.

Idea:
[X, Y] = 0

If [X, Y] \neq 0

$$(\delta Z)^m = \dots = R^m_{nab}X^aY^bZ^n\delta s\delta t + \mathcal{O}(\delta s^2\delta t, \delta t^2\delta s)$$

Holonomy

TUTORIAL 8 PARALLEL TRANSPORT & CURVATURE

Exercise 1.

Exercise 2. : Where connection coefficients appear

It was suggested in the tutorial sheets and hinted in the lecture that the following should be committed to memory.

Question : Recall the autoparallel equation for a curve γ .

(a)

$$\nabla_{v_\gamma} v_\gamma = 0$$

(b)

□

$$\begin{aligned} \nabla_{v_\gamma} v_\gamma &= \nabla_{\dot{\gamma}\frac{\partial}{\partial x^\mu}} v_\gamma = \dot{\gamma}^\nu \nabla_{\partial_\nu} v_\gamma = \dot{\gamma}^\nu \left[\frac{\partial v^\mu_\gamma}{\partial x^\nu} + \Gamma^\rho_{\mu\nu} v^\mu_\gamma \right] \frac{\partial}{\partial x^\rho} = \dot{\gamma}^\nu \left[\frac{\partial \dot{\gamma}^\rho}{\partial x^\nu} + \Gamma^\rho_{\mu\nu} \dot{\gamma}^\mu \right] \frac{\partial}{\partial x^\rho} = 0 \\ &\implies \boxed{\ddot{\gamma}^\rho + \Gamma^\rho_{\mu\nu} \dot{\gamma}^\mu \dot{\gamma}^\nu} \end{aligned}$$

as, for example, for $F(x(t))$,

$$\frac{dF(x(t))}{dt} = \dot{x} \frac{\partial F}{\partial x} = \frac{d}{dt} F$$

so that

$$\dot{\gamma}^\nu \frac{\partial v^\mu_\gamma}{\partial x^\nu} = \frac{d}{d\lambda} v^\mu_\gamma = \frac{d^2}{d\lambda^2} \gamma^\mu$$

Question : Determine the coefficients of the Riemann tensor with respect to a chart (U, x) .

Recall this manifestly covariant definition

□

$$Riem(\omega, Z, X, Y) = \omega(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z)$$

We want R^i_{jab} .
now

$$\nabla_X \nabla_Y Z = \nabla_X \left((Y^\mu \frac{\partial}{\partial x^\mu} Z^\rho + \Gamma^\rho_{\mu\nu} Z^\mu Y^\nu) \frac{\partial}{\partial x^\rho} \right) = (X^\alpha \frac{\partial}{\partial x^\alpha} (Y^\mu \frac{\partial}{\partial x^\mu} Z^\rho + \Gamma^\rho_{\mu\nu} Z^\mu Y^\nu) + \Gamma^\rho_{\alpha\beta} (Y^\mu \frac{\partial}{\partial x^\mu} Z^\alpha + \Gamma^\alpha_{\mu\nu} Z^\mu Y^\nu) X^\beta) \frac{\partial}{\partial x^\rho}$$

For $X = \partial_a, Y = \partial_b, Z = \partial_j$, then the partial derivatives of the coefficients of the input vectors become zero.

$$\implies \nabla_{\partial_a} \nabla_{\partial_b} \partial_j = \frac{\partial}{\partial x^a} (\Gamma^i_{jb}) + \Gamma^i_{\alpha a} \Gamma^\alpha_{jb}$$

Now

$$[X, Y]^i = X^j \frac{\partial}{\partial x^j} Y^i - Y^j \frac{\partial X^i}{\partial x^j}$$

For coordinate vectors, $[\partial_i, \partial_j] = 0 \ \forall i, j = 0, 1 \dots d$.

Thus

$$R^i{}_{jab} = \frac{\partial}{\partial x^a} \Gamma^i_{jb} - \frac{\partial}{\partial x^b} \Gamma^i_{ja} + \Gamma^i_{\alpha a} \Gamma^\alpha_{jb} - \Gamma^i_{\alpha b} \Gamma^\alpha_{ja}$$

Question : $\text{Ric}(X,Y) := \text{Riem}^m_{amb} X^a Y^b$ **define** $(0,2)$ -**tensor?**.

Yes, transforms as such:

EY developments. I roughly follow the spirit in Theodore Frankel’s **The Geometry of Physics: An Introduction** Second Ed. 2003, Chapter 9 Covariant Differentiation and Curvature, Section 9.3b. The Covariant Differential of a Vector Field. P.S. EY : 20150320 I would like a copy of the Third Edition but I don’t have the funds right now to purchase the third edition: go to my tilt crowdfunding campaign, <http://ernestyalumni.tilt.com>, and help with your financial support if you can or send me a message on my various channels and ernestyalumni gmail email address if you could help me get a hold of a digital or hard copy as a pro bono gift from the publisher or author.

The spirit of the development is the following:

“How can we express connections and curvatures in terms of forms?” -Theodore Frankel.

From Lecture 7, connection ∇ on vector field Y , in the “direction” X ,

$$\nabla_{\frac{\partial}{\partial x^k}} Y = \left(\frac{\partial Y^i}{\partial x^k} + \Gamma^i_{jk} Y^j \right) \frac{\partial}{\partial x^i}$$

Make the ansatz (approche, impostazione) that the connection ∇ acts on Y , the vector field, first:

$$\nabla Y(X) = \left(X^k \frac{\partial Y^i}{\partial x^k} + \Gamma^i_{jk} Y^j X^k \right) \frac{\partial}{\partial x^i} = X^k \left(\nabla_{\frac{\partial}{\partial x^k}} Y \right)^i \frac{\partial}{\partial x^i} = (\nabla_X Y)^i \frac{\partial}{\partial x^i} = \nabla_X Y$$

Now from Lecture 7, Definition for Γ ,

$$dx^i \left(\nabla_{\frac{\partial}{\partial x^k}} \frac{\partial}{\partial x^j} \right) = \Gamma^i_{jk}$$

Make this ansatz (approche, impostazine)

$$\nabla \frac{\partial}{\partial x^j} = (\Gamma^i_{jk} dx^k) \otimes \frac{\partial}{\partial x^i} \in \Omega^1(M, TM) = T^*M \otimes TM$$

where $\Omega^1(M, TM) = T^*M \otimes TM$ is the set of all TM or vector-valued 1-forms on M , with the 1-form being the following:

$$\Gamma^i_{jk} dx^k = \Gamma^i_j \in \Omega^1(M) \quad \begin{array}{l} i = 1 \dots \dim(M) \\ j = 1 \dots \dim(M) \end{array}$$

So Γ^i_j is a $\dim M \times \dim M$ matrix of 1-forms (EY !!!).

Thus

$$\nabla Y = (d(Y^i) + \Gamma^i_j Y^j) \otimes \frac{\partial}{\partial x^i}$$

So the connection is a (smooth) map from TM to the set of all vector-valued 1-forms on M , $\Omega^1(M, TM)$, and then, after “eating” a vector Y , yields the “covariant derivative”:

$$\nabla : TM \rightarrow \Omega^1(M, TM) = T^*M \otimes TM$$

$$\nabla : Y \mapsto \nabla Y$$

$$\nabla Y : TM \rightarrow TM$$

$$\nabla Y(X) \mapsto \nabla Y(X) = \nabla_X(Y)$$

Now

$$\left[\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j} \right] f = \frac{\partial}{\partial x^i} \left(\frac{\partial}{\partial x^j} \right) - \frac{\partial}{\partial x^j} \left(\frac{\partial}{\partial x^i} \right) = 0$$

(this is okay as on $p \in (U, x)$; x -coordinates on same chart (U, x))

EY : 20150320 My question is when is this nontrivial or nonvanishing (i.e. not equal to 0).

$$[e_a, e_b] = ?$$

for a frame (e_c) and would this be the difference between a tangent bundle TM vs. a (general) vector bundle?

Wikipedia helps here. cf. wikipedia, “Connection (vector bundle)”

$$\nabla : \Gamma(E) \rightarrow \Gamma(T^*M \otimes E) = \Omega^1(M, E)$$

$$\nabla e_a = \omega^c_{ab} f^b \otimes e_c$$

$$f^b \in T^*M \text{ (this is the dual basis for } TM \text{ and, note, this is for the manifold, } M$$

$$\nabla_{f_b} e_a = \omega^c_{ab} e_c \in E$$

$$\omega^c_a = \omega^c_{ab} f^b \in \Omega^1(M)$$

is the connection 1-form, with $a, c = 1 \dots \dim V$. EY : 20150320 This V is a vector space living on each of the fibers of E . I know that $\Gamma(T^*M \otimes E)$ looks like it should take values in E , but it’s meaning that it takes vector values of V . Correct me if I’m wrong: ernestyalumni at gmail and various social media.

Let $\sigma \in \Gamma(E)$, $\sigma = \sigma^a e_a$

$$\nabla \sigma = (d\sigma^c + \omega^c_{ab} \sigma^a f^b) \otimes e_c \text{ with}$$

$$d\sigma^c = \frac{\partial \sigma^c}{\partial x^b} f^b$$

$$\implies \nabla_X \sigma = \left(X^b \frac{\partial \sigma^c}{\partial x^b} + \omega^c_{ab} \sigma^a X^b \right) e_c = X^b \left(\frac{\partial \sigma^c}{\partial x^b} + \omega^c_{ab} \sigma^a \right) e_c$$

LECTURE 9: NEWTONIAN SPACETIME IS CURVED!

Axiom 1 (Newton I:). *A body on which no force acts moves uniformly along a straight line*

Axiom 2 (Newton II:). *Deviation of a body’s motion from such uniform straight motion is effected by a force, reduced by a factor of the body’s reciprocal mass.*

Remark:

- (1) 1st axiom - in order to be relevant - must be read as a measurement prescription for the geometry of space ...

(2) Since gravity universally acts on every particle, in a universe with at least two particles, gravity must not be considered a force if Newton I is supposed to remain applicable.

21.11. **Laplace’s questions.** Laplace *1749

†1827

Q: “Can gravity be encoded in a curvature of space, such that its effects show if particles under the influence of (no other) force we postulated to move along straight lines in this curved space?”

Answer: No!

Proof. gravity is a force point of view

$$\begin{aligned} m\ddot{x}^\alpha(t) &= F^\alpha(x(t)) \\ m\dot{x}^\alpha(t) &= \underbrace{mf^\alpha}_{F^\alpha}(x(t)) \end{aligned}$$

$-\partial_\alpha f^\alpha = 4\pi G\rho$ (Poisson)
 ρ mass density of matter

$$m\ddot{x}^\alpha(t) = \underbrace{mf^\alpha}_{F^\alpha}(x(t))$$

True?
(EY : 20150330) You know this, $F = Gm_1m_2/r^2$
Yes
weak equivalence principle

$$\ddot{x}^\alpha(t) - f^\alpha(x(t)) = 0$$

Laplace asks: Is this $(\ddot{x}(t))$ of the form

$$\ddot{x}^\alpha(t) + \Gamma^\alpha_{\beta\gamma}(x(t))\dot{x}^\beta(t)\dot{x}^\gamma(t) = 0$$

Conclusion: One cannot find Γ ’s such that Newton’s equation takes the form of an autoparallel.

□

Question (from audience) We can evaluate the autoparallel equation pointwise?! But at each point, we can set the Gammas to zero?!

Then, one should be able to write Newton’s second law in the usual form?

Prof. Schuller: you (observer) fall with the mass (i.e. accelerated reference frame) and so you transform Γ ’s to be 0. The problem with this is if you do the same experiment in the North pole and fall with the body. If someone else at the South Pole does the same experiment at the same time, with that same transformation (of reference frames), the effect of gravity cannot be transformed out.

In a homogeneous gravitational field, you can possibly transform away gravity, $\Gamma = 0$. But in an inhomogeneous gravitational field, no.

2. The full wisdom of Newton I. use also the information from Newton’s first law that particles (no force) move uniformly introduce the appropriate setting to talk about the difference easily
insight: in spacetime uniform & straight motion in space is simply **straight** motion
So let’s try in spacetime:
let $x : \mathbb{R} \rightarrow \mathbb{R}^3$

$$\begin{aligned} X : \mathbb{R} &\rightarrow \mathbb{R}^4 \\ t &\mapsto (t, x^1(t), x^2(t), x^3(t)) := \\ &:= (X^0(t), X^1(t), X^2(t), X^3(t)) \end{aligned}$$

be a particle’s trajectory in space \longleftrightarrow worldline (history) of the particle

That’s all it takes:

Trivial rewritings:

$$\dot{X}^0 = 1$$

$$\begin{aligned} \implies \begin{cases} \ddot{X}^0 &= 0 \\ \ddot{X}^\alpha - f^\alpha(X(t)) \cdot \dot{X}^0 \cdot \dot{X}^0 &= 0 \end{cases} & (\alpha = 1, 2, 3) \implies \begin{cases} a = 0, 1, 2, 3 \\ \ddot{X}^a + \Gamma^a_{bc} \dot{X}^b \dot{X}^c = 0 \end{cases} \\ & \text{antoparallel eqn in } \underline{\text{spacetime}} \end{aligned}$$

Yes, choosing $\Gamma^0_{ab} = 0$

$$\Gamma^\alpha_{\beta\gamma} = 0 = \Gamma^\alpha_{0\beta} = \Gamma^\alpha_{\beta 0}$$

only: $\Gamma^\alpha_{00} \stackrel{!}{=} -f^\alpha$

Question: Is this a coordinate-choice artifact?

No, since $R^\alpha_{0\beta 0} = -\frac{\partial}{\partial x^\beta} f^\alpha$ (only non-vanishing components) (tidal force tensor, – the Hessian of the force component)

Ricci tensor $\implies R_{00} = R^m_{0m0} = -\partial_\alpha f^\alpha = 4\pi G\rho$

Poisson: $-\partial_\alpha f^\alpha = 4\pi G \cdot \rho$

writing: $T_{00} = \frac{1}{2}\rho$

$$\implies \boxed{R_{00} = 8\pi GT_{00}}$$

Einstein in 1912 $R_{ab} \longrightarrow 8\pi GT_{ab}$

Conclusion: Laplace’s idea works in spacetime

Remark

$$\begin{aligned} \Gamma^\alpha_{00} &= -f^\alpha \\ R^\alpha_{\beta\gamma\delta} &= 0 \quad \alpha, \beta, \gamma, \delta = 1, 2, 3 \\ \boxed{R_{00} = 4\pi G\rho} \end{aligned}$$

Q: What about transformation behavior of LHS of

$$\underbrace{\ddot{x}^a + \Gamma^a_{bc} \dot{X}^b \dot{X}^c}_{\underbrace{(\nabla_{v_X} v_X)^a}_{:= a^a \text{ “acceleration vector”}}} = 0$$

3. The foundations of the geometric formulation of Newton’s axiom. new start

Definition 82. A *Newtonian spacetime* is a quintuple

$$(M, \mathcal{O}, \mathcal{A}, \nabla, t)$$

where $(M, \mathcal{O}, \mathcal{A})$ 4-dim. smooth manifold

$$t : M \rightarrow \mathbb{R} \text{ smooth function}$$

(i) “There is an absolute space”

$$(dt)_p \neq 0 \quad \forall p \in M$$

(ii) “absolute time flows uniformly”

$$\nabla dt \underbrace{=}_\text{space of (0,2)-tensor fields} 0 \quad \text{everywhere}$$

∇dt is a (0,2)-tensor field

(iii) add to axioms of Newtonian spacetime $\nabla = 0$ torsion free

Definition 83. *absolute space at time τ*

$$S_\tau := \{p \in M | t(p) = \tau\}$$
$$\xrightarrow{dt \neq 0} M = \coprod S_\tau$$

Definition 84. *A vector $X \in T_pM$ is called*

(a) *future-directed if*

$$dt(X) > 0$$

(b) *spatial if*

$$dt(X) = 0$$

(c) *past-directed if*

$$dt(X) < 0$$

picture
Newton I: The worldline of a particle under the influence of no force (gravity isn't one, anyway) is a future-directed autoparallel i.e.

$$\nabla_{v_X} v_X = 0$$
$$dt(v_X) > 0$$

and (iii) ∇ is torsion-free.

Newton II:

$$\nabla_{v_X} v_X = \frac{F}{m} \iff m \cdot \mathfrak{a} = F$$

where F is a spatial vector field:

$$dt(F) = 0$$

Convention: restrict attention to atlases $\mathcal{A}_{\text{stratified}}$ whose charts (\mathcal{U}, x) have the property

$$\begin{array}{l} x^0 : \mathcal{U} \rightarrow \mathbb{R} \\ x^1 : \mathcal{U} \rightarrow \mathbb{R} \\ \vdots \\ x^3 \end{array} \quad x^0 = t|_{\mathcal{U}} \quad \implies \quad \begin{array}{l} 0 \\ 0 = \nabla_{\frac{\partial}{\partial x^a}} dx^0 = -\Gamma^0_{ba} \end{array} \quad \begin{array}{l} \text{“absolute time flows uniformly”} \\ a = 0, 1, 2, 3 \end{array} \quad \nabla dt$$

Let's evaluate in a chart (\mathcal{U}, x) of a stratified atlas $\mathcal{A}_{\text{sheet}}$: Newton II:

$$\nabla_{v_X} v_X = \frac{F}{m}$$

in a chart.

$$(X^0)'' + \Gamma^0_{cd} (X^a)' (X^b)' \Big|_{\text{stratified atlas}} = 0$$
$$(X^\alpha)'' + \Gamma^\alpha_{\gamma\delta} X^{\gamma'} X^{\delta'} + \Gamma^\alpha_{00} X^{0'} X^{0'} + 2\Gamma^\alpha_{\gamma 0} X^{\gamma'} X^{0'} = \frac{F^\alpha}{m} \quad \alpha = 1, 2, 3$$

EY: 20160623: where the factor of 2 comes from torsion free ∇

$$\implies (X^0)''(\lambda) = 0 \implies X^0(\lambda) = a\lambda + b \quad \text{constants } a, b \quad \text{with}$$

$$X^0(\lambda) = (x^0 \circ X)(\lambda) \stackrel{\text{stratified}}{=} (t \circ X)(\lambda)$$

convention parametrize worldline by absolute time

$$\frac{d}{d\lambda} = a \frac{d}{dt}$$

$$a^2 \ddot{X}^\alpha + a^2 \Gamma^\alpha_{\gamma\delta} \dot{X}^\gamma \dot{X}^\delta + a^2 \Gamma^\alpha_{00} \dot{X}^0 \dot{X}^0 + 2a^2 \Gamma^\alpha_{\gamma 0} \dot{X}^\gamma \dot{X}^0 = \frac{F^\alpha}{m}$$
$$\implies \underbrace{\ddot{X}^\alpha + \Gamma^\alpha_{\gamma\delta} \dot{X}^\gamma \dot{X}^\delta + \Gamma^\alpha_{00} \dot{X}^0 \dot{X}^0 + 2\Gamma^\alpha_{\gamma 0} \dot{X}^\gamma \dot{X}^0}_{\mathfrak{a}^\alpha} = \frac{1}{a^2} \frac{F^\alpha}{m}$$

Student question: How can you get the Lagrange equations in this formalism?

Schuller says: In *all* systems they take the form like this:

$$L = \underbrace{T}_{\frac{m}{2} g(v_x, v_x)} + V$$

where v_x is velocity to the spatial curves x .

Schuller says: $\frac{m}{2} \sqrt{g(v_x, v_x)}$, parameter invariance, you get straight curves; if you take $\sqrt{}$ away, $\frac{m}{2} g(v_x, v_x)$ you get straight, *uniform* curves.

Schuller says: Only in inertial systems, which is defined as:

$$\nabla_{\frac{\partial}{\partial x^a}} \frac{\partial}{\partial x^b} = 0$$

inertial frame, non-accelerating, non-rotating.

Schuller says: Apart from g , a 2-tensor, you need to know ω .

Schuller references 2nd. year Mechanik in German: <https://www.video.uni-erlangen.de/course/id/272>

22. LECTURE 10: METRIC MANIFOLDS

cf. [Lecture 10: Metric Manifolds \(International Winter School on Gravity and Light 2015\)](#)

We establish a structure on a smooth manifold that allows one to assign vectors in each tangent space a length (and an angle between vectors in the same tangent space).

From this structure, one can then define a notion of length of a curve.

Then we can look at shortest curves.

Requiring then that the shortest curves coincide with the straightest curves (wrt ∇) will result in ∇ being determined by the metric structure.

$$g \stackrel{\text{straight=short}}{\underset{T=0}{\rightsquigarrow}} \nabla \rightsquigarrow \text{Riem}$$

22.1. Metrics.

Definition 85. *A metric g on a smooth manifold $(M, \mathcal{O}, \mathcal{A})$ is a $(0, 2)$ -tensor field satisfying*

- (i) *symmetry* $g(X, Y) = g(Y, X) \quad \forall X, Y \text{ vector fields}$
- (ii) *non-degeneracy: the musical map*

$$\text{“flat” } \flat : \Gamma(TM) \rightarrow \Gamma(T^*M)$$
$$X \mapsto \flat(X)$$

$$\text{where } \flat(X)(Y) := g(X, Y)$$

$$\flat(X) \in \Gamma(T^*M)$$

$$\text{In thought bubble: } \flat(X) = g(X, \cdot)$$

... is a C^∞ -isomorphism in other words, it is invertible.

Remark: $(\flat(X))_a$ or

X_a

$$(\flat(X))_a := g_{am} X^m$$

Thought bubble: $\flat^{-1} = \sharp$

$\flat^{-1}(\omega)^a := g^{am}\omega_m$
 $\flat^{-1}(\omega)^a := (g^{\text{“}^{-1}\text{”}})^{am}\omega_m \implies \text{not needed. (all of this is not needed)}$

Definition 86. The $(2,0)$ -tensor field $g^{\text{“}^{-1}\text{”}}$ with respect to a metric g is the symmetric

$$g^{\text{“}^{-1}\text{”}} : \Gamma(T^*M) \times \Gamma(T^*M) \rightarrow C^\infty(M)$$
$$(\omega, \sigma) \mapsto \omega(\flat^{-1}(\sigma)) \qquad \flat^{-1}(\sigma) \in \Gamma(TM))$$

chart: $g_{ab} = g_{ba}$
 $(g^{-1})^{am}g_{mb} = \delta_b^a$

Example: $(S^2, \mathcal{O}, \mathcal{A})$
chart (\mathcal{U}, x)
 $\varphi \in (0, 2\pi)$
 $\theta \in (0, \pi)$
define the metric

$$g_{ij}(x^{-1}(\theta, \varphi)) = \begin{bmatrix} R^2 & 0 \\ 0 & R^2 \sin^2 \theta \end{bmatrix}_{ij}$$

$R \in \mathbb{R}^+$
“the metric of the round sphere of radius R ”

22.2. **Signature.** Linear algebra:

$A^a{}_m v^m = \lambda v^a$

$g_{am}v^m = \lambda \cdot v^a?$ \rightsquigarrow

$$\begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix}$$

$$\begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & -1 & & \\ & & & & \ddots & \\ & & & & & -1 \\ & & & & & & 0 \\ & & & & & & & \ddots \\ & & & & & & & & 0 \end{pmatrix}$$

$(1,1)$ tensor has eigenvalues
 $(0,2)$ has signature (p,q) (well-defined)

$\left. \begin{matrix} (+ + +) \\ (+ + -) \\ (+ - -) \\ (- - -) \end{matrix} \right\} d + 1 \text{ if } p + q = \dim V$

Definition 87. A metric is called **Riemannian** if its signature is $(+ + \cdots +)$
Lorentzian if $(+ - \cdots -)$

22.3. **Length of a curve.** Let γ be a smooth curve.
Then we know its velocity $v_{\gamma, \gamma(\lambda)}$ at each $\gamma(\lambda) \in M$.

Definition 88. On a Riemannian metric manifold $(M, \mathcal{O}, \mathcal{A}, g)$, the **speed** of a curve at $\gamma(\lambda)$ is the number

$$(\sqrt{g(v_\gamma, v_\gamma)})_{\gamma(\lambda)} = s(\lambda)$$

F. Schuller: “I feel the need for speed.” -Top Gun.
(I feel the need for speed, then I feel the need for a metric)

Aside: $[v^a] = \frac{1}{T}$
 $[g_{ab}] = L^2$
 $[\sqrt{g_{ab}v^av^b}] = \sqrt{\frac{L^2}{T^2}} = \frac{L}{T}$

Definition 89. Let $\gamma : (0,1) \rightarrow M$ a smooth curve.
Then the **length of γ** is the number

$$\mathbb{R} \ni L[\gamma] := \int_0^1 d\lambda s(\lambda) = \int_0^1 d\lambda \sqrt{(g(v_\gamma, v_\gamma))_{\gamma(\lambda)}}$$

F. Schuller: “velocity is more fundamental than speed, speed is more fundamental than length”

Example: reconsider the round sphere of radius R
Consider its equator:

$\theta(\lambda) := (x^1 \circ \gamma)(\lambda) = \frac{\pi}{2}$
 $\varphi(\lambda) := (x^2 \circ \gamma)(\lambda) = 2\pi\lambda^3$
 $\theta'(\lambda) = 0$
 $\varphi'(\lambda) = 6\pi\lambda^2$

on the same chart $g_{ij} = \begin{bmatrix} R^2 & \\ & R^2 \sin^2 \theta \end{bmatrix}$

F.Schuller: do everything in this chart

$$L[\gamma] = \int_0^1 d\lambda \sqrt{g_{ij}(x^{-1}(\theta(\lambda), \varphi(\lambda)))(x^i \circ \gamma)'(\lambda)(x^j \circ \gamma)'(\lambda)} = \int_0^1 d\lambda \sqrt{R^2 \cdot 0 + R^2 \sin^2(\theta(\lambda))36\pi^2\lambda^4} =$$
$$= 6\pi R \int_0^1 d\lambda \lambda^2 = 6\pi R [\frac{1}{3}\lambda^3]_0^1 = 2\pi R$$

Theorem 26. $\gamma : (0,1) \rightarrow M$ and
 $\sigma : (0,1) \rightarrow (0,1)$ smooth bijective and increasing “reparametrization”

$$L[\gamma] = L[\gamma \circ \sigma]$$

Proof. \implies Tutorials

□

22.4. **Geodesics.**

Definition 90. A curve $\gamma : (0,1) \rightarrow M$ is called a **geodesic** on a Riemannian manifold $(M, \mathcal{O}, \mathcal{A}, g)$ if its a stationary curve with respect to a length functional L .

Thought bubble: in classical mechanics, deform the curve a little, ϵ times this deformation, to first order, it agrees with $L[\gamma]$

Theorem 27. γ geodesic iff it satisfies the Euler-Lagrange equations for the Lagrangian

$$\mathcal{L} : TM \rightarrow \mathbb{R}$$
$$X \mapsto \sqrt{g(X, X)}$$

In a chart, the Euler Lagrange equations take the form:

$$\left(\frac{\partial \mathcal{L}}{\partial \dot{x}^m}\right)^{\cdot} - \frac{\partial \mathcal{L}}{\partial x^m} = 0$$

F.Schuller: this is a chart dependent formulation
here:

$$\mathcal{L}(\gamma^i, \dot{\gamma}^i) = \sqrt{g_{ij}(\gamma(\lambda))\dot{\gamma}^i(\lambda)\dot{\gamma}^j(\lambda)}$$

Euler-Lagrange equations:

$$\frac{\partial \mathcal{L}}{\partial \dot{\gamma}^m} = \frac{1}{\sqrt{\dots}} g_{mj}(\gamma(\lambda)) \dot{\gamma}^j(\lambda)$$

$$\left(\frac{\partial \mathcal{L}}{\partial \dot{\gamma}^m}\right)^{\cdot} = \left(\frac{1}{\sqrt{\dots}}\right)^{\cdot} g_{mj}(\gamma(\lambda)) \cdot \dot{\gamma}^j(\lambda) + \frac{1}{\sqrt{\dots}} (g_{mj}(\gamma(\lambda)) \ddot{\gamma}^j(\lambda) + \dot{\gamma}^s (\partial_s g_{mj}) \dot{\gamma}^j(\lambda))$$

Thought bubble: reparametrize $g(\dot{\gamma}, \dot{\gamma}) = 1$ (it's a condition on my reparametrization)
By a clever choice of reparametrization $(\frac{1}{\sqrt{\dots}})^{\cdot} = 0$

$$\frac{\partial \mathcal{L}}{\partial \gamma^m} = \frac{1}{2\sqrt{\dots}} \partial_m g_{ij}(\gamma(\lambda)) \dot{\gamma}^i(\lambda) \dot{\gamma}^j(\lambda)$$

putting this together as Euler-Lagrange equations:

$$g_{mj} \ddot{\gamma}^j + \partial_s g_{mj} \dot{\gamma}^s \dot{\gamma}^j - \frac{1}{2} \partial_m g_{ij} \dot{\gamma}^i \dot{\gamma}^j = 0$$

Multiply on both sides $(g^{-1})^{qm}$

$$\ddot{\gamma}^q + (g^{-1})^{qm} (\partial_i g_{mj} - \frac{1}{2} \partial_m g_{ij}) \dot{\gamma}^i \dot{\gamma}^j = 0$$

$$\ddot{\gamma}^q + (g^{-1})^{qm} \frac{1}{2} (\partial_i g_{mj} + \partial_j g_{mi} - \partial_m g_{ij}) \dot{\gamma}^i \dot{\gamma}^j = 0$$

geodesic equation for γ in a chart.

$$(g^{-1})^{qm} \frac{1}{2} (\partial_i g_{mj} + \partial_j g_{mi} - \partial_m g_{ij}) =: \Gamma^q_{ij}(\gamma(\lambda))$$

Thought bubble: $\left(\frac{\partial \mathcal{L}}{\partial \xi_x^{a+\dim M}}\right)^{\cdot}_{\sigma(x)} - \left(\frac{\partial \mathcal{L}}{\partial x_i^a}\right)^{\cdot}_{\sigma(x)} = 0$

Definition 91. “Christoffel symbol” $^{L.C.}\Gamma$ are the connection coefficient functions of the so-called Levi-Civita connection $^{L.C.}\nabla$

We usually make this choice of ∇ if g is given.

$(M, \mathcal{O}, \mathcal{A}, g) \rightarrow (M, \mathcal{O}, \mathcal{A}, g, ^{L.C.}\nabla)$

abstract way: $\nabla g = 0$ and $T = 0$ (torsion)

$\implies \overline{\nabla} = ^{L.C.}\nabla$

Definition 92. (a) The Riemann-Christoffel curvature is defined by

$$R_{abcd} := g_{am} R^m_{bcd}$$

(b) Ricci: $R_{ab} = R^m_{amb}$

Thought bubble: with a metric, $^{L.C.}\nabla$

(c) (Ricci) scalar curvature:

$$R = g^{ab} R_{ab}$$

Thought bubble: $^{L.C.}\nabla$

Definition 93. Einstein curvature $(M, \mathcal{O}, \mathcal{A}, g)$

$$G_{ab} := R_{ab} - \frac{1}{2} g_{ab} R$$

Convention: $g^{ab} := (g^{-1})^{ab}$

F. Schuller: these indices are not being pulled up, because what would you pull them up with (student) Question: Does the Einstein curvature yield new information?

Answer:

$$g^{ab} G_{ab} = R_{ab} g^{ab} - \frac{1}{2} g_{ab} g^{ab} R = R - \delta_a^a R = R - \frac{1}{2} \dim M R = (1 - \frac{d}{2}) R$$

Tutorial 9: Metric manifolds. Exercise 3: Levi-Civita Connection. Suppose torsion-free $T = 0$ and metric-compatible connection $\nabla g = 0$

Question Recall $T = 0$ on a chart.

$$\Gamma^c_{ba} = \frac{1}{2} (g^{-1})^{cm} \left(\frac{\partial g_{bm}}{\partial x^a} + \frac{\partial g_{ma}}{\partial x^b} - \frac{\partial g_{ab}}{\partial x^m} \right)$$

or

$$\Gamma^a_{bc} = \frac{1}{2} (g^{-1})^{am} \left(\frac{\partial g_{bm}}{\partial x^c} + \frac{\partial g_{mc}}{\partial x^b} - \frac{\partial g_{bc}}{\partial x^m} \right)$$

23. SYMMETRY

EY : 20150321 This lecture tremendously and lucidly clarified, for me at least, what a symmetry of the Lie algebra is, and in comparing structures $(M, \mathcal{O}, \mathcal{A})$ vs. $(M, \mathcal{O}, \mathcal{A}, \nabla)$, clarified differences, and asking about differences is a good way to learn, the difference between \mathcal{L} and ∇ , respectively.

Feeling that the round sphere

$$(S^2, \mathcal{O}, \mathcal{A}, g^{\text{round}})$$

has rotational symmetry, while
the potato

$$(S^2, \mathcal{O}, \mathcal{A}, g^{\text{potato}})$$

does not.

23.1.

23.2. Important

23.3. **Flow of a complete vector field.** Let $(M, \mathcal{O}, \mathcal{A})$ smooth X vector field on M

Definition 94. A curve $\gamma : I \subseteq \mathbb{R} \rightarrow M$ is called an integral curve of X if

$$v_{\gamma, \gamma(\lambda)} = X_{\gamma(\lambda)}$$

Definition 95. A vector filed X is ***complete*** if all integral curves have $I = \mathbb{R}$ EY: 20150321 (i.e. domain is all of \mathbb{R})

Ex. minute 48:30 EY : reall good explanation by F.P.Schuller; take a pt. out for an incomplete vector field.

Theorem 28. compactly supported smooth vector field is complete.

Definition 96. The flow of a complete vector field X is a 1-parameter family

$$h^X = \mathbb{R} \times M \rightarrow M$$

where $\gamma_p : \mathbb{R} \rightarrow M$ is the integral curve of X with

$$\begin{aligned} \gamma(0) &= p \\ \text{Then for fixed } \lambda \in \mathbb{R} \end{aligned}$$

$$h^X_\lambda : M \rightarrow M \text{ smooth}$$

picture $h^X_\lambda(S) \neq S$ (if $X \neq 0$)

23.4. **Lie subalgebras of the Lie algebra $(\Gamma(TM), [\cdot, \cdot])$ of vector fields.**

- (a) $\Gamma(TM) = \{ \text{ set of all vector fields } \}$ $C^\infty(M)$ -module = \mathbb{R} -vector space
 $\implies [X, Y] \in \Gamma(TM)$ $[X, Y]f := X(Yf) - Y(Xf)$
- (i) $[X, Y] = -[Y, X]$
(ii) $[\lambda X + Z, Y] = \lambda[X, Y] + [Z, Y]$
(iii) $[X, [Y, Z]] + [Z, [X, Y]] + [Y, [Z, X]] = 0$
 $(\Gamma(TM), [\cdot, \cdot])$ Lie algebra
- (b) Let $X_1 \dots X_s$ for s (many) vector fields on M , such that

Tutorial 11 Symmetry. Exercise 1. : True or false?

- (a)
 -
 - $\phi^* : T^*N \rightarrow T^*M$ i.e. $\phi^*\nu(X) = \nu(\phi_*X)$ for smooth $\phi : M \rightarrow N$, so the pullback of a covector $\nu \in T^*N$ maps to a covector in T^*M .
 -
 -
 -
 -
- (b)
- (c)

Exercise 2. : Pull-back and push-forward

Question . Let’s check this locally

$$\begin{aligned} \phi^*(df)(X) &= (df)(\phi_*X) = (df)(X^i \frac{\partial y^j}{\partial x^i} \frac{\partial}{\partial y^j}) = X^i \frac{\partial y^j}{\partial x^i} \frac{\partial f}{\partial y^j} \text{ where } \phi_*X = X^i \frac{\partial y^j}{\partial x^i} \frac{\partial}{\partial y^j} \\ d(\phi^*f)(X) &= d(f(\phi))(X) = \frac{\partial f}{\partial y^j} \frac{\partial y^j}{\partial x^i} dx^i(X) = X^i \frac{\partial y^j}{\partial x^i} \frac{\partial f}{\partial y^j} \end{aligned}$$

So

$$\boxed{\phi^*(df) = d(\phi^*f)}$$

$$\forall p \in M, \forall X \in \mathfrak{X}(M)$$

The big idea is that this is a showing of the **naturality** of the pullback ϕ^* with d , i.e. that this commutes:

$$\begin{array}{ccc} \Omega^1(M) & \xleftarrow{\phi^*} & \Omega^1(N) \\ d \uparrow & & d \uparrow \\ C^\infty(M) & \xleftarrow{\phi^*} & C^\infty(N) \end{array}$$

Question .

$$(\phi_*)^a_b := (dy^a)(\phi_*(\frac{\partial}{\partial x^b}))$$

Let $g \in C^\infty(N)$

$$\begin{aligned} \phi_* \left(\frac{\partial}{\partial x^b} \right) g &= \frac{\partial x^b}{g} \phi(p) = \frac{\partial}{\partial x^b} g \phi x^{-1} x(p) = \frac{\partial}{\partial x^b} (g y y^{-1} \phi x^{-1})(x) = \\ &= \frac{\partial}{\partial x^b} (g y^{-1} (y \phi x^{-1}(x(p)))) = \frac{\partial g}{\partial y} \bigg|_y \frac{\partial y^a}{\partial x^b} \bigg|_x = \frac{\partial y^a}{\partial x^b} \frac{\partial g}{\partial y^a} \end{aligned}$$

Then

$$\phi_* \left(\frac{\partial}{\partial x^b} \right) = \frac{\partial y^a}{\partial x^b} \frac{\partial}{\partial y^a}$$

and so

$$(\phi_*)^a_b = \frac{\partial y^a}{\partial x^b}$$

Question .

Exercise 3. :Lie derivative-the pedestrian way

Question . While it is true that $\forall p \in S^2$, for $x(p) = (\theta, \varphi)$, and $(yix^{-1})(\theta, \varphi) = (y^1, y^2, y^3) \in \mathbb{R}^3$ and that, at this point

p , $(y^1)^2/a^2 + (y^2)^2/b^2 + (y^3)^2/c^3 = 1$, this doesn’t imply (EY: 20150321 I think) that, globally, it’s an ellipsoid (yet). In the familiar charts given, spherical chart $(U, x) \in \mathcal{A}$ and $(\mathbb{R}^3, y = \text{id}_{\mathbb{R}^3}) \in \mathcal{B}$ it looks like an ellipsoid, but change to another choice of charts, and it could look something very different.

Question .

Equip $(\mathbb{R}^3, \mathcal{O}_{\text{st}}, \mathcal{B})$ with the Euclidean metric g , and pullback g .
Note that the pullback of the inclusion from \mathbb{R}^3 onto S^2 for the Euclidean metric is the following:

$$i^*g \left(\frac{\partial}{\partial \theta^i}, \frac{\partial}{\partial \theta^j} \right) = g \left(i_* \frac{\partial}{\partial \theta^i}, i_* \frac{\partial}{\partial \theta^j} \right) = g \left(\frac{\partial x^a}{\partial \theta^i} \frac{\partial}{\partial x^a}, \frac{\partial x^b}{\partial \theta^j} \frac{\partial}{\partial x^b} \right) = g_{ab} \frac{\partial x^a}{\partial \theta^i} \frac{\partial x^b}{\partial \theta^j}$$

With $g_{ab} = \delta_{ab}$, the usual Euclidean metric, this becomes the following:

$$g_{ij}^{\text{ellipsoid}} = \frac{\partial x^a}{\partial \theta^i} \frac{\partial x^a}{\partial \theta^j}$$

At this point, one should get smart (we are in the 21st century) and use some sort of CAS (Computer Algebra System). I like Sage Math (version 6.4 as of 20150322). I also like the Sage Manifolds package for Sage Math.

I like Sage Math for the following reasons:

- Open source, so it’s open and freely available to anyone, which fits into my principle of making online education open and freely available to anyone, anytime
- Sage Math structures everything in terms of Category Theory and Categories and Morphisms naturally correspond to Classes and Class methods or functions in Object-Oriented Programming in Python and they’ve written it that way

and I like Sage Manifolds for roughly the same reasons, as manifolds are fit into a category theory framework that’s written into the Python code. e.g.

```
sage: S2 = Manifold(2, 'S^2', r'\mathbb{S}^2', start_index=1) ; print S2
sage: print S2
2-dimensional manifold 'S^2'
sage: type(S2)
<class 'sage.geometry.manifolds.manifold.Manifold_with_category'>
```

With code (I’ve provided for convenience; you can make your own as I wrote it based upon to example of S^2 on the sage-manifolds documentation website page), load it and do the following:

```
cf. https://github.com/ernestyalumni/diffgeo-by-sagemnfd/blob/master/S2.sage
http://sagemanifolds.obspm.fr/examples.html
sage: load("S2.sage")
sage: U_ep = S2.open_subset('U_{ep}')
sage: eps.<the,phi> = U_ep.chart()
sage: a = var('a')
sage: b = var('b')
sage: c = var('c')
sage: inclus = S2.diff_mapping(R3, {(eps, cart): [ a*cos(phi)*sin(the), b*sin(phi)*sin(the),c*cos(the) ]} , name="inc",latex_name=r'\mathcal{i}')
sage: inclus.pullback(h).display()
inc_*(h) = (c^2*sin(the)^2 + (a^2*cos(phi)^2 + b^2*sin(phi)^2)*cos(the)^2) dthe*dthe - (a^2 - b^2)*cos(phi)*cos(the)*sin(phi)*sin(the) dthe*dphi
- (a^2 - b^2)*cos(phi)*cos(the)*sin(phi)*sin(the) dphi*dthe + (b^2*cos(phi)^2 + a^2*sin(phi)^2)*sin(the)^2 dphi*dphi
sage: inclus.pullback(h)[2,2].expr()
(b^2*cos(phi)^2 + a^2*sin(phi)^2)*sin(the)^2
```

A new open subset U_{ep} was declared in S^2 , a new chart $(U_{\text{ep}}, (\theta, \phi))$ was declared, the constants, a, b, c , were declared, and the inclusion map given in the problem

$$y \circ \mathbf{i} \circ x^{-1} : (\theta, \phi) \mapsto (a \cos \phi \sin \theta, b \sin \phi \sin \theta, c \cos \theta)$$

Then the pullback of the inclusion map \rangle was done on the Euclidean metric h , defined earlier in the file

`S2.sage`

. Then one can access the components of this metric and do, for example,

`simplify_full()`, `full_simplify()`, `reduce_trig()`

on the expression.

In Python, I could easily do this, and give an answer quick in LaTeX:

```
sage: for i in range(1,3):
.....:     for j in range(1,3):
.....:         print inclus.pullback(h)[i,j].expr()
.....:         latex(inclus.pullback(h)[i,j].expr() )
.....:
c^2*sin(the)^2 + (a^2*cos(phi)^2 + b^2*sin(phi)^2)*cos(the)^2
```

(EY: I’ll suppress the LaTeX output but this sage math function gives you LaTeX code)

and so

$$\begin{aligned} i^*g = & c^2 \sin (the)^2 + \left(a^2 \cos (\phi)^2 + b^2 \sin (\phi)^2\right) \cos (the)^2 d\theta \otimes d\theta + \\ & -2\left(a^2 - b^2\right) \cos (\phi) \cos (the) \sin (\phi) \sin (the) d\theta \otimes d\phi + \\ & + \left(b^2 \cos (\phi)^2 + a^2 \sin (\phi)^2\right) \sin (the)^2 d\phi \otimes d\phi \end{aligned}$$

Question .

```
sage: polar_vees = eps.frame()
sage: X_1 = - sin(phi) * polar_vees[1] - cot( the ) * cos(phi) * polar_vees[2]
sage: X_2 = cos( phi ) * polar_vees[1] - cot( the ) * sin( phi ) * polar_vees[2]
sage: X_3 = polar_vees[2]
sage: X_2.lie_der(X_1).display()
(cos(the)^2 - 1)/sin(the)^2 d/dphi
sage: X_3.lie_der(X_1).display()
cos(phi) d/dthe - cos(the)*sin(phi)/sin(the) d/dphi
sage: X_3.lie_der(X_2).display()
sin(phi) d/dthe + cos(phi)*cos(the)/sin(the) d/dphi
Indeed, one can check on a scalar field  $f_{\text{eps}} \in C^\infty(S^2)$ :
sage: f_eps = S2.scalar_field({eps: function('f', the, phi ) }, name='f' )
sage: (X_1( X_2(f_eps)) - X_2(X_1(f_eps) ) ).display()
U_{ep} --> R
(the, phi) |--> -D[1](f)(the, phi)
sage: X_2.lie_der(X_1) == -X_3
True
sage: X_3.lie_der(X_1) == X_2
True
sage: X_3.lie_der(X_2) == -X_1
True
```

$$\implies \boxed{[X_i, X_j] = -\epsilon_{ijk} X_k}$$

So $\text{span}_{\mathbb{R}}\{X_1, X_2, X_3\}$ equipped with $[\ , \]$ constitute a Lie subalgebra on S^2 (It’s closed under $[\ , \]$

24. INTEGRATION

24.1.

24.2.

24.3. Volume forms.

Definition 97. *On a smooth manifold $(M, \mathcal{O}, \mathcal{A})$ a $(0, \dim M)$ -tensor field Ω is called a volume form if*

- (a) Ω vanishes nowhere (i.e. $\Omega \neq 0 \ \forall p \in M$)
- (b) *totally antisymmetric*

$$\Omega(\dots, \underbrace{X}_{ith}, \dots, \underbrace{Y}_{jth}, \dots) = -\Omega(\dots, \underbrace{Y}_{ith}, \dots, \underbrace{X}_{jth}, \dots)$$

In a chart:

$$\Omega_{i_1 \dots i_d} = \Omega_{[i_1 \dots i_d]}$$

Example $(M, \mathcal{O}, \mathcal{A}, g)$ metric manifold
construct volume form Ω from g
In any chart: (U, x)

$$\Omega_{i_1 \dots i_d} := \sqrt{\det(g_{ij}(x))} \epsilon_{i_1 \dots i_d}$$

where **Levi-Civita symbol** $\epsilon_{i_1 \dots i_d}$ is defined as $\epsilon_{123 \dots d} = +1$
 $\epsilon_{1 \dots d} = \epsilon_{[i_1 \dots i_d]}$

Proof. (well-defined) Check: What happens under a change of charts

$$\begin{aligned} \Omega(y)_{i_1 \dots i_d} &= \sqrt{\det(g(y)_{ij})} \epsilon_{i_1 \dots i_d} = \\ &= \sqrt{\det(g_{mn}(x) \frac{\partial x^m}{\partial y^i} \frac{\partial x^n}{\partial y^j}) \frac{\partial y^{m_1}}{\partial x^{i_1}} \dots \frac{\partial y^{m_d}}{\partial x^{i_d}}} \epsilon_{[m_1 \dots m_d]} = \\ &= \sqrt{|\det g_{ij}(x)|} \left| \det \left(\frac{\partial x}{\partial y} \right) \right| \det \left(\frac{\partial y}{\partial x} \right) \epsilon_{i_1 \dots i_d} = \sqrt{\det g_{ij}(x) \epsilon_{i_1 \dots i_d}} \operatorname{sgn} \left(\det \left(\frac{\partial x}{\partial y} \right) \right) \end{aligned}$$

EY : 20150323
Consider the following:

$$\begin{aligned} \Omega(y)(Y_{(1)} \dots Y_{(d)}) &= \Omega(y)_{i_1 \dots i_d} Y_{(1)}^{i_1} \dots Y_{(d)}^{i_d} = \\ &= \sqrt{\det(g_{ij}(y))} \epsilon_{i_1 \dots i_d} Y_{(1)}^{i_1} \dots Y_{(d)}^{i_d} = \\ &= \sqrt{\det(g_{mn}(x)) \frac{\partial x^m}{\partial y^i} \frac{\partial x^n}{\partial y^j} \epsilon_{i_1 \dots i_d} \frac{\partial y^{i_1}}{\partial x^{m_1}} \dots \frac{\partial y^{i_d}}{\partial x^{m_d}}} X^{m_1} \dots X^{m_d} = \\ &= \sqrt{\det(g_{mn}(x)) \frac{\partial x^m}{\partial y^i} \frac{\partial x^n}{\partial y^j}} \det \left(\frac{\partial y}{\partial x} \right) \epsilon_{m_1 \dots m_d} X^{m_1} \dots X^{m_d} = \\ &= \sqrt{\det(g_{mn}(x))} \left| \det \left(\frac{\partial x}{\partial y} \right) \right| \det \left(\frac{\partial y}{\partial x} \right) \epsilon_{m_1 \dots m_d} X^{m_1} \dots X^{m_d} = \\ &= \sqrt{\det(g_{mn}(x))} \epsilon_{m_1 \dots m_d} \operatorname{sgn} \left(\det \left(\frac{\partial x}{\partial y} \right) \right) X^{m_1} \dots X^{m_d} = \operatorname{sgn}(\det \left(\frac{\partial x}{\partial y} \right)) \Omega_{m_1 \dots m_d}(x) X^{m_1} \dots X^{m_d} \end{aligned}$$

If $\det \left(\frac{\partial y}{\partial x} \right) > 0$,

$$\Omega(y)(Y_{(1)} \dots Y_{(d)}) = \Omega(x)(X_{(1)} \dots X_{(d)})$$

This works also if Levi-Civita symbol $\epsilon_{i_1 \dots i_d}$ doesn't change at all under a change of charts. (around 42:43 <https://youtu.be/2XpnbvPy-Zg>)

Alright, let's require,
restrict the smooth atlas \mathcal{A}
to a subatlas (\mathcal{A}^\uparrow still an atlas)

$$\mathcal{A}^\uparrow \subseteq \mathcal{A}$$

s.t. $\forall (U, x), (V, y)$ have chart transition maps $y \circ x^{-1}$
 $x \circ y^{-1}$

s.t. $\det \left(\frac{\partial y}{\partial x} \right) > 0$
such \mathcal{A}^\uparrow called an **oriented** atlas

$$(M, \mathcal{O}, \mathcal{A}, g) \implies (M, \mathcal{O}, \mathcal{A}^\uparrow, g)$$

Note: associated bundles.
Note also: $\det \left(\frac{\partial y^b}{\partial x^a} \right) = \det(\partial_a(y^b x^{-1}))$ $\frac{\partial y^b}{\partial x^a}$ is an endomorphism on vector space V .

$\varphi : V \rightarrow V$
 $\det \varphi$ independent of choice of basis
 g is a $(0, 2)$ tensor field, not endomorphism (not independent of choice of basis) $\sqrt{|\det(g_{ij}(y))|}$

Definition 98. Ω be a volume form on $(M, \mathcal{O}, \mathcal{A}^\uparrow)$ and consider chart (U, x)

Definition 99. $\omega_{(X)} := \Omega_{i_1 \dots i_d} \epsilon^{i_1 \dots i_d}$ same way $\epsilon^{12 \dots d} = +1$
 $\epsilon^{[\dots]}$
one can show

$$\omega_{(y)} = \det \left(\frac{\partial x}{\partial y} \right) \omega_{(x)}$$

scalar density

24.4. Integration on one chart domain U .

Definition 100.

(102)

$$\int_U f : \stackrel{(U, y)}{=} \int_{y(U)} d^d \beta \omega_{(y)}(y^{-1}(\beta)) f_{(y)}(\beta)$$

Proof. : Check that it's (well-defined), how it changes under change of charts

$$\begin{aligned} \int_U f : \stackrel{(U, y)}{=} \int_{y(U)} d^d \beta \omega_{(y)}(y^{-1}(\beta)) f_{(y)}(\beta) &= \stackrel{(U, y)}{=} \int_{x(U)} \int d^d \alpha \left| \det \left(\frac{\partial y}{\partial x} \right) \right| f_{(x)}(\alpha) \omega_{(x)}(x^{-1}(\alpha)) \det \left(\frac{\partial x}{\partial y} \right) = \\ &= \int_{x(U)} d^d \alpha \omega_{(x)}(x^{-1}(x)) f_{(x)}(\alpha) \end{aligned}$$

□

On an oriented metric manifold $(M, \mathcal{O}, \mathcal{A}^\uparrow, g)$

$$\int_U f := \int_{x(U)} d^d \alpha \underbrace{\sqrt{\det(g_{ij}(x))(x^{-1}(\alpha))}}_{\sqrt{g}} f_{(x)}(\alpha)$$

24.5. Integration on the entire manifold.

25. LECTURE 13: RELATIVISTIC SPACETIME

Recall, from Lecture 9, the definition of Newtonian spacetime

$$(M, \mathcal{O}, \mathcal{A}, \nabla, t)$$

$$\begin{aligned} &\nabla \text{ torsion free} \\ &t \in C^\infty(M) \\ &dt \neq 0 \\ &\nabla dt = 0 \quad (\text{uniform time}) \end{aligned}$$

and the definition of relativistic spacetime (before Lecture)

$$(M, \mathcal{O}, \mathcal{A}^\uparrow, \nabla, g, T)$$

$$\begin{aligned} &\nabla \text{ torsion-free} \\ &g \text{ Lorentzian metric}(+ - - -) \\ &T \text{ time-orientation} \end{aligned}$$

25.1. Time orientation.

Definition 101. $(M, \mathcal{O}, \mathcal{A}^\uparrow, g)$ a Lorentzian manifold. Then a time-orientation is given by a vector field T that

- (i) does **not** vanish anywhere

(ii) $g(T, T) > 0$

Newtonian vs. relativistic

Newtonian

X was called future-directed if

$$dt(X) > 0$$

$\forall p \in M$, take half plane, half space of $T_p M$

also stratified atlas so make planes of constant t straight

relativistic

half cone $\forall p, q \in M$, half-cone $\subseteq T_p M$

This definition of spacetime

Question

I see how the cone structure arises from the new metric. I don’t understand however, how the T , the time orientation, comes in

Answer

$(M, \mathcal{O}, \mathcal{A}, g)$ $g \leftarrow (+ - - -)$

requiring $g(X, X) > 0$, select cones

T chooses which cone

This definition of spacetime has been made to enable the following physical postulates:

- (P1) The worldline γ of a massive particle satisfies
- (i) $g_{\gamma(\lambda)}(v_{\gamma, \gamma(lambda)}, v_{\gamma, \gamma(\lambda)}) > 0$

(ii) $g_{\gamma(\lambda)}(T, v_{\gamma, \gamma(\lambda)}) > 0$
- (P2) Worldlines of massless particles satisfy
- (i) $g_{\gamma(\lambda)}(v_{\gamma, \gamma(\lambda)}, v_{\gamma, \gamma(\lambda)}) = 0$

(ii) $g_{\gamma(\lambda)}(T, v_{\gamma, \gamma(\lambda)}) > 0$
- picture: spacetime:

Answer (to a question) T is a smooth vector field, T determines future vs. past, “general relativity: we have such a time orientation; smoothness makes it less arbitrary than it seems” -FSchuller,

Claim: 9/10 of a metric are determined by the cone

spacetime determined by distribution, only one-tenth error

25.2. Observers. $(M, \mathcal{O}, \mathcal{A}^\uparrow, \nabla, g, T)$

Definition 102. An observer is a worldline γ with

$$g(v_\gamma, v_\gamma) > 0$$

$$g(T, v_\gamma) > 0$$

together with a choice of basis

$$v_{\gamma, \gamma(\lambda)} \equiv e_0(\lambda), e_1(\lambda), e_2(\lambda), e_3(\lambda)$$

of each $T_{\gamma(\lambda)} M$ where the observer worldline passes, if $g(e_a(\lambda), e_b(\lambda)) = \eta_{ab} = \begin{bmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & -1 \end{bmatrix}_{ab}$

precise: observer = smooth curve in the frame bundle LM over M

25.2.1. Two physical postulates.

(P3) A **clock** carried by a specific observer (γ, e) will measure a **time**

$$\tau := \int_{\lambda_0}^{\lambda_1} d\lambda \sqrt{g_{\gamma(\lambda)}(v_{\gamma, \gamma(\lambda)}, v_{\gamma, \gamma(\lambda)})}$$

between the two “events”

$$\gamma(\lambda_0)$$

“start the clock”

and

$$\gamma(\lambda_1)$$

“stop the clock”

Compare with Newtonian spacetime:

$$t(p) = 7$$

Thought bubble: proper time/eigentime τ

$$M = \mathbb{R}^4$$

Application/Example. $\mathcal{O} = \mathcal{O}_{\text{st}}$

$$\mathcal{A} \ni (\mathbb{R}^4, \text{id}_{\mathbb{R}^4})$$

$$g : g_{(x)ij} = \eta_{ij} \quad ; \quad T_{(x)}^i = (1, 0, 0, 0)^i$$

$$\implies \Gamma_{(x) \ jk}^i = 0 \text{ everywhere}$$

$$\implies (M, \mathcal{O}, \mathcal{A}^\uparrow, g, T, \nabla) \quad \text{Riem} = 0$$

\implies spacetime is flat

This situation is called special relativity.

Consider two observers:

$$\gamma : (0, 1) \rightarrow M$$

$$\gamma_{(x)}^i = (\lambda, 0, 0, 0)^i$$

$$\delta : (0, 1) \rightarrow M$$

$$\alpha \in (0, 1) : \delta_{(x)}^i = \begin{cases} (\lambda, \alpha \lambda, 0, 0)^i & \lambda \leq \frac{1}{2} \\ (\lambda, (1 - \lambda)\alpha, 0, 0)^i & \lambda > \frac{1}{2} \end{cases}$$

let’s calculate:

$$\tau_\gamma := \int_0^1 \sqrt{g_{(x)ij} \dot{\gamma}_{(x)}^i \dot{\gamma}_{(x)}^j} = \int_0^1 d\lambda 1 = 1$$

$$\tau_\delta := \int_0^{1/2} d\lambda \sqrt{1 - \alpha^2} + \int_{1/2}^1 \sqrt{1^2 - (-\alpha)^2} = \int_0^1 \sqrt{1 - \alpha^2} = \sqrt{1 - \alpha^2}$$

Note: piecewise integration

Taking the clock postulate (P3) seriously, one better come up with a realistic clock design that supports the postulate.
idea.

- (P4) Postulate
- 2 little mirrors
- Let (γ, e) be an observer, and δ be a *massive* particle worldline that is parametrized s.t. $g(v_\gamma, v_\gamma) = 1$ (for parametrization/normalization convenience)
- Suppose the observer and the particle *meet* somewhere (in spacetime)

$$\delta(\tau_2) = p = \gamma(\tau_1)$$

This observer measures the 3-velocity (spatial velocity) of this particle as

$$(103) \quad v_\delta : \epsilon^\alpha(v_{\delta, \delta(\tau_2)})e_\alpha \quad \alpha = 1, 2, 3$$

where $\epsilon^0, \boxed{\epsilon^1, \epsilon^2, \epsilon^3}$ is the unique dual basis of $e_0, \boxed{e_1, e_2, e_3}$

EY:20150407

There might be a major correction to Eq. (103) from the Tutorial 14 : Relativistic spacetime, matter, and Gravitation, see the second exercise, Exercise 2, third question:

$$(104) \quad v := \frac{\epsilon^\alpha(v_\delta)}{\epsilon^0(v_\delta)}e_\alpha$$

Consequence: An observer (γ, e) will extract quantities measurable in his laboratory from objective spacetime quantities always like that.

Ex: F Faraday (0, 2)-tensor of electromagnetism:

$$F(e_a, e_b) = F_{ab} = \begin{bmatrix} 0 & E_1 & E_2 & E_3 \\ -E_1 & 0 & B_3 & -B_2 \\ -E_2 & -B_3 & 0 & B_1 \\ -E_3 & B_2 & -B_1 & 0 \end{bmatrix}$$

observer frame e_a, e_b

$E_\alpha := F(e_0, e_\alpha)$
 $B^\gamma := F(e_\alpha, e_\rho)\epsilon^{\alpha\beta\gamma}$ where $\epsilon^{123} = +1$ totally antisymmetric

25.3. **Role of the Lorentz transformations.** Lorentz transformations emerge as follows:

Let (γ, e) and $(\tilde{\gamma}, \tilde{e})$ be observers with $\gamma(\tau_1) = \tilde{\gamma}(\tau_2)$
(for simplicity $\gamma(0) = \tilde{\gamma}(0)$)
Now

$$\begin{array}{ll} e_0, \dots, e_1 & \text{at } \tau = 0 \\ \text{and } \tilde{e}_0, \dots, \tilde{e}_1 & \text{at } \tau = 0 \end{array}$$

both bases for the same $T_{\gamma(0)}M$

Thus: $\tilde{e}_a = \Lambda^b_a e_b \quad \Lambda \in GL(4)$

Now:

$$\begin{aligned} \eta_{ab} = g(\tilde{e}_a, \tilde{e}_b) &= g(\Lambda^m_a e_m, \Lambda^n_b e_n) = \\ &= \Lambda^m_a \Lambda^n_b \underbrace{g(e_m, e_n)}_{\eta_{mn}} \end{aligned}$$

i.e. $\Lambda \in O(1, 3)$

Result: Lorentz transformations relate the *frames* of *any two observers* at the same point.
“ $\tilde{x}^\mu - \Lambda^\mu_\nu x^\nu$ ” is utter nonsense

Tutorial. I didn’t see a tutorial video for this lecture, but I saw that the Tutorial sheet number 14 had the relevant topics. Go there.

two types of matter
point matter
field matter
point matter
massive point particle
more of a phenomenological importance
field matter
electromagnetic field
more fundamental from the GR point of view
both classical matter types

26.1. **Point matter.** Our postulates (P1) and (P2) already constrain the possible particle worldlines.
But what is their precise law of motion, possibly in the presence of “forces”,

- (a) without external forces

$$S_{\text{massive}}[\gamma] := m \int d\lambda \sqrt{g_{\gamma(\lambda)}(v_{\gamma, \gamma(\lambda)}, v_{\gamma, \gamma(\lambda)})}$$

with:

$$g_{\gamma(\lambda)}(T_{\gamma(\lambda)}, v_{\gamma, \gamma(\lambda)}) > 0$$

dynamical law Euler-Lagrange equation

similarly

$$S_{\text{massless}}[\gamma, \mu] = \int d\lambda \mu g(v_{\gamma, \gamma(\lambda)}, v_{\gamma, \gamma(\lambda)})$$

$$\begin{array}{ll} \delta_\mu & g(v_{\gamma, \gamma(\lambda)}, v_{\gamma, \gamma(\lambda)}) = 0 \\ \delta_\gamma & \text{e.o.m.} \end{array}$$

Reason for describing equations of motion by actions is that composite systems have an action that is the sum of the actions of the parts of that system, possibly including “interaction terms.”

Example.

$$S[\gamma] + S[\delta] + S_{\text{int}}[\gamma, \delta]$$

- (b) presence of external forces
or rather presence of fields to which a particle “couples”
Example

$$S[\gamma; A] = \int d\lambda m \sqrt{g_{\gamma(\lambda)}(v_{\gamma, \gamma(\lambda)}, v_{\gamma, \gamma(\lambda)})} + qA(v_{\gamma, \gamma(\lambda)})$$

where A is a **covector field** on M . A fixed (e.g. the electromagnetic potential)

Consider Euler-Lagrange eqns. $L_{\text{int}} = qA_{(x)}\dot{\gamma}^m_{(x)}$

$$m(\nabla_{v_\gamma} v_\gamma)_a + \underbrace{\left(\frac{\partial L_{\text{int}}}{\partial \dot{\gamma}^m_{(x)}} \right)}_* - \frac{\partial L_{\text{int}}}{\partial \gamma^m_{(x)}} = 0 \implies \boxed{\begin{array}{l} m(\nabla_{v_\gamma} v_\gamma)^a = \underbrace{-qF^a_m \dot{\gamma}^m}_{\text{Lorentz force on a charged particle in an electromagnetic field}} \end{array}}$$

$$\frac{\partial L}{\partial \dot{\gamma}^a} = qA_{(x)a}, \quad \left(\frac{\partial L}{\partial \dot{\gamma}^m_{(x)}} \right) = q \cdot \frac{\partial}{\partial x^m}(A_{(x)m}) \cdot \dot{\gamma}^m_{(x)}$$

$$\frac{\partial L}{\partial \gamma^a} = q \cdot \frac{\partial}{\partial x^a}(A_{(x)m}) \dot{\gamma}^m$$

$$* = q \left(\frac{\partial A_a}{\partial x^m} - \frac{\partial A_m}{\partial x^a} \right) \dot{\gamma}^m_{(x)} = q \cdot F_{(x)am} \dot{\gamma}^m_{(x)}$$

$F \leftarrow$ Faraday

$$S[\gamma] = \int (m\sqrt{g(v_\gamma,v_\gamma)} + qA(v_\gamma))d\lambda$$

26.2. **Field matter.**

Definition 103. *Classical (non-quantum) field matter is any tensor field on spacetime where equations of motion derive from an action.*

Example:

$$S_{\text{Maxwell}}[A] = \frac{1}{4} \int_M d^4x \sqrt{-g} F_{ab} F_{cd} g^{ac} g^{bd}$$

- A $(0,1)$ -tensor field
- = thought cloud: for simplicity one chart covers all of M
- for $\sqrt{-g}$ $(+ - - -)$

$F_{ab} := 2\partial_{[a}A_{b]} = 2(\nabla_{[a}A)_{b]}$
Euler-Lagrange equations for fields

$$0 = \frac{\partial \mathcal{L}}{\partial A_m} - \frac{\partial}{\partial x^s} \left(\frac{\partial \mathcal{L}}{\partial \partial_s A_m} \right) + \frac{\partial}{\partial x^s} \frac{\partial}{\partial x^t} \frac{\partial^2 \mathcal{L}}{\partial \partial_t \partial_s A_m}$$

Example ...

$$(\nabla_{\frac{\partial}{\partial x^m}} F)^{ma} = j^a$$

inhomogeneous Maxwell
thought bubble $j = qv_\gamma$

$$\partial_{[a}F_{b]} - ()$$

homogeneous Maxwell
Other example well-liked by textbooks

$$S_{\text{Klein-Gordon}}[\phi] := \int_M d^4x \sqrt{-g} [g^{ab}(\partial_a \phi)(\partial_b \phi) - m^2 \phi^2]$$

ϕ $(0,0)$ -tensor field

26.3. **Energy-momentum tensor of matter fields.** At some point, we want to write down an action for the metric tensor field itself.

But then, this action $S_{\text{grav}}[g]$ will be added to any $S_{\text{matter}}[A,\phi,\dots]$ in order to describe the total system.

$$S_{\text{total}}[g,A] = S_{\text{grav}}[g] + S_{\text{Maxwell}}[A,g]$$

$$\delta A \quad \implies \text{Maxwell's equations}$$

$$\delta g_{ab} \quad : \quad \boxed{\frac{1}{16\pi G} G^{ab}} + (-2T^{ab}) = 0$$

G Newton’s constant

$$G^{ab} = 8\pi G_N T^{ab}$$

Definition 104. $S_{\text{matter}}[\Phi,g]$ is a matter action, the ***so-called energy-momentum tensor*** is

$$T^{ab} := \frac{-2}{\sqrt{-g}} \left(\frac{\partial \mathcal{L}_{\text{matter}}}{\partial g_{ab}} - \partial_s \frac{\partial \mathcal{L}_{\text{matter}}}{\partial \partial_s g_{ab}} + \dots \right)$$

- of $\frac{-2}{\sqrt{g}}$ is Schrödinger minus (EY : 20150408 F.Schuller’s joke? but wise)
- choose all sign conventions s.t.

$$T(\epsilon^0,\epsilon^0) > 0$$

Example: For S_{Maxwell} :

$$T_{ab} = F_{am}F_{bn}g^{mn} - \frac{1}{4}F_{mn}F^{mn}g_{ab}$$

$$T_{ab} \equiv T_{\text{Maxwell}ab}$$

$$T(e_0,e_0) = \underline{E}^2 + \underline{B}^2$$

$$T(e_0,e_\alpha) = (E \times B)_\alpha$$

Fact: One often does not specify the fundamental action for some matter, but one is rather satisfied to assume certain properties / forms of

$$T_{ab}$$

Example Cosmology: (homogeneous & isotropic)
perfect fluid

of pressure p and density ρ modelled by

$$T^{ab} = (\rho + p)u^a u^b - pg^{ab}$$

radiative fluid
What is a fluid of photons:

$$T_{\text{Maxwell}}^{ab}g_{ab} = 0$$

observe: $T_{\text{p.f.}}^{ab}g_{ab} \stackrel{!}{=} 0$
 $= (\rho + p)u^a u^b g_{ab} - p \underbrace{g^{ab}g_{ab}}_4$

$$\leftrightarrow \rho_p 0 4 p = 0$$

$$\rho = 3p$$

$p = \frac{1}{3}\rho$
Reconvene at 3 pm? (EY : 20150409 I sent a Facebook (FB) message to the International Winter School on Gravity and Light: there was no missing video; it continues on Lecture 15 immediately)

Tutorial 14: Relativistic Spacetime, Matter and Gravitation. Exercise 2: Lorentz force law.

Question electromagnetic potential.

27. LECTURE 15: EINSTEIN GRAVITY

Recall that in Newtonian spacetime, we were able to reformulate the Poisson law $\Delta\phi = 4\pi G_N\rho$ in terms of the Newtonian spacetime curvature as

$$R_{00} = 4\pi G_N\rho$$

R_{00} with respect to ∇_{Newton}

G_N = Newtonian gravitational constant

This prompted Einstein to postulate < 1915 that the relativistic field equations for the Lorentzian metric g of (relativistic) spacetime

$$R_{ab} = 8\pi G_N T_{ab} \text{✓}$$

However, this equation suffers from a problem

LHS: $(\nabla_a R)^{ab} \neq 0$

generically

RHS:

$$(\nabla_a T)^{ab} = 0$$

thought bubble: = formulated from an action

Einstein tried to argue this problem away.

Nevertheless, the equations cannot be upheld.

27.1. **Hilbert.** Hilbert was a specialist for variational principles.

To find the appropriate left hand side of the gravitational field equations, Hibert suggested to start from an action

$$S_{\text{Hilbert}}[g] = \int_M \sqrt{-g} R_{ab} g^{ab}$$

thought bubble = “simplest action”

aim: varying this w.r.t. metric g_{ab} will result in some tensor

$$G^{ab} = 0$$

27.2. **Variation of S_{Hilbert} .**

$$0 \stackrel{!}{=} \underbrace{\delta}_{g_i} S_{\text{Hilbert}}[g] = \int_M \underbrace{[\delta\sqrt{-g}g^{ab}R_{ab}]}_1 + \underbrace{\sqrt{-g}\delta g^{ab}R_{ab}}_2 + \underbrace{\sqrt{-g}g^{ab}\delta R_{ab}}_3$$

$$\text{and 1 : } \delta\sqrt{-g} = \frac{-(\det g)g^{mn}\delta g_{mn}}{2\sqrt{-g}} = \frac{1}{2}\sqrt{-g}g^{mn}\delta g_{mn}$$

thought bubble

$$\delta\det(g) = \det(g)g^{mn}\delta g_{mn}$$

e.g. from

$$\det(g) = \exp \text{tr} \ln g$$

ad 2: $g^{ab}g_{bc} = \delta_c^a$

$$\implies (\delta g^{ab})g_{bc} + g^{ab}(\delta g_{bc}) = 0$$

$$\implies \delta g^{ab} = -g^{am}g^{bn}\delta g_{mn}$$

ad 3:

$$\begin{aligned} \Delta R_{ab} &\stackrel{\text{normal coords at point}}{=} \delta\partial_b\Gamma^m_{am} - \delta\partial_m\Gamma^m_{ab} + \Gamma\Gamma - \Gamma\Gamma = \\ &= \partial_b\delta\Gamma^m_{am} - \partial_m\delta\Gamma^m_{ab} = \\ &= \nabla_b(\delta\Gamma^m_{am}) - \nabla_m(\delta\Gamma^m_{ab}) \\ &\implies \sqrt{-g}g^{ab}\delta R_{ab} = \sqrt{-g} \end{aligned}$$

“if you formulate the variation properly, you’ll see the variation δ commute with ∂_b ” EY : 20150408 I think one uses the integration at the bounds, integration by parts trick

$\Gamma^i_{(x)jk} - \tilde{\Gamma}^i_{(x)jk}$ are the components of a (1,2)-tensor.

Notation: $(\nabla_b A)^i{}_g =: A^i{}_{j;b}$

$$\begin{aligned} &\implies \sqrt{-g}g^{ab}\delta R_{ab} \\ &\stackrel{\nabla g=0}{=} \sqrt{-g}(g^{ab}\delta\Gamma^m_{am})_{;b} - \sqrt{-g}(g^{ab}\delta\Gamma^m_{ab})_{;m} = \sqrt{-g}A^b{}_{;b} - \sqrt{-g}B^m{}_{;m} \end{aligned}$$

Question: Why is the difference of coefficients a tensor?

Answer:

$$\Gamma^i_{(y)jk} = \frac{\partial y^i}{\partial x^m} \frac{\partial x^m}{\partial y^j} \frac{\partial x^q}{\partial y^k} \Gamma^m_{(x) ,nq} + \frac{\partial y^i}{\partial x^m} \frac{\partial^2 x^m}{\partial y^j \partial y^k}$$

Collecting terms, one obtains

$$\begin{aligned} 0 &\stackrel{!}{=} \delta S_{\text{Hilbert}} = \int_M \left[\frac{1}{2} \sqrt{-g} g^{mn} \delta g_{mn} g^{ab} R_{ab} - \sqrt{-g} g^{am} g^{bn} \delta g_{mn} R_{ab} + \underbrace{(\sqrt{-g} A^a)_{,a}}_{\text{surface}} - \underbrace{(\sqrt{-g} B^b)_{,b}}_{\text{surface term}} \right] \\ &= \int_M \sqrt{-g} \delta \underbrace{g_{mn}}_{\text{arbitrary variation}} \left[\frac{1}{2} g^{mn} R - R^{mn} \right] \implies G^{mn} = R^{mn} - \frac{1}{2} g^{mn} R \end{aligned}$$

Hence Hilbert, from this “mathematical” argument, concluded that one may take

$$R_{ab} - \frac{1}{2} g_{ab} R = 8\pi G_N T_{ab}$$

Einstein equations

$$S_{E-H}[g] = \int_M \sqrt{-g} R$$

27.3. **3. Solution of the $\nabla_a T^{ab} = 0$ issue.** One can show (\rightarrow Tutorials) that the Einstein curvature

$$G_{ab} = R_{ab} - \frac{1}{2} g_{ab} R$$

satisfy the so-called contracted differential Bianchi identity

$$(\nabla_a G)^{ab} = 0$$

27.4. **Variants of the field equations.**

(a) a simple rewriting:

$$R_{ab} - \frac{1}{2} g_{ab} R = 8\pi G_N T_{ab} = T_{ab}$$

$$G_N = \frac{1}{8\pi}$$

Contract on both sides g^{ab}

$$R_{ab} - \frac{1}{2} g_{ab} R = T_{ab} || g^{ab}$$

$$R - 2R = T := T_{ab} g^{ab}$$

$$\implies R = -T$$

$$\implies R_{ab} + \frac{1}{2} g_{ab} T = T_{ab}$$

$$\iff R_{ab} = (T_{ab} - \frac{1}{2} T g_{ab}) =: \hat{T}_{ab}$$

(b)

$$R_{ab} = \widehat{T}_{ab}$$

$$S_{E-H}[g] := \int_M \sqrt{-g}(R + 2\Lambda)$$

thought bubble: Λ cosmological constant

History:

1915: $\Lambda < 0$ (Einstein) in order to get a non-expanding universe

>1915: $\Lambda = 0$ Hubble

today $\Lambda > 0$ to account for an accelerated expansion

$\Lambda \neq 0$ can be interpreted as a contribution

$-\frac{1}{2}\Lambda g$ to the energy-momentum

“dark energy”

Question: surface terms scalar?

Answer: for a careful treatment of the surface terms which we discarded, see, e.g. E. Poisson, “A relativist’s toolkit”

C.U.P. “excellent book”

Question: What is a constant on a manifold?

Answer: $\int \sqrt{-g}\Lambda = \Lambda \int \sqrt{-g}1$

[back to dark energy]

[Weinberg, QCD, calculated]

idea: 1 could arise as the vacuum energy of the standard model fields

$\Lambda_{\text{calculated}} = 10^{120} \times \Lambda_{\text{obs}}$

“worst prediction of physics”

Tutorials: check that

- Schwarzschild metric (1916)
- FRW metric
- pp-wave metric
- Reisner-Nordstrom

\implies are solutions to Einstein’s equations

in high school

$$m\ddot{x} + m\omega^2 x^2 = 0$$

$$x(t) = \cos(\omega t)$$

ET: [elementary tutorials]

study motion of particles & observers in Schwarzschild S.T.

Satellite: Marcus C. Werner

Gravitational lensing

odd number of pictures Morse theory (EY:20150408 Morse Theory !!!)

Domenico Giulini

Hamiltonian form Canonical Formulations

Key to Quantum Gravity

TUTORIAL 13 SCHWARZSCHILD SPACETIME

EY : 20150408 I’m not sure which tutorial follows which lecture at this point.

The tutorial video is excellent itself. Here, I want to encourage the use of CAS to do calculations. There are many out there.

Again, I’m partial to the Sage Manifolds package for Sage Math which are both open-source and based on Python. I’ll use that here.

Exercise 1. Geodesics in a Schwarzschild spacetime

Question Write down the Lagrangian.

Load “Schwarzschild.sage” in Sage Math, which will always be available freely here <https://github.com/ernestyalumni/diffgeo-by-sagemnfd/blob/master/Schwarzschild.sage>:

```
sage: load("Schwarzschild.sage")
4-dimensional manifold 'M'
open subset 'U_sph' of the 4-dimensional manifold 'M'
Levi-Civita connection 'nabla_g' associated with the Lorentzian metric 'g' on the 4-dimensional manifold 'M'
```

and so on.

Look at the code and I had defined the Lagrangian to be

L

. To get out the coefficients of L of the components of the tangent vectors to the curve, i.e. t', r', θ', ϕ' , denoted

tp, rp, thp, php

in my .sage file, do the following:

```
sage: L.expr().coefficients(tp)[1][0].factor().full_simplify()
(2*G_N*M_0 - r)/r
sage: L.expr().coefficients(rp)[1][0].factor().full_simplify()
-r/(2*G_N*M_0 - r)
sage: L.expr().coefficients(php)[1][0].factor().full_simplify()
r^2
sage: L.expr().coefficients(thp)[1][0].factor().full_simplify()
r^2*sin(th)^2
```

Question There are 4 Euler-Lagrange equations for this Lagrangian. Derive the one with respect to the function $t(\lambda)!$.

```
sage: L.expr().diff(t)
0
```

This confirms that $\frac{\partial L}{\partial t} = 0$

For $\frac{d}{d\lambda} \frac{\partial L}{\partial t'}$, then one needs to consider this particular workaround for Sage Math (computer technicality). One takes derivatives with respect to declared variables (declared with var) and then substitute in functions that are dependent upon λ , and then take the derivative with respect to the parameter λ . This does that:

```
sage: L.expr().diff( thp ).factor().subs( r == gamma1 ).subs( thp == gamma3.diff( tau ) ).subs( th == gamma3 ).diff(tau)\
....: .factor()
2*(2*cos(gamma3(tau))*gamma1(tau)*D[0](gamma3)(tau)^2 + 2*sin(gamma3(tau))*D[0](gamma1)(tau)*D[0](gamma3)(tau)
+ gamma1(tau)*sin(gamma3(tau))*D[0, 0](gamma3)(tau))*gamma1(tau)*sin(gamma3(tau))
```

Question Show that the Lie derivative of g with respect to the vector fields $K_t := \frac{\partial}{\partial t}$.

The first line defines the vector field by accessing the frame defined on a chart with spherical coordinates and getting the time vector. The second line is the Lie derivative of g with respect to this vector field.

```
sage: K_t = espher[0]
sage: g.lie_der(K_t).display() # 0, as desired
0
```

EY : 20150410 My question is this: $\forall X \in \Gamma(TM)$ i.e. X is a vector field on M , or, specifically, a section of the tangent bundle, then does

$$\mathcal{L}_X g = 0$$

instantly mean that X is a symmetry for (M, g) ? $\mathcal{L}_X g$ is interpreted geometrically as how g changes along the flow generated by X , and if it equals 0, then g doesn’t change.

28.

29.

30. CANONICAL FORMULATION OF GR I

Dynamical and Hailtonian formulation of General Relativity.

Purpose

- (1) formulate and solve initial-value problems
- (2) integrate Einstein’s Equations by numerical codes
- (3) characterize degrees of freedom
- (4) characterize isolated systems, associated symmetry groups and conserved quantities, like Energy/Mass, Momenta (linear and angular), Poincaré charges
- (5) starting point for “canonical quantization” program.

How. We will rewrite Einstein’s Eq. in form of a *constrained Hamiltonian system*.

(− + ++)

$$T^{\mu\nu} = \begin{pmatrix} W & \frac{1}{c}S^m \\ cg^m & \mathfrak{t}^{mn} \end{pmatrix}$$

- W = Energy density (1 component)
- g^m = Momentum density, (3 components)
- S^m = Energy current-density (3 components)
- \mathfrak{t}^{mn} = Momentum current-density (6 components)

$$T^{\mu\nu} = T^{\nu\mu} \implies S^m = c^2 g^m$$

10 independent komp. (components)

Phys. dim. $[T^{\mu\nu}] = \frac{J}{m^3}$
 $[G^{\mu\nu}] = \frac{1}{m^2}$

$$k = \frac{\text{curvature}}{\text{Energy} \cdot \text{density}}$$

$$[k] = \frac{1}{m^2} / \frac{J}{m^3}, \quad {}^2k = \frac{\text{Curvature}}{\text{mass density}} = \left(\frac{1}{1.5 \text{ AU}}\right)^2 / \text{Density of water}$$
$$= \left(\frac{1}{10 \text{ km}}\right)^2 / \text{Nuclear density in core of neutron star} \simeq 5 \cdot 10^{17} \text{ kg}/m^3$$

If “Ein” for Einstein Tensor, $G_{\mu\nu} = \text{Ein}\left(\frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu}\right)$

$$\text{Ein}(v, w) = \frac{1}{4}[\text{Ein}(v + w, v + w) - \text{Ein}(v - w, v - w)]$$
$$\text{Ein}(w, w) = -g(w, w) \sum_{\perp w} \text{Sec}$$

where $\perp w$ is take the sum over any triple of mutually perp. 2-planes in $\perp w$

$\text{Sec}(\text{Span}\{v, w\}) = \frac{\text{Riem}(v, w, v, w)}{[g(v, w)]^2 - g(v, v)g(w, w)}$
“sectional curvature”

Identity: $\nabla_\mu G^{\mu\nu} = 0$ (follows from twice-contracted II. Bianchi Identity
 $\sum_{\lambda\mu\nu} \text{cycl} \nabla_\lambda R_{\alpha\beta\mu\nu} = 0$)

$$\underbrace{\partial_0 G^{0\nu}}_{\text{contains at most 1st time der.}} + \underbrace{\partial_k G^{k\nu} + \Gamma G + \Gamma G}_{\text{contains at most 2nd. time derivatives}} \equiv 0$$

\implies 4 out of 10 Einstein Eq. do not evolve the fields but rather constrain the initial data. The space-space components (6 Eqs.) are the evolution Eqs.

- 10 Einstein Eq. - 4 constraints (underdetermined elliptic type)
- \ - 6 evolution equations (undetermined hyperbolic type)

31.

32.

33.

34. LECTURE 22: BLACK HOLES

Only depends on Lectures 1-15, so does lecture on “Wednesday”
Schwarzschild solution also vacuum solution (from tutorial EY : oh no, must do tutorial)
Study the Schwarzschild as a vacuum solution of the Einstein equation:
 $m = G_N M$ where M is the “mass”

$$g = \left(1 - \frac{2m}{r}\right) dt \otimes dt - \frac{1}{1 - \frac{2m}{r}} dr \otimes dr - r^2(d\theta \otimes d\theta + \sin^2 \theta d\varphi \otimes d\varphi)$$

in the so-called Schwarzschild coordinates

t	r	θ	φ
$(-\infty, \infty)$	$(0, \infty)$	$(0, \pi)$	$(0, 2\pi)$

What staring at this metric for a while, two questions naturally pose themselves:

(i) What exactly happens $r = 2m$?

t	r	θ	φ
$(-\infty, \infty)$	$(0, 2m) \cup (2m, \infty)$	$(0, \pi)$	$(0, 2\pi)$

(ii) Is there anything (in the real world) beyond $t \rightarrow -\infty$?

$$t \rightarrow +\infty$$

idea: Map of Linz, blown up
Insight into these two issues is afforded by stopping to stare.
Look at *geodesic* of g , instead.

34.1. **Radial null geodesics.** null - $g(v_\gamma, v_\gamma) = 0$
Consider null geodesic in “Schd”

$$S[\gamma] = \int d\lambda \left[\left(1 - \frac{2m}{r}\right) \dot{t}^2 - \left(1 - \frac{2m}{r}\right)^{-1} \dot{r}^2 - r^2(\dot{\theta}^2 + \sin^2 \theta \dot{\varphi}^2) \right]$$

with $[\dots] = 0$
and one has, in particular, the t -eqn. of motion:

$$\left(\left(1 - \frac{2m}{r}\right) \dot{t}\right)' = 0$$

⇒

$$\left(1 - \frac{2m}{r}\right) \dot{t} = k = \text{const.}$$

Consider radial null geodesics

$$\theta \stackrel{!}{=} \text{const.} \qquad \varphi = \text{const.}$$

From \square and \square

$$\implies \dot{r}^2 = k^2 \leftrightarrow \dot{r} = \pm k$$

$$\implies r(\lambda) = \pm k \cdot \lambda$$

Hence, we may consider

$$\tilde{t}(r) := t(\pm k \lambda)$$

Case A: \oplus

$$\frac{d\tilde{t}}{dr} = \frac{\tilde{\dot{t}}}{\dot{r}} = \frac{k}{\left(1 - \frac{2m}{r}\right)k} = \frac{r}{r - 2m}$$

$$\implies \tilde{t}_+(r) = r + 2m \ln|r - 2m|$$

(**outgoing** null geodesics)

Case b. \pm (Circle around $-$, consider $-$):

$$\tilde{t}_-(r) = -r - 2m \ln|r - 2m|$$

(**ingoing** null geodesics)

Picture

34.2. **Eddington-Finkelstein.** Brilliantly simple idea:

change (on the domain of the Schwarzschild coordinates) to different coordinates, s.t.

in those new coordinates,

ingoing null geodesics appear as straight lines, of slope -1

This is achieved by

$$\bar{t}(t, r, \theta, \varphi) := t + 2m \ln|r - 2m|$$

Recall: ingoing null geodesic has

$$\tilde{t}(r) = -(r + 2m \ln|r - 2m|) \qquad (Schdcoords)$$

$$\iff \bar{t} - 2m \ln|r - 2m| = -r - 2m \ln|r - 2m| + \text{const.}$$

$$\therefore \bar{t} = -r + \text{const.}$$

(Picture)

outgoing null geodesics

$$\bar{t} = r + 4m \ln|r - 2m| + \text{const.}$$

Consider the new chart (V, g) while (U, x) was the Schd chart.

$$\bigcup_{\text{Schd}} \{ \text{horizon} \} = V$$

“chart image of the horizon”

Now calculate the *Schd metric* g w.r.t. Eddington-Finkelstein coords.

$$\bar{t}(t, r, \theta, \varphi) = t + 2m \ln|r - 2m|$$

$$\bar{r}(t, r, \theta, \varphi) = r$$

$$\bar{\theta}(t, r, \theta, \varphi) = \theta$$

$$\bar{\varphi}(t, r, \theta, \varphi) = \varphi$$

EY : 20150422 I would suggest that after seeing this, one would calculate the metric by your favorite CAS. I like the Sage Manifolds package for Sage Math.

[Schwarzschild_BH.sage on github](#)

[Schwarzschild_BH.sage on Patreon](#)

[Schwarzschild_BH.sage on Google Drive](#)

```
sage: load('Schwarzschild_BH.sage')
4-dimensional manifold 'M'
expr = expr.simplify_radical()
Levi-Civita connection 'nabla_g' associated with the Lorentzian metric 'g' on the 4-dimensional manifold 'M'
Launched png viewer for Graphics object consisting of 4 graphics primitives
```

Then calculate the Schwarzschild metric g but in Eddington-Finkelstein coordinates. Keep in mind to calculate the set of coordinates that uses \bar{t} , not \tilde{t} :

```
sage: gI.display()
gI = (2*m - r)/r dt*dt - r/(2*m - r) dr*dr + r^2 dth*dth + r^2*sin(th)^2 dph*dph
sage: gI.display( X_EF_I_null.frame())
gI = (2*m - r)/r dtbar*dtbar + 2*m/r dtbar*dr + 2*m/r dr*dtbar + (2*m + r)/r dr*dr + r^2 dth*dth + r^2*sin(th)^2 dph*dph
```

REFERENCES

[1] William Feller. **An Introduction to Probability Theory and Its Applications**, Vol. 1, 3rd Edition. 1968. ISBN-13: 978-0471257080

[2] Jeffrey M. Lee. **Manifolds and Differential Geometry**, *Graduate Studies in Mathematics* Volume: 107, American Mathematical Society, 2009. ISBN-13: 978-0-8218-4815-9

[3] John Lee, **Introduction to Smooth Manifolds** (Graduate Texts in Mathematics, Vol. 218), 2nd edition, Springer, 2012, ISBN-13: 978-1441999818

[4] Victor Guillemin, Alan Pollack. **Differential Topology**. American Mathematical Society. 2010. ISBN-13: 978-0821851937 https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=2&cad=rja&uact=8&ved=0ahUKEwjG96q9z63JAhWMLYgKHempDoMQFggmMAE&url=http%3A%2F%2Fwww.mat.unimi.it%2Fusers%2Fdedo%2Ftop%2520diff%2FGuillemin-Pollack_Differential%2520topology.pdf&usg=AFQjCNF5imOH5xeXRSK60qzM7zT97sdIsw

[5] Anant R. Shastri. **Elements of Differential Topology**. CRC Press. 2011. ISBN-13: 978-0415339209

[6] Morris W. Hirsch, **Differential Topology** (Graduate Texts in Mathematics), Graduate Texts in Mathematics (Book 33), Springer (September 16, 1997). ISBN-13: 978-0387901480

[7] P.-A. Absil, R. Mahony, and R. Sepulchre. **Optimization Algorithms on Matrix Manifolds**. Princeton University Press. 2008. ISBN 978-0-691-13298-3 <https://press.princeton.edu/titles/8586.html>

[8] John Baez, Javier P Muniain. **Gauge Fields, Knots and Gravity** (Series on Knots and Everything), World Scientific Publishing Company (October 24, 1994), ISBN-13: 978-9810220341

[9] Joseph J. Rotman, **Advanced Modern Algebra** (Graduate Studies in Mathematics), (Book 114), American Mathematical Society; 2 edition, (August 10, 2010), ISBN-13: 978-0821847411

[10] Andrew Baker, “Representations of Finite Groups”, 07112011 <http://www.maths.gla.ac.uk/~ajb/dvi-ps/groupreps.pdf>

[11] Yvette Kosmann-Schwarzbach, **Groups and Symmetries: From Finite Groups to Lie Groups**, Springer, 2010. e-ISBN 978-0-387-78866-1 <http://www.caam.rice.edu/~yad1/miscellaneous/References/Math/Groups/Groups%20and%20Symmetries.pdf>

[12] Agostino Prástaro. **Geometry of PDEs and Mechanics**. World Scientific Publishing Co. 1996. QC125.2.P73 1996 530.1'55353–dc20.
Agostino Prástaro. *Dipartimento di Metodi e Modelli Matematici per le Scienze Applicate, Università degli Studi di Roma “La Sapienza”*.

[13] Clemens Koppensteiner. *Complex Manifolds: Lecture Notes*. <http://www.caramdir.at/uploads/math/piii-cm/complex-manifolds.pdf>

[14] Stefan Vandoren. **Lectures on Riemannian Geometry, Part II: Complex Manifolds**. <http://www.staff.science.uu.nl/~vando101/MRIlectures.pdf>

[15] Kentaro Hori (Author, Editor), Sheldon Katz (Editor), Albrecht Klemm (Editor), Rahul Pandharipande (Editor), Richard Thomas (Editor), Cumrun Vafa (Editor), Ravi Vakil (Editor), Eric Zaslow (Editor). **Mirror Symmetry** (Clay Mathematics Monographs, V. 1). Clay Mathematics Monographs (Book 1). American Mathematical Society (August 19, 2003). ISBN-10: 0821829556 ISBN-13: 978-0821829554 <https://web.archive.org/web/20060919020706/http://math.stanford.edu/~vakil/files/mirrorfinal.pdf>

[16] Lawrence Conlon. **Differentiable Manifolds** (Modern Birkhäuser Classics). 2nd Edition. Birkhäuser; 2nd edition (October 10, 2008). ISBN-13: 978-0817647667

[17] Andrew Clarke and Bianca Santoro. *Holonomy Groups in Riemannian Geometry*. [arXiv:1206.3170](https://arxiv.org/abs/1206.3170) [math.DG]

[18] Urs Schreiber and Konrad Waldorf. *Parallel Transport and Functors*. [arXiv:0705.0452](https://arxiv.org/abs/0705.0452) [math.DG]

[19] Y. Eliashberg, N. Mishachev, *Introduction to the h-principle*, Grad. Studies in Math. Vol 48 (AMS 2002)

[20] Walter Greiner. **Classical Mechanics: Point Particles and Relativity** (Classical Theoretical Physics). Springer New York; 2004 edition (December 4, 2003). ISBN-13: 978-0387955865

[21] Vladimir I. Arnold, Valery V. Kozlov, Anatoly I. Neishtadt. **Mathematical Aspects of Classical and Celestial Mechanics**. Third Edition. Springer.

[22] Yvonne Choquet-Bruhat. **General Relativity and the Einstein Equations** (Oxford Mathematical Monographs) 1st Edition. Oxford Mathematical Monographs (February 4, 2009) ISBN-13: 978-0199230723

[23] T. Frankel, **The Geometry of Physics**, Cambridge University Press, Third Edition, 2012.

[24] Shigeyuki Morita. **Geometry of Differential Forms** (Translations of Mathematical Monographs, Vol. 201). American Mathematical Society (August 28, 2001). ISBN-10: 0821810456

[25] Tom M. Apostol. **Calculus, Vol. 1: One-Variable Calculus, with an Introduction to Linear Algebra**. Wiley; 2nd edition (January 16, 1991).ISBN-13: 978-0471000051

[26] Roger R. Bate, Donald D. Mueller, Jerry E. White. **Fundamentals of Astrodynamics** (Dover Books on Aeronautical Engineering). Publisher: Dover Publications; Revised ed. edition (June 1, 1971). ISBN-13: 978-0486600611