

# GE23131-Programming Using C-2024

Quiz navigation

1

2

3

Show one page at a time

Finish review

Status	Finished
Started	Friday, 17 January 2025, 2:05 PM
Completed	Friday, 17 January 2025, 2:22 PM
Duration	17 mins 3 secs

Question 1

Correct

Marked out of 3.00

Flag question

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that  $A[i] - A[j] = k, i \neq j$ .

Input Format

- 1. First line is number of test cases T. Following T lines contain:
- 2. N, followed by N integers of the array
- 3. The non-negative integer k

Output format

Print 1 if such a pair exists and 0 if it doesn't.

Example

Input:

3 1 3 5

4

Output:

1

Input:

1

3 1 3 5

99

Output:

0

**Answer:** (penalty regime: 0 %)

```
1 #include<stdio.h>
2 int main(){
3     int t;
4     scanf("%d",&t);
5     while(t--){
6         int n;
7         scanf("%d",&n);
8         int a[n];
9         for(int i=0;i<n;i++){
10             scanf("%d",&a[i]);
11         }
```


```
14 scanf("%d",&k);
15 int flag=0;
16 for(int i=0;i<n;i++){
17     for(int j=i+1;j<n;j++){
18         if((a[i]-a[j]==k) || (a[j]-a[i]==k)){
19             flag=1;
20             break;
21         }
22     }
23     if(flag) break;
24 }
25 printf("%d\n",flag);
26 }
27 }
```

	Input	Expected	Got	
✓	1 3 1 3 5 4	1	1	✓
✓	1 3 1 3 5 99	0	0	✓

Passed all tests! ✓

Sam loves chocolates and starts buying them on the 1st day of the year. Each day of the year, x, is numbered from 1 to Y. On days when x is odd, Sam will buy x chocolates; on days when x is even, Sam will not purchase any chocolates.

Marked out of  
5.00

 [Flag question](#)

new line. This is a function-only challenge, so input is handled for you by the locked stub code in the editor.

### Input Format

The program takes an array of integers as a parameter.

The locked code in the editor handles reading the following input from stdin, assembling it into an array of integers (arr), and calling calculate(arr).

The first line of input contains an integer, T (the number of test cases). Each line i of the T subsequent lines describes the ith test case as an integer, Ni (the number of days).

### Constraints

$$1 \leq T \leq 2 \times 10^5$$

$$1 \leq N \leq 2 \times 10^6$$

$$1 \leq x \leq N \leq Y$$

### Output Format

For each test case, Ti in arr, your calculate method should print the total number of chocolates Sam purchased by day Ni on a new line.

### Sample Input 0

2

3

Sample Output 0

1

1

4

Explanation

Test Case 0:  $N = 1$

Sam buys 1 chocolate on day 1, giving us a total of 1 chocolate. Thus, we print 1 on a new line.

Test Case 1:  $N = 2$

Sam buys 1 chocolate on day 1 and 0 on day 2. This gives us a total of 1 chocolate. Thus, we print 1 on a new line.

Test Case 2:  $N = 3$

Sam buys 1 chocolate on day 1, 0 on day 2, and 3 on day 3. This gives us a total of 4 chocolates. Thus, we print 4 on a new line.

**Answer:** (penalty regime: 0 %)

```
4 scanf("%d",&t);
5 while(t--){
6     int n,c=0;
7     scanf("%d",&n);
8     for(int i=0;i<=n;i++){
9         if(i%2!=0) c=c+i;
10    }printf("%d\n",c);
11 }
12 }
```

	Input	Expected	Got	
✓	3	1	1	✓
	1	1	1	
	2	4	4	
	3			

100	1849	1849
86	729	729
54	400	400
40	25	25
9	1521	1521
77	25	25
9	49	49
13	2401	2401
98		

Passed all tests! ✓

Question **3**

Correct

Marked out of  
7.00

Flag question

The number of goals achieved by two football teams in matches in a league is given in the form of two lists. Consider:

- Football team A, has played three matches, and has scored { 1 , 2 , 3 } goals in each match respectively.
- Football team B, has played two matches, and has scored { 2, 4 } goals in each match respectively.
- Your task is to compute, for each match of team B, the total number of matches of team A, where team A has scored less than or equal to the number of goals scored by team B in that match.
- In the above case:
- For 2 goals scored by team B in its first match, team A has 2 matches with scores 1 and 2.
- For 4 goals scored by team B in its second match, team A has 3 matches with scores 1, 2 and 3.

Hence, the answer: {2, 3}.

positive integers, one for each  $\text{maxes}[i]$  representing the total number of elements  $\text{nums}[j]$  satisfying  $\text{nums}[j] \leq \text{maxes}[i]$  where  $0 \leq j < n$  and  $0 \leq i < m$ , in the given order.

It has the following:

$\text{nums}[\text{nums}[0], \dots, \text{nums}[n-1]]$ : first array of positive integers

$\text{maxes}[\text{maxes}[0], \dots, \text{maxes}[m-1]]$ : second array of positive integers

Constraints

- $2 \leq n, m \leq 105$
- $1 \leq \text{nums}[j] \leq 109$ , where  $0 \leq j < n$ .
- $1 \leq \text{maxes}[i] \leq 109$ , where  $0 \leq i < m$ .

Input Format For Custom Testing

Input from stdin will be processed as follows and passed to the function.

The first line contains an integer  $n$ , the number of elements in  $\text{nums}$ .

The next  $n$  lines each contain an integer describing  $\text{nums}[j]$  where  $0 \leq j < n$ .

The next line contains an integer  $m$ , the number of elements in  $\text{maxes}$ .

The next  $m$  lines each contain an integer describing  $\text{maxes}[i]$  where  $0 \leq i < m$ .

Sample Case 0



4

1

4

2

4

2

3

5

Sample Output 0

2

4

Explanation 0

We are given  $n = 4$ ,  $\text{nums} = [1, 4, 2, 4]$ ,  $m = 2$ , and  $\text{maxes} = [3, 5]$ .

1. For  $\text{maxes}[0] = 3$ , we have 2 elements in  $\text{nums}$  ( $\text{nums}[0] = 1$  and  $\text{nums}[2] = 2$ ) that are  $\leq \text{maxes}[0]$ .
2. For  $\text{maxes}[1] = 5$ , we have 4 elements in  $\text{nums}$  ( $\text{nums}[0] = 1$ ,  $\text{nums}[1] = 4$ ,  $\text{nums}[2] = 2$ , and  $\text{nums}[3] = 4$ ) that are  $\leq \text{maxes}[1]$ .

Thus, the function returns the array  $[2, 4]$  as the answer.

Sample Case 1

5  
2  
10  
5  
4  
8  
4  
3  
1  
7  
8

Sample Output 1

1  
0  
3  
4

Explanation 1

We are given,  $n = 5$ ,  $nums = [2, 10, 5, 4, 8]$ ,  $m = 4$ , and  $maxes = [3, 1, 7, 8]$ .

1. For  $maxes[0] = 3$ , we have 1 element in  $nums$  ( $nums[0] = 2$ ) that is  $\leq maxes[0]$ .

nums[3] = 4) that are  $\leq$  maxes[2].

4. For maxes[3] = 8, we have 4 elements in nums (nums[0] = 2, nums[2] = 5, nums[3] = 4, and nums[4] = 8) that are  $\leq$  maxes[3].

Thus, the function returns the array [1, 0, 3, 4] as the answer.

**Answer:** (penalty regime: 0 %)

```

1  #include<stdio.h>
2  int main()
3  {
4      int s1,s2,ans;
5      scanf("%d",&s1);
6      int ta[s1];
7      for(int i=0;i<s1;i++)
8          scanf("%d",&ta[i]);
9      scanf("%d",&s2);
10     int tb[s2];
11     for(int i=0;i<s2;i++)
12         scanf("%d",&tb[i]);
13     for(int j=0;j<s2;j++)
14     {
15         ans=0;
16         for(int i=0;i<s1;i++){
17             if(tb[j]>=ta[i])
18                 ans++;
19         }printf("%d\n",ans);
20     }
21 }
```

	Input	Expected	Got	

	4			
	2			
	4			
	2			
	3			
	5			
✓	5	1	1	✓
	2	0	0	
	10	3	3	
	5	4	4	
	4			
	8			
	4			
	3			
	1			
	7			
	8			

Passed all tests! ✓

Finish review