

# Mid-Term Exam

Subject:- Chemical Engineering  
Thermodynamics-I

Submitted by:- Muhammad Yasir

Registration NO:- CHEN19111008

Submitted To:- Engr. Dr. Aamir Alaud Din

Class:- Bs Chemical Engineering

Date:- 27-November-2020

KFUEIT RYK

Muhammad Yasir

chen19111008

## — (Q.NO.1) —

### 1. System:-

Refer to the subject matter of analysis. Thermodynamic system or system refers to definite quantity of matter, enclosed by a boundary on which we focus our attention for thermodynamic analysis.

e.g. The gas in the container is system.

### 2. Surroundings:-

we define surroundings as. The part of the universe other than the system. It interacts with the system.

e.g. The walls of the container are part of the surroundings.

### 3. Adiabatic process:-

In the adiabatic process, no heat passes through boundaries of the system.

Muhammad Yasir

chen19111008

#### 4. Isolated system:-

A system in which both mass and energy cannot flow across the system boundary.

#### 5. Extensive Property:-

The property of the system which depends upon the mass of the system.

We will use capital letter for the extensive property.

e.g Mass, Volume, Internal energy etc.

Muhammad Yasir

chen19111008

Q. No. 2Given Data:-

$$p = 7 \text{ bar}$$

$$\hat{h} = 2600 \text{ KJ/kg}$$

Required:-

$$x = ?$$

$$\hat{V} = ?$$

$$\hat{u} = ?$$

Solution:-

Now, we find  $x$  as.

$$\hat{h} = hf + x hfg$$

putting values, we get.

$$2600 = 697.1 + x (2064.9)$$

$$2600 - 697.1 = x (2064.9)$$

$$1902.9 = x (2064.9)$$

$$x = \frac{1902.9}{2064.9}$$

$$x = 0.921$$



Muhammad Yasir

chen/9111008

Now,

$$\begin{aligned}
 \hat{u} &= u_f + x u_{fg} \\
 &= 696.3 + (0.921)(2571.1 - 696.3) \\
 &= 696.3 + (0.921)(1874.8) \\
 &= 696.3 + 1726.6908
 \end{aligned}$$

$$\boxed{\hat{u} = 2420 \text{ kJ/kg}}$$

also, for  $\hat{v}$ 

$$\begin{aligned}
 \hat{v} &= v_f + x v_{fg} \\
 &= 0.001108 + (0.921)(0.273 - 0.001108) \\
 &= 0.001108 + (0.921)(0.271892) \\
 &= 0.001108 + 0.25041
 \end{aligned}$$

$$\boxed{\hat{v} = 0.2515 \text{ m}^3/\text{kg}}$$

Muhammad Yasir chen19111008

— (Q. No. 3) —

Given Data:-

$$m = 10 \text{ Kg}$$

$$P_1 = 20 \text{ bar}$$

$$V_1 = 1.0 \text{ m}^3$$

$$P_2 = 100 \text{ bar}$$

Required:-

$$(a) \quad \dot{W} = ?$$

$$(b) \quad \dot{q} = ?$$

$$(a) \quad \dot{W} = ?$$

$$P_2 = \frac{100 \text{ bar} \mid 100 \text{ KPa}}{1 \text{ bar}} = 10,000 \text{ KPa} = 10 \text{ MPa}$$

$$\dot{V}_2 = ?$$

Since,

$$P_1 \dot{V}_1^{1.5} = P_2 \dot{V}_2^{1.5} \Rightarrow \dot{V}_2^{1.5} = \frac{P_1 \dot{V}_1^{1.5}}{P_2}$$

$$\dot{V}_2 = \left( \frac{P_1 \dot{V}_1^{1.5}}{P_2} \right)^{1/1.5}$$

$$= \left[ \frac{2 \text{ MPa} \mid \left( \frac{0.1 \text{ m}^3}{\text{kg}} \right)^{1.5} \mid 1}{10 \text{ MPa}} \right]^{1/1.5}$$

Muhammad Yasir

chen19111008

$$\begin{aligned}
 \dot{V}_2 &= \left( (0.2 \times 10.1)^{1.5} \right)^{1/1.5} \text{ m}^3/\text{Kg} \\
 &= (0.2 \times 0.0316)^{1/1.5} \text{ m}^3/\text{Kg} \\
 &= (0.2 \times 0.0316)^{0.667}
 \end{aligned}$$

$$\dot{V}_2 = 0.0342 \text{ m}^3/\text{Kg}$$

$$\dot{W} = \int_{0.1}^{0.0342} P E \cdot d\dot{V} = - \int_{0.1}^{0.0342} P \cdot d\dot{V}$$

$$P \dot{V}^{1.5} = P_1 \dot{V}_1^{1.5} \Rightarrow P = \frac{P_1 \dot{V}_1^{1.5}}{\dot{V}^{1.5}}$$

$$P = \frac{2 \text{ MPa} \left( \frac{0.1 \text{ m}^3}{\text{Kg}} \right)^{1.5}}{\left( \frac{\text{Kg}}{\dot{V} \text{ m}^3} \right)^{1.5}}$$

$$P = \frac{0.0632}{\dot{V}^{1.5}} \text{ MPa}$$

$$\dot{W} = - \int_{0.1}^{0.0342} \frac{0.0632}{\dot{V}^{1.5}} d\dot{V}$$

$$= \left. \frac{-0.0632}{-0.5} \frac{1}{\dot{V}^{0.5}} \right|_{0.1}^{0.0342}$$

$$= \frac{-0.0632}{-0.5} \left[ \frac{1}{(0.0342)^{0.5}} - \frac{1}{(0.1)^{0.5}} \right]$$

Muhammad Yasir

chen19111008

$$= \frac{0.2838 \text{ MPa} \cdot \text{m}^3}{\text{kg}} \left| \frac{1000 \text{ KPa}}{1 \text{ MPa}} \right| \left| \frac{1 \text{ kN}}{1 \text{ KPa} \cdot \text{m}^2} \right|$$

$$\hat{w} = 283.8 \text{ KJ/kg}$$

$$(b) \hat{q} = ?$$

since,

$$\Delta \hat{u} = \hat{q} + \hat{w}$$

At state,  $p_1 = 2 \text{ MPa}$ ,  $T_1 = 213.6^\circ \text{C}$ 

$$\hat{u}_1 = ?$$

$\hat{u}_1 \text{ (KJ/kg)}$	$T(^{\circ}\text{C})$
2600.3	212.4
$u_1$	213.6
2628.3	225

After Interpolation, we get

$$\hat{u}_1 = 2602.97 \text{ KJ/kg}$$



Muhammad Yasir chen19111008

At State 2,  $p_2 = 10 \text{ MPa}$ ,  $\hat{v}_2 = 0.0342 \text{ m}^3/\text{kg}$

$\hat{u}_2 \text{ (KJ/kg)}$	$\hat{v}_2 \text{ (m}^3/\text{kg)}$
3045.8	0.0328
$\hat{u}_2$	0.0342
3144.5	0.0356

After interpolation, we get

$$\boxed{\hat{u}_2 = 3095.15 \text{ KJ/kg}}$$

Since,

$$\begin{aligned} \Delta \hat{u} &= \hat{u}_2 - \hat{u}_1 \\ &= (3095.15 - 2602.97) \text{ KJ/kg} \end{aligned}$$

$$\boxed{\Delta \hat{u} = 492.18 \text{ KJ/kg}}$$

putting values, we get

$$\Delta \hat{u} = \hat{q}_1 + \hat{w}$$

$$\hat{q}_1 = \Delta \hat{u} - \hat{w}$$

$$\hat{q}_1 = (492.18 - 283.8) \text{ KJ/kg}$$

$$\boxed{\hat{q}_1 = 208.38 \text{ KJ/kg}}$$