

Babax Maqbool

CHE19111049

Chemical Engineering 3rd semester.

System:-

A "System" Everything under consideration. Everything is called universe. The part of universe in which we are interested called our system.

Surrounding:-

When we left the system from our universe the remaining universe is called surrounding. A boundary (Imaginary and real) separates the system and surrounding.

Adiabatic process:-

When at least one thermodynamic property of a system changes it means

Babax Maqbool

CHENIQUIN

BS chemical Eng


the state of a system change,
So, the system undergoes the
process.

In adiabatic process the heat
remain constant =

$Q = 0$ (when the system undergoes
from state 1 to state 2 Q remain
same).

Isolated System:-

A system in which no
mass or no energy inter the
system. Our system called
isolated.



Extensive property:-

An extensive property
is a physical quantity whose

CHE 111049
Babax Maqbool
BS chemical Eng

value is proportional to the size of the system it describes, or to the quantity of matter in the system. For example:- the mass of a sample is an extensive quantity. it depends on the amount of substance.

QNo: 2

Given data:

$$P = 7 \text{ bar}$$

$$h = 2600 \text{ kJ/kg} \quad E: 0.110$$

$$\hat{u} = ?$$

$$\hat{v} = ?$$

$$x = ?$$

$$\hat{h} = \hat{h}_f + x \hat{h}_{fg}$$

$$\frac{2600 \text{ kJ}}{\text{kg}} = \frac{697 \text{ kJ}}{\text{kg}} + x \frac{(2067) \text{ kJ}}{\text{kg}}$$

$$1903 \text{ kJ/kg}$$

$$x \text{ kg} \quad | \quad 2067 \text{ kJ/kg}$$

$$x = 0.92065$$

$$x = 0.921$$

$$\hat{u} = x \hat{u}_g$$

CHEN19111049

$$0.921 (0.2728) \text{ m}^3/\text{kg}$$

$$\hat{v} = 0.2511 \frac{\text{m}^3}{\text{kg}}$$

Now we calculate internal energy.

$$\hat{U} = \hat{U}_f + x \hat{U}_g$$

$$\hat{U} = \frac{696 \text{ kJ}}{\text{kg}} + 0.921 (2573 - 696) \frac{\text{kJ}}{\text{kg}}$$

$$\hat{U} = \frac{696 \text{ kJ}}{\text{kg}} + 0.921 (1877) \frac{\text{kJ}}{\text{kg}}$$

$$\hat{U} = \frac{696 \text{ kJ}}{\text{kg}} + 1728.717 \frac{\text{kJ}}{\text{kg}}$$

$$\hat{U} = 2424.717 \frac{\text{kJ}}{\text{kg}}$$

Q No: 3

$$m = 10.0 \text{ kg}$$

$$P_1 = 20 \text{ bar}$$

$$V_1 = 1.0 \text{ m}^3$$

$$P_2 = 100 \text{ bar}$$

$$P_v \text{ relation} = P v^5 = \text{const}$$

$$(1) \quad T = ?$$

$$P_1 = \frac{20 \text{ bar} \times 100 \text{ K}}{10 \text{ bar}} = 2000 \text{ KPa}$$

$$P_2 = 2 \text{ MPa}$$

$$\hat{v}_1 = \frac{1.0 \text{ m}^3}{10 \text{ kg}} = 0.1 \text{ m}^3/\text{kg}$$

CHEN19111049

$$\hat{w} = \int_{0.1}^{0.0342} P d\hat{v} = - \int_{0.1}^{0.0342} P d\hat{v}$$

$$P \hat{v}^{1.5} = P_1 \hat{v}_1^{1.5} \Rightarrow P = \frac{P_1 \hat{v}_1^{1.5}}{\hat{v}^{1.5}}$$

$$P = \left(2 \text{ MPa} \left(\frac{0.1 \text{ m}^3}{\text{kg}} \right)^{1.5} \right) \left(\frac{\text{kg}}{\text{m}^3} \right)^{1.5}$$

$$P = 0.0632 \text{ MPa}$$

$$\hat{w} = - \int_{0.1}^{0.0342} 0.0632 d\hat{v}$$

$$= -0.0632 \left[\frac{1}{-0.5} \right]_{0.1}^{0.0342}$$

$$= -0.0632 \left[\frac{1}{0.0342^{0.5}} - \frac{1}{0.1^{0.5}} \right]$$

$$= 0.2838 \text{ MPa m}^3 / \text{kg} \quad \left| \frac{1000 \text{ kPa}}{1 \text{ kg}} \right| \frac{1 \text{ kN}}{11 \text{ kPa} \cdot \text{m}^3}$$

$$= 283.8 \text{ kJ/kg}$$

$$\hat{w} = 283.8 \text{ kJ/kg}$$

(c) $q = ?$

$$\Delta \hat{u} = \hat{q} + \hat{w}$$

State 1, $P_1 = 2 \text{ MPa}$, $T = 213.6^\circ\text{C}$.

$u_1 = ?$

u_1 (kJ/kg) T (°C)

2600.3 212.4

u_1 213.6

CHEM19111049

From steam tables, at $P = \text{MPa}$

$$\hat{v}_1 = 0.0012, \hat{v}_g = 0.0996$$

Since, $\hat{v}_1 > \hat{v}_g$ at $P = \text{MPa}$,
the steam is superheated.

$$T(^{\circ}\text{C}) = \hat{v}(\text{m}^3/\text{kg})$$

$$\frac{212.4}{0.0996}$$

$$T = 0.1$$

$$225 \quad 0.1038$$

$$y = [(y_2 - y_1) \left(\frac{x - x_1}{x_2 - x_1} \right)] + y_1$$

$$T = (225 - 212.4) \left(\frac{0.1 - 0.0996}{0.1038 - 0.0996} \right) + 212.4$$

$$T = 213.6^{\circ}\text{C}$$

(b) $c \hat{v} = ?$

$$P_2 = \frac{100}{690} \left| \frac{100 \text{ kPa}}{1 \text{ bar}} \right| = 10.000 \text{ kPa} = 10 \text{ MPa}$$

$$\hat{v}_2 = ?$$

$$P_1 \hat{v}_1^{1.5} = P_2 \hat{v}_2^{1.5} \Rightarrow \hat{v}_2^{1.5} = \frac{P_1 \hat{v}_1^{1.5}}{P_2}$$

$$\hat{v}_2 = \left(\frac{P_1 \hat{v}_1^{1.5}}{P_2} \right)^{1/1.5}$$

$$\hat{v}_2 = (0.2 \times 0.1)^{1/1.5} \text{ m}^3/\text{kg}$$

$$= (0.2 \times 0.0316)^{1/1.5} = (0.2 \times 0.0316)^{0.667}$$

$$\hat{v}_2 = 0.0342 \text{ m}^3/\text{kg}$$

CHEN19111049

$$2625.3$$

$$225$$

$$u_1 = 2602.97 \text{ kJ/kg}$$

At state 2, $P_2 = 10 \text{ MPa}$, $\hat{v}_2 = 0.342 \text{ m}^3/\text{kg}$

$$\hat{u}_2 \text{ (kJ/kg)}$$

$$\hat{v}_2 \text{ (m}^3/\text{kg)}$$

$$\text{kg}$$

$$3045.8$$

$$0.8328$$

$$\hat{u}_2$$

$$0.0342$$

$$3144.5$$

$$0.0356$$

$$\hat{u}_2 = 3095.15 \text{ kJ/kg}$$

$$\Delta \hat{u} = \hat{u}_2 - \hat{u}_1$$

$$= (3095.15 - 2602.97) \text{ kJ/kg}$$

$$\Delta \hat{u} = 492.18 \text{ kJ/kg}$$

$$\Delta \hat{u} = \hat{q} + \hat{w}$$

$$\hat{q} = \Delta \hat{u} - \hat{w}$$

$$\hat{q} = (492.18 - 283.8) \text{ kJ/kg}$$

$$\hat{q} = 208.38 \text{ kJ/kg}$$