

a
Donor deviations, not raw level
high capacity
low capacity
Deviation from mean curve (log)
basal
AA
AA+G3
AA+G16.7
IBMX
wash
KCl
Fraction (1 min)
b
Fitted window correlation
c
Most variance is capacity
Penn. glucagon
Penn. insulin
Vand. glucagon
Vand. insulin
Stimulus-specific share of variance (%)
d
Stimulus structure is real
observed
null 95th centile
Likelihood ratio
Penn. diag vs capacity
Penn. full vs diag
Vand. diag vs capacity
Vand. full vs diag
e
Simulation retains more axes
Axes retained, conventional traits
real data (3)
A null
B indep.
C fitted
D pos. ctrl
f
Only axis 1 is reproduced
A null
B indep.
C fitted
D pos. ctrl
Congruence with real axes
equivalence, 0.85
Axis 1
Axis 2
Axis 3
g
Model recovers the truth
simulated truth
estimate
Glucose-suppression SD, model
h
Precision of each phenotype
Penn. Amino acids
Penn. Glucose supp.
Penn. Low glucose
Penn. IBMX
Penn. KCl
Vand. IBMX
Vand. KCl
Vand. Glucose supp.
Vand. Adrenaline
Reliability of the donor effect
i
Traits computed identically
Basal
G16.7 II
G16.7 AUC
Epi AUC
KCl AUC
KCl SI
Epi SI
IBMX AUC
IBMX SI
Spearman correlation with PanKbase