

a
Laboratory 1 glucagon
No diabetes
Type 2 diabetes
Type 1 diabetes
Glucagon release
Time (min)
b
Capacity and stimulus contrasts
Capacity
Amino acids
Glucose supp.
Low glucose
cAMP (IBMX)
KCl
0.87
0.51
0.33
0.77
0.71
reliability
Between-preparation SD (log units)
c
Laboratory 2 glucagon
No diabetes
Type 2 diabetes
Type 1 diabetes
Glucagon release
Time (min)
d
Suppression by group
Glucose suppression (log units)
No diabetes
Type 2 diabetes
e
Two contrasts, one random effect
r = -0.15
Glucose suppression (log units)
Capacity (log units)
f
Suppression by HbA1c tertile
Glucose suppression (log units)
≤ 36
36 to 44
> 44
HbA1c tertile (mmol/mol)
g
Both sexes
Glucose suppression (log units)
Male
Female
HbA1c (mmol/mol)
h
Weaker suppression with HbA1c
0.26 SD per 10 mmol/mol
95% CI 0.17, 0.34
p = 2.9e-09, censored fit
Glucose suppression (log units)
HbA1c (mmol/mol)
i
HbA1c associations, glucagon
Penn. Amino acids
Penn. Glucose supp.
Penn. KCl
Penn. Low glucose
Penn. cAMP (IBMX)
Penn. Capacity
Vand. Adrenaline
Vand. Glucose supp.
Vand. KCl
Vand. cAMP (IBMX)
Vand. Capacity
SD per 10 mmol/mol HbA1c (95% CI)