
a
Same donors, two laboratories
r = 0.33 (95% CI 0.09, 0.56)
p = 0.0045, n = 74
Laboratory 2, suppression
Laboratory 1, suppression
b
Only matched stimuli agree
cAMP / cAMP
KCl / KCl
cAMP / cAMP at 16.7 mM
capacity / capacity
suppression / suppression
suppression / KCl
cAMP / suppression
KCl / suppression
amino acids / suppression
amino acids / adrenaline
matched
mismatched
cross-stimulus
Correlation between laboratories
c
Disattenuation is uninformative
observed
disattenuated
r = 1
Bootstrap replicates
Correlation
d
Shared construct, different gain
Laboratory 1
Laboratory 2
fixed at 1
0.54 (SE 0.22), p = 0.013
Loading on the shared phenotype
e
Who the joint model uses
127 donors in the joint model
68
59
Donors
both laboratories
one laboratory
f
Three routes, one answer
Laboratory 1 alone
Laboratory 2 alone
Joint latent model
n = 112
n = 83
n = 127
SD per 10 mmol/mol HbA1c (95% CI)
g
Stable in laboratory 1 only
Laboratory 1
Laboratory 2
Stimulus-specific variance (%)
full
2 factors
1 factor
capacity only
h
Coordination is preserved
window slope, HbA1c
window slope, T2D
fraction correlation, HbA1c
fraction correlation, T2D
composite, HbA1c
composite, T2D
SD per 10 mmol/mol HbA1c or for T2D
i
Two measures of coordination
Spearman ρ = 0.31
n = 128
Fraction-correlation measure
Window-slope measure