
a
Protocol and observed OCR transitions
n = 89
median, IQR
45
75
108
135
165
basal
AAM
G3
G16
FCCP
OCR relative to basal
Time (min)
b
Response transitions, median and IQR
AAM
3 mM glucose
16.7 mM glucose
FCCP
azide, antimycin A
Observed response transition (min)
c
Duplicated traces, both excluded
HPAP-185 / HPAP-190
HPAP-106 / HPAP-108
HPAP-049 / HPAP-052
OCR + offset
Time (min)
d
No reliable non-mitochondrial floor
floor reached
still falling
median 0.74
33 of 87 still falling
2 with no inhibitor response
Post-inhibitor minimum / baseline
e
Spare capacity by group and sex
n=28
n=24
n=16
n=6
Spare capacity (log OCR)
F
M
control
T2D
f
Glucose response by group and sex
16.7 mM G response (log)
g
Chen et al. parameters, both feature sets
P
project Basal
published
project Maximal (see caption)
project Spare capacity
project High-glucose response
0.062
0.059
0.758
0.912
0.023
0.250
0.015
0.021
T2D × male interaction (SD of feature)
h
Oxygen does not predict glucagon
Female
Male
Glucagon suppression (log)
n = 59
Spare capacity (log OCR)
i
Attenuation after adjustment
3 mM G response
Spare capacity
16.7 mM glucose
16.7 mM G response
AAM response
Basal
3 mM glucose
Overall level
Maximal (FCCP)
Amino acids
Insulin response (same run)
Change in HbA1c coefficient (%)