# An EJS Simulation of a Mass Spectrometer

Student: Aaron Robinson

Advisors:

Dr. Martin DeWitt

Dr. Aaron Titus

March 26, 2011





# Ion Motion in EM trap

Newton's 2nd law of motion

$$F = ma$$
,

Lorentz force


$$F = q[E + (v \times B)],$$

Ion Acceleration

$$\frac{dv}{dt} = \frac{q}{m}[E + (v \times B)].$$

ODE Form: 
$$\frac{dv_x}{dt} = \frac{q}{m} [E_x + (v_y B_z - v_z B_y)]$$
$$\frac{dv_y}{dt} = \frac{q}{m} [E_y + (v_z B_x - v_x B_z)]$$
$$\frac{dv_z}{dt} = \frac{q}{m} [E_z + (v_x B_y - v_y B_x)]$$


### Ion trap schematic



http://www.chem.ucsb.edu/~devries/groupsite/labicr.htm


ODE Form: 
$$\frac{dv_x}{dt} = \frac{q}{m} [E_x + (v_y B_z - v_x B_y)]$$
$$\frac{dv_y}{dt} = \frac{q}{m} [E_y + (v_x B_z - v_x B_z)]$$
$$\frac{dv_z}{dt} = \frac{q}{m} [E_z + (v_x B_y - v_x B_z)]$$

### Ion trap schematic



http://www.chem.ucsb.edu/~devries/groupsite/labicr.htm

# Cyclotron motion



#### Cyclotron equation

$$\omega_c = \frac{qB_0}{m}$$

- Same m/q = Same frequency
- Independent of initial velocity



# Acknowledgment

- Anuj Shah
- Summer undergraduate research internship (SULI) program
- Pacific Northwest National Laboratory
- High Point University

#### References

 A. G. Marshall and F. R. Verdun, Fourier Transforms in NMR, Optical, and Mass Spectrometry. Elsevier Science, 1990.

