

Problem 26

Show that the Casoratian $W(n)$ given by:

$$W(n) = \begin{vmatrix} x_1(n) & x_2(n) & \cdots & x_k(n) \\ x_1(n+1) & x_2(n+1) & \cdots & x_k(n+1) \\ \vdots & & & \vdots \\ x_1(n+k-1) & x_2(n+k-1) & \cdots & x_k(n+k-1) \end{vmatrix} \quad (1)$$

may be given by the formula:

$$W(n) = \begin{vmatrix} x_1(n) & x_2(n) & \cdots & x_k(n) \\ \Delta x_1(n) & \Delta x_2(n) & \cdots & \Delta x_k(n) \\ \vdots & & & \vdots \\ \Delta^{k-1} x_1(n) & \Delta^{k-1} x_2(n) & \cdots & \Delta^{k-1} x_k(n) \end{vmatrix}$$

Attempt:

We know by definition that, $\Delta x_i(n) = x_i(n+1)$, and that $\Delta x_i(n+k-1) = x_i(n+k)$

Problem 27

Consider the second order difference equation

$$u(n+2) - \frac{n+3}{n+2}u(n+1) + \frac{2}{n+2}u(n) = 0$$

- (a) show that $u_1(n) = \frac{2^n}{n!}$ is a solution of the equation.
- (b) Use the above formula to find another solution $u_2(n)$ of the equation.

Attempt:

We know that $x^{(k)}$ is defined as follows:

$$x^{(k)} = x(x-1)(x-2)\cdots(x-k+1), \quad k \in \mathbb{Z}^+$$

$$(x+1)^{(k)} = (x+1)(x)(x-1)\cdots(x-k+2), \quad k \in \mathbb{Z}^+$$

This can be similarly defined as:

$$x^{(k)} = \frac{x!}{(x-k)!} = x(x-1)(x-2)\cdots(x-k+1) \quad (2)$$

and as:

$$x^{(k)} = \prod_{j=1}^k x - j + 1 = x(x-1)(x-2)\cdots(x-k+1) \quad (3)$$

$$(x+1)^{(k)} = \prod_{j=1}^{k-1} x - j + 1 = (x+1)(x)(x-2)\cdots(x-k+2)$$

and from the lecture notes that:

$$\begin{aligned} \Delta x^{(k)} &= (x+1)^{(k)} - x^{(k)} \\ &= (x+1)(x)(x-1)\cdots(x-k+2) - x(x-1)\cdots(x-k+2)(x-k+1) \\ &= (x)(x-1)\cdots(x-k+2)[x+1 - (x-k+1)] \\ &= x(x-1)\cdots(x-k+2)k \\ &= kx^{(k-1)} \end{aligned}$$

(i) If we express the above as $\Delta^1 x^{(k)}$, then we can do the following:

$$\begin{aligned}
 \Delta [\Delta x^{(k)}] &= \Delta [kx^{(k-1)}] \\
 &= k\Delta x^{(k-1)} \\
 &= k[(x+1)^{(k-1)} - x^{(k-1)}] \\
 &= k\left[(x+1)(x)(x-1)\cdots(x-(k-1)+2) \right. \\
 &\quad \left. - x(x-1)\cdots(x-(k-1)+2)(x-(k-1)+1)\right] \\
 &= k\left[(x)(x-1)\cdots(x-(k-1)+2)\right]\left((x+1) - (x-(k-1)+1)\right)
 \end{aligned}$$

The term on the left in square brackets is identically $x^{(k+2)}$ since:

$$x^{(k-2)} = x(x-1)\cdots(x-(k-2)+1) = x(x-1)\cdots(x-k+3)$$

and the term on the right simplifies to:

$$\left((x+1) - (x-(k-1)+1)\right) = k-1$$

giving us:

$$\Delta^2 x^{(k)} = k(k-1)x^{(k-2)}$$

Now let's assume that this pattern holds up to some $(n-1)$, then:

$$\Delta^{n-1} x^{(k)} = k(k-1)\cdots((k-(n-1)-1))x^{(k-(n-1))}$$

then:

$$\begin{aligned}
 \Delta [\Delta^{n-1} x^{(k)}] &= \Delta [k(k-1)\cdots(k-(n-1)-1)x^{(k-(n-1))}] \\
 &= \Delta [k(k-1)\cdots(k-n)x^{(k-n+1)}]
 \end{aligned}$$

We can express the quantity on the left inside the brackets as:

$$k(k-1)\cdots(k-n) = \frac{k!}{(k-n+1)!} \quad (4)$$

and since it is a scalar quantity that does not depend on x and is thus not effected by the Δ operator, we can move it outside. (since the Δ operator is linear)

$$\begin{aligned}
 \Delta [\Delta^{n-1} x^{(k)}] &= \frac{k!}{(k-n+1)!} \Delta x^{(k-n+1)} \\
 &= \frac{k!}{(k-n+1)!} [(x+1)^{(k-n+1)} - x^{(k-n+1)}]
 \end{aligned}$$

We know that the structure of the operator as before in $\Delta^2 x^{(k)}$, the bracketed part will have the form from (3), where $(x+1)^{(k)}$ shifts both indicies down by one:

$$\prod_{j=0}^{k-n} x - j + 1 - \prod_{j=1}^{k-n+1} x - j + 1$$

which simplifies to:

$$\left[\prod_{j=1}^{k-n} x - j + 1 \right] [(x+1) - (x - (k-n+1) + 1)] = \left[\prod_{j=1}^{k-n} x - j + 1 \right] (k-n+1)$$

going back to what we had, we can substitute into the bracketed part:

$$\begin{aligned} \Delta [\Delta^{n-1} x^{(k)}] &= \frac{k!}{(k-n-1)!} [(x+1)^{(k-n+1)} - x^{(k-n+1)}] \\ &= \frac{k!}{(k-n-1)!} \left[\prod_{j=1}^{k-n} x - j + 1 \right] (k-n+1) \end{aligned}$$

Simplifying further using (3) again we then get:

$$\Delta^n x^{(k)} = \frac{k!}{(k-n-1)!} x^{(k-n)} (k-n+1) \quad (5)$$

$$\Delta^n x^{(k)} = k(k-1) \cdots (k-n)(k-n+1) x^{(k-n)}$$

(ii) Using (5), if $n = k$:

$$\Delta^k x^{(k)} = \frac{k!}{(k-k-1)!} x^{(k-k)} (k-k+1)$$

and using (4)

$$\begin{aligned} \Delta^k x^{(k)} &= \frac{k!}{(-1)!} \frac{x!}{(x - (k-k))!} (0+1) \\ &= \frac{k!}{(-1)!} (1)(1) \\ &= \frac{k!}{(-1)!} \end{aligned}$$

The term in brackets is zero, since the indicies are backwards. For the term on the left, we know that negative interger factorials have the form:

$$n! = \frac{(-1)^{-n-1}}{(-n-1)!}$$

and that $-1! = 1$, so this gives us:

$$\Delta x^{(k)} = k!$$

Problem 28

Write the general solution of the difference equation:

$$(E - 3)^2(E^2 + 4)x(n) = 0$$

Attempt:

since $\Delta = E - I$,

$$\begin{aligned} \Delta \left[\frac{x(n)}{y(n)} \right] &= (E - I) \left[\frac{x(n)}{y(n)} \right] \\ &= E \left[\frac{x(n)}{y(n)} \right] - I \left[\frac{x(n)}{y(n)} \right] \\ &= \left[\frac{x(n+1)}{y(n+1)} \right] - \left[\frac{x(n)}{y(n)} \right] \\ &= \left[\frac{x(n+1)y(n) - x(n)y(n+1)}{y(n)y(n+1)} \right] \\ &= \left[\frac{x(n+1)y(n) + [y(n)x(n) - y(n)x(n)] - x(n)y(n+1)}{y(n)y(n+1)} \right] \\ &= \left[\frac{x(n+1)y(n) - y(n)x(n) - x(n)y(n+1) + y(n)x(n)}{y(n)y(n+1)} \right] \\ &= \left[\frac{y(n)[x(n+1) - x(n)] - x(n)[y(n+1) - y(n)]}{y(n)y(n+1)} \right] \\ &= \left[\frac{y(n)\Delta x(n) - x(n)\Delta y(n)}{y(n)Ey(n)} \right] \end{aligned}$$

Problem 29

Find a homogenous difference equation whose solution is:

$$y(n) = 2^{n-1} - 5^{n+1}$$

Attempt:

1. if $X_1(n)$ and $X_2(n)$ are solutions to ??, then:

$$X_1(k+n) + p_1(n)X_1(n+k-1) + \cdots + p_k(n)X_1(n) = 0$$

$$X_2(k+n) + p_1(n)X_2(n+k-1) + \cdots + p_k(n)X_2(n) = 0$$

So if we plug in $x(n) = X_1(n) + X_2(n)$ for (??)

$$\begin{aligned} x(n) &= X_1(k+n) + X_2(k+n) + p_1(n)X_1(n+k-1) + p_1(n)X_2(n+k-1) \\ &\quad + \cdots + p_k(n)X_1(n) + p_k(n)X_2(n) \\ &= X_1(k+n) + p_1(n)X_1(n+k-1) + \cdots + p_k(n)X_1(n) \\ &\quad + X_2(k+n) + p_1(n)X_2(n+k-1) + \cdots + p_k(n)X_2(n) \\ &= 0 + 0 \end{aligned}$$

But this is just equal to the two given solutions meaning the sum of each (when we split them) is zero, so this is also a solution.

2. Doing something similar:

$$\begin{aligned} \tilde{x}(n) &= aX_1(k+n) + p_1(n)aX_1(n+k-1) + \cdots + p_k(n)aX_1(n) \\ &= a(X_1(k+n) + p_1(n)X_1(n+k-1) + \cdots + p_k(n)X_1(n)) \\ &= a(0) \\ &= 0 \end{aligned}$$

So this is also a solution.

Problem 30

Find a partic

Attempt:(a) let $n = 1$

$$\begin{vmatrix} 1 & 1 & 3 \\ -2 & 2 & 3 \\ 4 & 4 & 3 \end{vmatrix} = 1 \begin{vmatrix} 2 & 3 \\ 4 & 3 \end{vmatrix} - (-2) \begin{vmatrix} 1 & 3 \\ 4 & 3 \end{vmatrix} + 4 \begin{vmatrix} 1 & 3 \\ 2 & 3 \end{vmatrix} \\ = 6 - 12 + 6 - 24 + 12 - 24 = -12 - 6 - 18 = -36$$

since this $\neq 0$ these are linearly independent

(b)

$$\begin{vmatrix} 0 & 1 & 3 \\ 0 & 2 & 3 \\ 0 & 4 & 3 \end{vmatrix} = 0 \begin{vmatrix} 2 & 3 \\ 4 & 3 \end{vmatrix} - 0 \begin{vmatrix} 1 & 3 \\ 4 & 3 \end{vmatrix} + 0 \begin{vmatrix} 1 & 3 \\ 2 & 3 \end{vmatrix} = 0$$

So this is linearly dependent.

(c)

$$\begin{vmatrix} 1 & 3 & 4 \\ 2 & 3 & 8 \\ 4 & 3 & 16 \end{vmatrix} = 1 \begin{vmatrix} 3 & 8 \\ 3 & 16 \end{vmatrix} - 2 \begin{vmatrix} 3 & 4 \\ 3 & 16 \end{vmatrix} + 4 \begin{vmatrix} 3 & 4 \\ 3 & 8 \end{vmatrix} \\ = 48 - 24 + -96 + 24 + 96 - 48 = 0$$

So this is linearly dependent