

**Third Semester B. E. (Computer Science and Engineering /
Information Technology) Examination**

LINEAR ALGEBRA AND STATISTICS

Time : 3 Hours]

[Max. Marks : 60

Instructions to Candidates :—

- (1) All questions are compulsory.
- (2) Use of Non – programmable calculator is permitted.
- (3) Use of area under normal curve table and student's t distribution table is permitted.

1. Solve any Three :—

- (a) Find $F(x, y)$, where the linear map $F: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is defined by $F(1, 2) = (3, -1)$ and $F(0, 1) = (2, 1)$.
5 (CO 1, 2)
- (b) Prove that kernel of a linear mapping is a subspace of domain of linear mapping.
5 (CO 1, 2)
- (c) Find the basis as well as dimension of the kernel and the image of linear map $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$ defined by :
 $T(x, y, z, t) = (x + 2y + 3z + 2t, 2x + 7z + 5t, x + 2y + 6z + 5t)$.
5 (CO 1, 2)
- (d) Let $F: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be defined by $F(x, y, z) = (2y + z, x - 4y, 3x)$. Find the matrix representing F relative to the bases
 $S = \{(1, 1, 1), (1, 1, 0), (1, 0, 0)\}$
5 (CO 1, 2)

2. Solve any Three :—

- (a) Let S consist of the following vectors in \mathbb{R}^4 :
 $u_1 = (1, 2, 1, 3), u_2 = (16, -13, 1, 3), u_3 = (1, 1, -9, 2), u_4 = (1, 1, 0, -1)$
 - (i) Determine whether S is orthogonal or not.
 - (ii) Find basis of \mathbb{R}^4 .
5 (CO 1, 2)

- (b) Define inner product space.
Verify that the following defines an inner product in \mathbb{R}^2 :
 $\langle u, v \rangle = x_1y_1 - x_1y_2 - x_2y_1 + 3x_2y_2$, where $u = (x_1, x_2)$, $v = (y_1, y_2)$.
 5 (CO 1, 2)
- (c) Suppose $v = (1, 3, 5, 7)$. Find the projection of v onto W or, in other words, find $w \in W$ that minimizes $\|v - w\|$, where W is the subspace of \mathbb{R}^4 spanned by : $u_1 = (1, 1, 1, 1)$ and $u_2 = (1, -3, 4, -2)$.
 5 (CO 1, 2)
- (d) Find the singular value decomposition of matrix $A = \begin{bmatrix} 5 & 5 \\ -1 & 7 \end{bmatrix}$.
 5 (CO 1, 2)

3. Solve :—

- (a) The joint probability density function of the two continuous random variables X and Y is given by :

$$f(x, y) = \begin{cases} k x^3 y^3 & , \quad 0 \leq x \leq 2, \quad 0 \leq y \leq 2 \\ 0 & , \quad \text{elsewhere} \end{cases}$$

 Find k and the marginal densities of X and Y .
 5 (CO 3)
- (b) A salesman S sells in only three cities A , B and C . Suppose S never sells in the same city on successive days. If S sells in city A , then the next day S sells in city B . However, if S sells in either B or C , then the next day S is twice as likely to sell in city A as in the other city. Find out how often, in the long run, S sells in each city.
 5 (CO 3)
- (c) Explain the Poisson process.
 5 (CO 3)

4. Solve any **Three** :—

- (a) A certain stimulus administered to each of 12 patients resulted in the following change in blood pressure :
 5, 2, 8, -1, 3, 0, -2, 1, 5, 0, 4, 6.
 Can it be concluded that the stimulus will increase the blood pressure ?
 5 (CO 3)

- (b) Twenty people were attacked by a disease and only 18 survived. Will you reject the hypothesis that the survival rate, if attacked by this disease, is 85% in favor of the hypothesis that it is more, at 5% level ?
(Use Large Sample Test) 5 (CO 3)
- (c) A cigarette manufacturing firm claims that its brand A of the cigarettes outsells its brand B by 8%. If it is found that 42 out of a sample of 100 smokers prefer brand A and 18 out of another random sample of 100 smokers prefer brand B, test whether the 8% difference is a valid claim.
(Use 5% level of significance). 5 (CO 3)
- (d) In a certain factory there are two independent processes manufacturing the same item. The average weight in a sample of 250 items produced from one process is found to be 120 with a standard deviation of 12, while the corresponding figures in a sample of 400 items from the other process are 124 and 14. Obtain the standard error of difference between the two sample means : Is this difference significant ? Also find the 99% confidence limits for the difference in the average weights of items produced by the two processes respectively. 5 (CO 3)