

How Common is Autism?

Autism statistics from the U.S. Centers for Disease Control and Prevention (CDC) released in March 2014 identify around 1 in 68 American children as on the autism spectrum – a ten-fold increase in prevalence in 40 years. Careful research shows that this increase is only partly explained by improved diagnosis and awareness. Studies also show that autism is four to five times more common among boys than girls. An estimated 1 out of 42 boys and 1 in 189 girls are diagnosed with autism in the United States.

ASD affects over 2 million individuals in the U.S. and tens of millions worldwide. Moreover, government autism statistics suggest that prevalence rates have increased 10% to 17% annually in recent years. There is no established explanation for this continuing increase, although improved diagnosis and environmental influences are two reasons often considered.

What Causes Autism?

Not long ago, the answer to this question would have been “we have no idea.” Research is now delivering the answers. First and foremost, we now know that there is no one cause of autism, just as there is no one type of autism. Over the last five years, scientists have identified a number of rare gene changes or mutations associated with autism. Research has identified more than 100 autism risk genes. In around 15% of cases, a specific genetic cause of a person’s autism can be identified. However, most cases involve a complex and variable combination of genetic risk and environmental factors that influence early brain development.



In other words, in the presence of a genetic pre-disposition to autism, a number of non-genetic or environmental influences further increase a child’s risk. The clearest evidence of these environmental risk factors involves events before and during birth. They include advanced parental age at time of conception (both mom and dad), maternal illness during pregnancy, extreme prematurity, very low birth weight and certain difficulties during birth, particularly those involving periods of oxygen deprivation to the baby’s brain. Mothers exposed to high levels of pesticides and air pollution may also be at higher risk of having a child with ASD. It is important to keep in mind that these factors, by themselves, do not cause autism. Rather, in combination with genetic risk factors, they appear to modestly increase risk.

A small but growing body of research suggests that autism risk is lower among children whose mothers took prenatal vitamins (containing folic acid) in the months before and after conception.

Increasingly, researchers are looking at the role of the immune system in autism. Autism Speaks is working to increase awareness and investigation of these and other issues where further research has the potential to improve the lives of those who struggle with autism.

While the causes of autism are complex, it is abundantly clear that it is not caused by bad parenting. Dr. Leo Kanner, the psychiatrist who first described autism as a unique condition in 1943, believed that it was caused by cold, unloving mothers. Bruno Bettelheim, a renowned professor of child development, perpetuated this misinterpretation of autism. Their promotion of the idea that unloving mothers caused their children’s autism created a generation of parents who carried the tremendous burden of guilt for their child’s disability. In the 1960s and 70s, Dr. Bernard Rimland, the father of a son with autism who later founded the Autism Society of America and the Autism Research Institute, helped the medical community understand that autism is a biological disorder and is not caused by cold parents.