

4.13.72

AI25BTECH11003 - Bhavesh Gaikwad

Question: A non-zero vector \mathbf{a} is parallel to the line of intersection of the plane determined by the vectors $\hat{i}, \hat{i} + \hat{j}$ and the plane determined by the vectors $\hat{i} - \hat{j}, \hat{i} + \hat{k}$. The angle between \mathbf{a} and the vector $\hat{i} - 2\hat{j} + 2\hat{k}$ is _____.

(1996)

Solution:

$$\text{Let } \mathbf{A} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \mathbf{D} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

Let \mathbf{n}_1 be the perpendicular vector to Plane-1.

$$\text{Let } \mathbf{n}_1 = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

$$\therefore \mathbf{A}^\top \mathbf{n}_1 = 0 \text{ \& } \mathbf{B}^\top \mathbf{n}_1 = 0 \quad (0.1)$$

From Equation 0.1,

$$x_1 = 0 \text{ \& } x_2 + x_1 = 0 \Rightarrow x_1 = x_2 = 0 \quad (0.2)$$

$$\mathbf{n}_1 = \begin{pmatrix} 0 \\ 0 \\ x_3 \end{pmatrix} \quad \text{OR} \quad \mathbf{n}_1 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad (0.3)$$

Let \mathbf{n}_2 be the perpendicular vector to Plane-2.

$$\text{Let } \mathbf{n}_2 = \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix}$$

$$\therefore \mathbf{C}^\top \mathbf{n}_2 = 0 \text{ \& } \mathbf{D}^\top \mathbf{n}_2 = 0 \quad (0.4)$$

From Equation 0.4,

$$z_1 - z_2 = 0 \text{ \& } z_1 + z_3 = 0 \quad (0.5)$$

$$\mathbf{n}_2 = \begin{pmatrix} -z_3 \\ -z_3 \\ z_3 \end{pmatrix} \quad \text{OR} \quad \mathbf{n}_2 = \begin{pmatrix} -1 \\ -1 \\ 1 \end{pmatrix} \quad (0.6)$$

Let \mathbf{n}_3 be the parallel vector to the line of intersection of planes.

$$\text{Let } \mathbf{n}_3 = \begin{pmatrix} k_1 \\ k_2 \\ k_3 \end{pmatrix}$$

Since, \mathbf{n}_3 is perpendicular to both \mathbf{n}_1 and \mathbf{n}_2 .

$$\therefore \mathbf{n}_3^\top \mathbf{n}_1 = 0 \text{ \& } \mathbf{n}_3^\top \mathbf{n}_2 = 0 \quad (0.7)$$

The angle between \mathbf{a} and $1\hat{i} - 2\hat{j} + 2\hat{k}$ is 45° .

(0.8)

$$k_3 = 0 \text{ \& } k_3 - k_1 - k_2 = 0 \quad \Rightarrow k_2 = -k_1 \quad (0.9)$$

$$\mathbf{n}_3 = \begin{pmatrix} k_1 \\ k_2 \\ 0 \end{pmatrix} \quad OR \quad \mathbf{n}_3 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad (0.10)$$

From Equation 0.10,

$$\therefore \mathbf{a} = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} \quad (0.11)$$

Let $\mathbf{u} = \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}$ [Already given in the Question]

We know,

$$\mathbf{a}^\top \mathbf{u} = \|\mathbf{a}\| \|\mathbf{u}\| \cos(\theta) \quad (0.12)$$

$$\|\mathbf{a}\| = \sqrt{\mathbf{a}^\top \mathbf{a}} = \sqrt{2}, \|\mathbf{u}\| = \sqrt{\mathbf{u}^\top \mathbf{u}} = 3 \quad (0.13)$$

From Equation 0.12 and 0.13,

$$\cos(\theta) = \frac{3}{3\sqrt{2}} \quad \Rightarrow \theta = 45^\circ \quad (0.14)$$

The angle between \mathbf{a} and $1\hat{i} - 2\hat{j} + 2\hat{k}$ is 45° .

(0.15)




Fig. 0.1: Plane