8.2.33

EE25BTECH11065-Yoshita J

September 19,2025

Question

Find the equation of the conic with length of major axis 26, foci $(\pm 5,0)$.

Theoretical Solution

The equation of a conic is

$$x^T V x + 2u^T x + f = 0 (1)$$

where

$$V = ||n||^2 I - e^2 n n^T$$
 (2)

The foci are

$$F_1 = \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \quad F_2 = \begin{pmatrix} -5 \\ 0 \end{pmatrix}$$
 (3)

The centre is

$$u = \frac{F_1 + F_2}{2} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \tag{4}$$

The axis vector is

$$n = F_1 - F_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \tag{5}$$

Theoretical Solution

Therefore, substituting $n = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ in (2), we get

$$V = \begin{pmatrix} 1 - e^2 & 0 \\ 0 & 1 \end{pmatrix} \tag{6}$$

From the formula for the length of the major axis,

$$2\sqrt{\frac{|f|}{\lambda_1}}\tag{7}$$

where $\lambda_1 = 1 - e^2$. Hence

$$26 = 2\sqrt{\frac{|f|}{1 - e^2}} \tag{8}$$

The relation between focus and eccentricity is

$$\pm ce^2 = 5 \tag{9}$$

Theoretical Solution

The distance *c* is

$$c = \pm \frac{1}{e} \sqrt{\frac{|f|}{|e^2 - 1|}} \tag{10}$$

Thus from (8)–(10), solving the unknowns (c, e, f) we get

$$e = \frac{5}{13}, \quad c = \pm 5, \quad |f| = 144.$$
 (11)

Let $x = \begin{pmatrix} 0 \\ \alpha \end{pmatrix}$ be a vertex on the minor axis. Substituting in (1):

$$\frac{12^2}{1} + f = 0 \implies f = -144. \tag{12}$$

Hence the equation of the conic is

$$x^{T} \begin{pmatrix} \frac{144}{169} & 0\\ 0 & 1 \end{pmatrix} x - 144 = 0.$$
 (13)

C Code

```
#include <stdio.h>
#include <math.h>
#define PI 3.1415926535
double calculate circular sector area() {
   double radius = 2.0;
   double angle in radians = PI / 6.0;
   double area = 0.5 * radius * radius * angle in radians;
   return area;
```

Python Code

```
import numpy as np
 import matplotlib.pyplot as plt
 a = 13
 b = 12
 c = 5
theta = np.linspace(0, 2 * np.pi, 200)
 x = a * np.cos(theta)
y = b * np.sin(theta)
plt.figure(figsize=(10, 8))
ax = plt.gca()
 ax.plot(x, y, label='Ellipse: x^2/169 + y^2/144 = 1')
 ax.plot(0, 0, 'ko', label='Center (0, 0)')
 lax.plot(c, 0, 'ro', label='Focus 1 (5, 0)')
 ax.plot(-c, 0, 'ro', label='Focus 2 (-5, 0)')
    EE25BTECH11065-Yoshita J
                                  8.2.33
                                                     September 19,2025
```

Python Code

```
ax.set title('Plot of the Ellipse', fontsize=16)
ax.set xlabel('X-axis')
ax.set ylabel('Y-axis')
ax.set aspect('equal', adjustable='box')
ax.grid(True, linestyle='--')
ax.legend()
ax.set xlim(-a - 2, a + 2)
ax.set_ylim(-b - 2, b + 2)
ax.axhline(0, color='black', linewidth=0.5)
ax.axvline(0, color='black', linewidth=0.5)
plt.show()
```

